

Chapter 1

Equations and Inequalities

Section 1.1 Linear Equations

Classroom Example 1 (page 89)

$$-4(3x - 5) = 3 - (8x + 7)$$

$$-12x + 20 = 3 - 8x - 7$$

$$-12x + 20 = -4 - 8x$$

$$24 = 4x$$

$$6 = x$$

Solution set: $\{6\}$

Classroom Example 2 (page 89)

$$\frac{3x+6}{10} - \frac{1}{2}x = \frac{2}{5}x + \frac{33}{5}$$

$$10\left(\frac{3x+6}{10} - \frac{1}{2}x\right) = 10\left(\frac{2}{5}x + \frac{33}{5}\right)$$

$$3x + 6 - 5x = 4x + 66$$

$$-2x + 6 = 4x + 66$$

$$-6x = 60$$

$$x = -10$$

Solution set: $\{-10\}$

Classroom Example 3 (page 90)

$$(a) \quad 6x - 9 = 4x + 13$$

$$2x = 22 \Rightarrow x = 11$$

conditional equations; $\{11\}$

$$(b) \quad 10 + 14x = 7(2x - 5)$$

$$10 + 14x = 14x - 35$$

$$10 = -35$$

contradiction; \emptyset

$$(c) \quad -3(2x - 1) + 5x = 3 - x$$

$$-6x + 3 + 5x = 3 - x$$

$$-x + 3 = 3 - x$$

identity; $\{\text{all real numbers}\}$

Classroom Example 4 (page 91)

$$(a) \quad d = rt \Rightarrow \frac{d}{r} = t$$

$$(b) \quad S = kr^2 + kr\ell$$

$$S = k(r^2 + r\ell)$$

$$\frac{S}{r^2 + r\ell} = k$$

$$(c) \quad 11y + 8 = 2(4y + 5w) - 6z$$

$$11y + 8 = 8y + 10w - 6z$$

$$3y = 10w - 6z - 8$$

$$y = \frac{10w - 6z - 8}{3}$$

Classroom Example 5 (page 92)

$$P = \$2580, t = \frac{9}{12} = \frac{3}{4} \text{ yr, and } r = 0.06$$

$$I = Prt = 2580(0.06)\left(\frac{3}{4}\right) = \$116.10$$

Section 1.2 Applications and Modeling with Linear Equations

Classroom Example 1 (page 95)

Let x = the length of the original rectangle. Then,

$x - 2$ = the width of the original rectangle.

$x + 3$ = the length of the new rectangle, so

$(x - 2) + 3 = x + 1$ = the width of the new rectangle.

The perimeter of the new rectangle is

$$2(x + 3) + 2(x + 1) = 2x + 6 + 2x + 2 = 4x + 8. \text{ The}$$

perimeter of the new rectangle is 4 in. less than 8 times the width of the original rectangle, so we have

$$4x + 8 = 8(x - 2) - 4. \text{ Solving, we have}$$

$$4x + 8 = 8(x - 2) - 4 \Rightarrow 4x + 8 = 8x - 16 - 4 \Rightarrow$$

$$4x + 8 = 8x - 20 \Rightarrow 28 = 4x \Rightarrow 7 = x$$

Thus, the length of the original rectangle is 7 in. and the width of the original rectangle is 5 in.

Classroom Example 2 (page 96)

Let x = the distance to her grandmother's house.

	r	t	d
Going	40	$\frac{x}{40}$	x
Returning	48	$\frac{x}{48}$	x

The driving time returning was 1 hour less than the driving time going, so we have

$$\frac{x}{40} - 1 = \frac{x}{48}. \text{ Solving, we have}$$

$$\frac{x}{40} - 1 = \frac{x}{48}$$

$$40 \cdot 48 \left(\frac{x}{40} - 1 \right) = 40 \cdot 48 \left(\frac{x}{48} \right)$$

$$48x - 1920 = 40x$$

$$-1920 = -8x \Rightarrow 240 = x$$

The distance from Krissa's home to her grandmother's home is 240 mi.

Classroom Example 3 (page 97)

Let x = the amount of 25% antifreeze solution (in liters).

Strength	Gallons of Solution	Gallons of Pure Antifreeze
25%	x	$0.25x$
10%	5	$0.10 \cdot 5 = 0.5$
15%	$x + 5$	$0.15(x + 5)$

The number of gallons of pure antifreeze in the 25% solution plus the number of gallons of pure antifreeze in the 10% solution must equal the number of gallons of pure antifreeze in the 15% solution.

$$0.25x + 0.5 = 0.15(x + 5)$$

$$0.25x + 0.5 = 0.15x + 0.75$$

$$0.1x = 0.25 \Rightarrow x = 2.5 \text{ L}$$

2.5 liters of the 25% solution should be added.

Classroom Example 4 (page 98)

Let x = amount invested at 2.4%.

Then $28,000 - x$ = amount invested at 3.1%.

Amount in Account	Interest Rate	Interest
x	2.4%	$0.024x$
$28,000 - x$	3.1%	$0.031(28,000 - x)$
		784

The amount of interest from the 2.4% account plus the amount of interest from the 3.1% account must equal the total amount of interest.

$$0.024x + 0.031(28,000 - x) = 784$$

$$0.024x + 868 - 0.031x = 784$$

$$-0.007x = -84$$

$$x = 12,000$$

Owen invested \$12,000 at 2.4% and $\$28,000 - \$12,000 = \$16,000$ at 3.1%.

Classroom Example 5 (page 99)

$$P = 1.06F + 7.18$$

$$P = 70, \text{ so we have}$$

$$70 = 1.06F + 7.18$$

$$62.82 = 1.06F \Rightarrow 59.26 \approx F$$

The flow rate must be approximately 59.26 L per second to remove 70% of the contaminants.

Classroom Example 6 (page 99)

- (a) The year 2007 is 7 years after the year 2000.

Let $x = 7$ and find the value of y .

$$y = 331x + 5091$$

$$y = 331(7) + 5091 = 7408$$

The per capita health care expenditures in the year 2007 were \$7408.

- (b) Let $y = 9100$ and find the value of x

$$y = 331x + 5091$$

$$9100 = 331x + 5091$$

$$4009 = 331x$$

$$x \approx 12.1$$

Per capita health care expenditures are projected to reach \$9100 about 12.1 years after 2000, or in 2012.

Section 1.3 Complex Numbers**Classroom Example 1 (page 106)**

(a) $\sqrt{-81} = i\sqrt{81} = 9i$

(b) $\sqrt{-55} = i\sqrt{55}$

(c) $\sqrt{-98} = i\sqrt{98} = i\sqrt{49 \cdot 2} = 7i\sqrt{2}$

Classroom Example 2 (page 107)

(a) $\sqrt{-21} \cdot \sqrt{-21} = i\sqrt{21} \cdot i\sqrt{21} = i^2 (\sqrt{21})^2 = -21$

(b) $\sqrt{-5} \cdot \sqrt{-30} = i\sqrt{5} \cdot i\sqrt{30} = i^2 \sqrt{150}$
 $= i^2 \sqrt{25 \cdot 6} = -5\sqrt{6}$

(c) $\frac{\sqrt{-42}}{\sqrt{-3}} = \frac{i\sqrt{42}}{i\sqrt{3}} = \sqrt{\frac{42}{3}} = \sqrt{14}$

(d) $\frac{\sqrt{-63}}{\sqrt{21}} = \frac{i\sqrt{63}}{\sqrt{21}} = i\sqrt{\frac{63}{21}} = i\sqrt{3}$

Classroom Example 3 (page 107)

$$\begin{aligned}\frac{15 - \sqrt{-75}}{5} &= \frac{15 - i\sqrt{75}}{5} = \frac{15 - i\sqrt{25 \cdot 3}}{5} \\ &= \frac{15 - 5i\sqrt{3}}{5} = \frac{5(3 - i\sqrt{3})}{5} = 3 - i\sqrt{3}\end{aligned}$$

Classroom Example 4 (page 108)

$$(a) \quad (4 - 5i) + (-5 + 8i) = -1 + 3i$$

$$(b) \quad (-10 + 7i) - (5 - 3i) = -15 + 10i$$

Classroom Example 5 (page 109)

$$\begin{aligned}(a) \quad (5 + 3i)(2 - 7i) &= 10 + 5(-7i) + (3i)(2) + (3i)(-7i) \\ &= 10 - 35i + 6i - 21i^2 \\ &= 10 - 29i + 21 = 31 - 29i\end{aligned}$$

$$\begin{aligned}(b) \quad (4 - 5i)^2 &= 4^2 - 2(4)(5i) + (5i)^2 \\ &= 16 - 40i + 25i^2 \\ &= 16 - 40i - 25 = -9 - 40i\end{aligned}$$

$$\begin{aligned}(c) \quad (9 - 8i)(9 + 8i) &= 9^2 - (8i)^2 \quad \text{difference of two squares} \\ &= 81 - 64i^2 \\ &= 81 - 64(-1) \\ &= 81 + 64 = 145, \text{ or } 145 + 0i\end{aligned}$$

Classroom Example 6 (page 110)

$$\begin{aligned}(a) \quad \frac{5 - 5i}{3 + i} &= \frac{5 - 5i}{3 + i} \cdot \frac{3 - i}{3 - i} = \frac{15 - 5i - 15i + 5i^2}{9 - i^2} \\ &= \frac{15 - 20i - 5}{9 - (-1)} = \frac{10 - 20i}{10} = 1 - 2i\end{aligned}$$

$$(b) \quad \frac{15}{-i} = \frac{15}{-i} \cdot \frac{i}{i} = \frac{15i}{-i^2} = \frac{15i}{1} = 15i, \text{ or } 0 + 15i$$

Classroom Example 7 (page 111)

$$(a) \quad i^{33} = i^{32} \cdot i = (i^4)^8 \cdot i = 1^8 \cdot i = i$$

$$\begin{aligned}(b) \quad i^{-14} &= i^{-16} \cdot i^2 = (i^{16})^{-1} \cdot i^2 = \left((i^4)^4\right)^{-1} \cdot i^2 \\ &= 1^{-1} \cdot i^2 = i^2 = -1\end{aligned}$$

Section 1.4 Quadratic Equations**Classroom Example 1 (page 114)**

$$\begin{aligned}10x^2 + x - 2 &= 0 \\ (2x + 1)(5x - 2) &= 0 \\ 2x + 1 = 0 \quad \text{or} \quad 5x - 2 &= 0 \\ 2x = -1 \quad \text{or} \quad 5x = 2 \\ x = -\frac{1}{2} \quad \text{or} \quad x = \frac{2}{5}\end{aligned}$$

$$\text{Solution set: } \left\{-\frac{1}{2}, \frac{2}{5}\right\}$$

Classroom Example 2 (page 115)

$$(a) \quad x^2 = 29 \Rightarrow x = \pm\sqrt{29}$$

$$(b) \quad x^2 = -144 \Rightarrow x = \pm 12i$$

$$\begin{aligned}(c) \quad (x - 8)^2 &= 24 \Rightarrow x - 8 = \pm\sqrt{24} \Rightarrow \\ x - 8 &= \pm 2\sqrt{6} \Rightarrow x = 8 \pm 2\sqrt{6}\end{aligned}$$

Classroom Example 3 (page 116)

$$\begin{aligned}x^2 + 10x - 20 &= 0 \\ x^2 + 10x &= 20 \\ x^2 + 10x + \left(\frac{1}{2} \cdot 10\right)^2 &= 20 + \left(\frac{1}{2} \cdot 10\right)^2 \\ x^2 + 10x + 25 &= 20 + 25 \\ (x + 5)^2 &= 45 \\ x + 5 &= \pm\sqrt{45} = \pm 3\sqrt{5} \\ x &= -5 \pm 3\sqrt{5}\end{aligned}$$

Classroom Example 4 (page 116)

$$\begin{aligned}4x^2 + 6x + 5 &= 0 \\ x^2 + \frac{6}{4}x + \frac{5}{4} &= 0 \\ x^2 + \frac{6}{4}x &= -\frac{5}{4} \\ x^2 + \frac{6}{4}x + \left(\frac{1}{2} \cdot \frac{6}{4}\right)^2 &= -\frac{5}{4} + \left(\frac{1}{2} \cdot \frac{6}{4}\right)^2 \\ x^2 + \frac{3}{2}x + \frac{9}{16} &= -\frac{5}{4} + \frac{9}{16} \\ \left(x + \frac{3}{4}\right)^2 &= -\frac{11}{16} \\ x + \frac{3}{4} &= \pm i\sqrt{\frac{11}{16}} = \pm \frac{\sqrt{11}}{4}i \\ x &= -\frac{3}{4} \pm \frac{\sqrt{11}}{4}i\end{aligned}$$

Classroom Example 5 (page 117)

$$x^2 + 6x = 3 \Rightarrow x^2 + 6x - 3 = 0$$

$$a = 1, b = 6, c = -3$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-6 \pm \sqrt{6^2 - 4(1)(-3)}}{2(1)} \\ &= \frac{-6 \pm \sqrt{36 + 12}}{2} = \frac{-6 \pm \sqrt{48}}{2} \\ &= \frac{-6 \pm 4\sqrt{3}}{2} = -3 \pm 2\sqrt{3} \end{aligned}$$

Classroom Example 6 (page 118)

$$4x^2 = 3x - 5 \Rightarrow 4x^2 - 3x + 5 = 0$$

$$a = 4, b = -3, c = 5$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(4)(5)}}{2(4)} \\ &= \frac{3 \pm \sqrt{9 - 80}}{8} = \frac{3 \pm \sqrt{-71}}{8} = \frac{3 \pm i\sqrt{71}}{8} \\ &= \frac{3}{8} \pm \frac{\sqrt{71}}{8}i \end{aligned}$$

Classroom Example 7 (page 118)

$$x^3 - 125 = (x - 5)(x^2 + 5x + 25) \quad \begin{array}{l} \text{difference of cubes} \\ \text{(section R.4)} \end{array}$$

$$x - 5 = 0 \quad \text{or} \quad x^2 + 5x + 25 = 0$$

$$x = 5 \quad \text{or} \quad x = \frac{-5 \pm \sqrt{5^2 - 4(1)(25)}}{2(1)}$$

$$x = \frac{-5 \pm \sqrt{25 - 100}}{2}$$

$$x = \frac{-5 \pm \sqrt{-75}}{2}$$

$$x = \frac{-5 \pm 5\sqrt{3}i}{2}$$

$$x = -\frac{5}{2} \pm \frac{5\sqrt{3}}{2}i$$

$$\text{Solution set: } \left\{ 5, -\frac{5}{2} \pm \frac{5\sqrt{3}}{2}i \right\}$$

Classroom Example 8 (page 119)

$$\begin{aligned} \text{(a)} \quad V &= \frac{1}{3}\pi r^2 h \Rightarrow 3V = \pi r^2 h \Rightarrow \frac{3V}{\pi h} = r^2 \Rightarrow \\ r &= \pm \sqrt{\frac{3V}{\pi h}} \Rightarrow r = \pm \sqrt{\frac{3V}{\pi h}} \cdot \frac{\sqrt{\pi h}}{\sqrt{\pi h}} \Rightarrow \\ r &= \frac{\pm \sqrt{3V\pi h}}{\pi h} \end{aligned}$$

$$\text{(b)} \quad 2my^2 - ny = 3p \Rightarrow 2my^2 - ny - 3p = 0$$

Use the quadratic formula with $a = 2m$, $b = -n$, and $c = -3p$ to solve for y :

$$\begin{aligned} y &= \frac{-(-n) \pm \sqrt{(-n)^2 - 4(2m)(-3p)}}{2(2m)} \\ &= \frac{n \pm \sqrt{n^2 + 24mp}}{4m} \end{aligned}$$

Classroom Example 9 (page 120)

$$\text{(a)} \quad 4x^2 - 12x + 9 = 0$$

$$a = 4, b = -12, c = 9$$

$$b^2 - 4ac = (-12)^2 - 4(4)(9) = 0$$

Therefore, there is one distinct rational solution.

$$\text{(b)} \quad 3x^2 + x = -5 \Rightarrow 3x^2 + x + 5 = 0$$

$$a = 3, b = 1, c = 5$$

$$b^2 - 4ac = 1^2 - 4(3)(5) = -59$$

Therefore, there are two distinct nonreal complex solutions.

$$\text{(c)} \quad 2x^2 = 6x + 7 \Rightarrow 2x^2 - 6x - 7 = 0$$


$$a = 2, b = -6, c = -7$$

$$b^2 - 4ac = (-6)^2 - 4(2)(-7) = 92$$

Therefore, there are two distinct irrational solutions.

Section 1.5 Applications and Modeling with Quadratic Equations**Classroom Example 1 (page 124)**

Let x = the width of the rectangle. Then $2x$ = the length of the rectangle. The box is formed by cutting 10 cm + 10 cm from both the length and the width. Thus the width of the bottom of the box is $x - 20$, the length of the bottom of the box is $2x - 20$, and the height of the box is 10 cm.



(continued on next page)

(continued)

$$\begin{aligned}
 V &= lwh \\
 7500 &= (2x - 20)(x - 20)(10) \\
 7500 &= 20x^2 - 600x + 4000 \\
 0 &= 20x^2 - 600x - 3500 \\
 0 &= x^2 - 30x - 175 \\
 0 &= (x + 5)(x - 35) \\
 x + 5 &= 0 \quad \text{or} \quad x - 35 = 0 \\
 x &= -5 \quad \text{or} \quad x = 35
 \end{aligned}$$

Reject the negative solution because length cannot be negative. The dimensions of the box are 35 cm by 70 cm.

Classroom Example 2 (page 125)

Let x = the length of the shorter leg.
 Then $2x - 10$ = the length of the longer leg, and
 $(2x - 10) + 20 = 2x + 10$ = the length of the hypotenuse. Using the Pythagorean theorem, we have

$$\begin{aligned}
 x^2 + (2x - 10)^2 &= (2x + 10)^2 \\
 x^2 + 4x^2 - 40x + 100 &= 4x^2 + 40x + 100 \\
 5x^2 - 40x + 100 &= 4x^2 + 40x + 100 \\
 x^2 - 80x &= 0 \\
 x(x - 80) &= 0
 \end{aligned}$$

$x = 0$ or $x - 80 = 0 \Rightarrow x = 80$. Since length cannot be 0, we reject that solution. The shorter leg is 80 m, the longer leg is $2(80) - 10 = 150$ m, and the hypotenuse is $2(80) + 10 = 170$ m.

Classroom Example 3 (page 126)

$$\begin{aligned}
 \text{(a)} \quad s &= -4.9t^2 + 73.5t \\
 100 &= -4.9t^2 + 73.5t \Rightarrow \\
 4.9t^2 - 73.5t + 100 &= 0 \\
 \text{Solve using the quadratic formula, with} \\
 a &= 4.9, b = -73.5, \text{ and } c = 100. \\
 t &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-73.5) \pm \sqrt{(-73.5)^2 - 4(4.9)(100)}}{2(4.9)} \\
 &= \frac{73.5 \pm \sqrt{5402.25 - 1960}}{9.8} \\
 &= \frac{73.5 \pm \sqrt{3442.25}}{9.8} \approx 1.51 \text{ or } 13.49
 \end{aligned}$$

The projectile will be 100 m above the ground after 1.51 sec and after 13.49 sec.

$$\begin{aligned}
 \text{(b)} \quad &\text{When the projectile returns to the ground, the height } s \text{ will be 0 ft, so let } s = 0 \text{ and solve for } t: \\
 &0 = -4.9t^2 + 73.5t \Rightarrow \\
 &t(-4.9t + 73.5) = 0 \Rightarrow t = 0 \text{ or} \\
 &-4.9t + 73.5 = 0 \Rightarrow t = 15
 \end{aligned}$$

The first solution, 0, represents the time at which the projectile was on the ground before being launched. The projectile will return to the ground 15 sec after being launched.

Classroom Example 4 (page 127)

$$\begin{aligned}
 \text{(a)} \quad S &= 0.0538x^2 - 0.807x + 8.84 \\
 \text{For 2013, } x &= 13. \\
 S &= 0.0538(13)^2 - 0.807(13) + 8.84 \approx 7.4 \\
 \text{million. This is \$0.2 million less than the actual} \\
 &\text{figure of \$7.6 million.}
 \end{aligned}$$

$$\begin{aligned}
 \text{(b)} \quad &\text{Let } S = 5.9, \text{ then solve for } x: \\
 5.9 &= 0.0538x^2 - 0.807x + 8.84 \Rightarrow \\
 0 &= 0.0538x^2 - 0.807x + 2.94 \\
 x &= \frac{0.807 \pm \sqrt{(-0.807)^2 - 4(0.0538)(2.94)}}{2(0.0538)} \\
 &= \frac{0.807 \pm \sqrt{0.018561}}{0.1076} \\
 &\approx \frac{0.807 \pm 0.1362}{0.1076} \\
 x &\approx 8.8 \text{ or } x \approx 6.2
 \end{aligned}$$

According to the model, sales reached 5.9 million about 6.2 years after 2000, or in 2006, and about 8.8 years after 2000, or in 2008.

Section 1.6 Other Types of Equations and Applications**Classroom Example 1 (page 136)**

$$\begin{aligned}
 \text{(a)} \quad &\frac{2x - 3}{2} + \frac{5x}{x + 1} = x \\
 &\text{The least common denominator is } 2(x + 1), \text{ which equals 0 when } x = -1. \text{ Therefore, } -1 \\
 &\text{cannot be a solution of the equation.} \\
 &2(x + 1)\left(\frac{2x - 3}{2} + \frac{5x}{x + 1}\right) = 2(x + 1)x \\
 &(x + 1)(2x - 3) + 2(5x) = 2x^2 + 2x \\
 &2x^2 - 3x + 2x - 3 + 10x = 2x^2 + 2x \\
 &2x^2 + 9x - 3 = 2x^2 + 2x \\
 &7x = 3 \\
 &x = \frac{3}{7}
 \end{aligned}$$

The restriction $x \neq -1$ does not affect the result. Therefore, the solution set is $\left\{\frac{3}{7}\right\}$.

(b) $\frac{x}{x-5} + 5 = \frac{5}{x-5}$

The least common denominator is $x - 5$, which equals 0 when $x = 5$. Therefore, 5 cannot be a solution of the equation.

$$\begin{aligned}(x-5)\left(\frac{x}{x-5} + 5\right) &= (x-5)\left(\frac{5}{x-5}\right) \\ x + 5(x-5) &= 5 \\ 6x - 25 &= 5 \Rightarrow 6x = 30 \Rightarrow x = 5\end{aligned}$$

The only possible solution is 5. However, the variable is restricted to real numbers except 5. Therefore, the solution set is \emptyset .

Classroom Example 2 (page 137)

(a) $\frac{x-5}{x-3} + \frac{1}{x} = \frac{-7}{x^2-3x}$

The least common denominator is

$x(x-3) = x^2 - 3x$, which is equal to 0 if $x = 3$ or $x = 0$. Therefore, 0 and 3 cannot possibly be solutions of this equation.

$$\begin{aligned}x(x-3)\left(\frac{x-5}{x-3} + \frac{1}{x}\right) &= x(x-3)\left(\frac{-7}{x^2-3x}\right) \\ x(x-5) + (x-3) &= -7 \\ x^2 - 5x + x - 3 &= -7 \\ x^2 - 4x + 4 &= 0 \\ (x-2)(x-2) &= 0 \Rightarrow x = 2\end{aligned}$$

The restrictions $x \neq 0$ or $x \neq 3$ do not affect the result. Therefore, the solution set is $\{2\}$.

(b) $\frac{x}{x+5} + \frac{5}{x-5} = \frac{50}{x^2-25}$

The least common denominator is

$(x+5)(x-5) = x^2 - 25$, which is equal to 0 if $x = \pm 5$.

$$\begin{aligned}(x+5)(x-5)\left(\frac{x}{x+5} + \frac{5}{x-5}\right) &= (x+5)(x-5)\left(\frac{50}{x^2-25}\right) \\ x(x-5) + 5(x+5) &= 50 \\ x^2 - 5x + 5x + 25 &= 50 \\ x^2 &= 25 \Rightarrow x = \pm 5\end{aligned}$$

Because of the restriction $x \neq \pm 5$, the solution set is \emptyset .

Classroom Example 3 (page 138)

Let x = the amount of time (in hours) it takes Lisa and Keith to rake the leaves.

	r	t	Part of the Job Accomplished
Lisa	$\frac{1}{5}$	x	$\frac{1}{5}x$
Keith	$\frac{1}{4}$	x	$\frac{1}{4}x$

Lisa and Keith must accomplish 1 job (raking the leaves), so we must solve the following equation.

$$\begin{aligned}\frac{1}{5}x + \frac{1}{4}x &= 1 \\ 20\left(\frac{1}{5}x + \frac{1}{4}x\right) &= 20 \cdot 1 \\ 4x + 5x &= 20 \\ 9x &= 20 \Rightarrow x = \frac{20}{9} = 2\frac{2}{9}\end{aligned}$$

It takes Lisa and Keith $2\frac{2}{9}$ hr working together to rake the leaves.

Classroom Example 4 (page 140)

$$\begin{aligned}x - \sqrt{4x+12} &= 0 \\ x &= \sqrt{4x+12} \\ x^2 &= 4x+12 \\ x^2 - 4x - 12 &= 0 \\ (x-6)(x+2) &= 0 \Rightarrow x = 6 \text{ or } x = -2 \\ \text{Check } x = -2: \\ -2 - \sqrt{4(-2)+12} &\stackrel{?}{=} 0 \\ -2 - \sqrt{-8+12} &= 0 \\ -2 - \sqrt{4} &= 0 \\ -2 - 2 &= 0 \\ -4 &\neq 0\end{aligned}$$

Thus, -2 is not a solution.

Check $x = 6$:

$$\begin{aligned}6 - \sqrt{4(6)+12} &\stackrel{?}{=} 0 \\ 6 - \sqrt{24+12} &= 0 \\ 6 - \sqrt{36} &= 0 \\ 6 - 6 &= 0 \\ 0 &= 0\end{aligned}$$

Thus, 6 is a solution.

Solution set: $\{6\}$

Classroom Example 5 (page 141)

$$\begin{aligned}
\sqrt{3x+1} - \sqrt{x+4} &= 1 \\
\sqrt{3x+1} &= \sqrt{x+4} + 1 \\
(\sqrt{3x+1})^2 &= (\sqrt{x+4} + 1)^2 \\
3x+1 &= (\sqrt{x+4})^2 + 2\sqrt{x+4} + 1 \\
3x+1 &= x+4 + 2\sqrt{x+4} + 1 \\
2x-4 &= 2\sqrt{x+4} \\
(2x-4)^2 &= (2\sqrt{x+4})^2 \\
4x^2 - 16x + 16 &= 4x + 16 \\
4x^2 - 20x &= 0 \\
x^2 - 5x &= 0 \Rightarrow x(x-5) = 0 \Rightarrow x = 0 \text{ or } x = 5
\end{aligned}$$

Check $x = 0$:

$$\begin{aligned}
\sqrt{3(0)+1} - \sqrt{(0)+4} &\stackrel{?}{=} 1 \\
1 - 2 &= 1 \\
-1 &\neq 1
\end{aligned}$$

Thus, $x = 0$ is not a solution.Check $x = 5$:

$$\begin{aligned}
\sqrt{3(5)+1} - \sqrt{(5)+4} &\stackrel{?}{=} 1 \\
4 - 3 &= 1 \Rightarrow 1 = 1
\end{aligned}$$

Thus, $x = 5$ is a solution. The solution set is $\{5\}$.**Classroom Example 6 (page 142)**

$$\begin{aligned}
\sqrt[3]{5x^2 - 12x + 6} - \sqrt[3]{x} &= 0 \\
\sqrt[3]{5x^2 - 12x + 6} &= \sqrt[3]{x} \\
(\sqrt[3]{5x^2 - 12x + 6})^3 &= (\sqrt[3]{x})^3 \\
5x^2 - 12x + 6 &= x \\
5x^2 - 13x + 6 &= 0 \\
(5x-3)(x-2) &= 0 \Rightarrow x = \frac{3}{5} \text{ or } x = 2
\end{aligned}$$

Check $x = \frac{3}{5}$:

$$\begin{aligned}
\sqrt[3]{5\left(\frac{3}{5}\right)^2 - 12\left(\frac{3}{5}\right) + 6} - \sqrt[3]{\frac{3}{5}} &\stackrel{?}{=} 0 \\
\sqrt[3]{\frac{9}{5} - \frac{36}{5} + 6} - \sqrt[3]{\frac{3}{5}} &= 0 \\
\sqrt[3]{\frac{3}{5}} - \sqrt[3]{\frac{3}{5}} &= 0 \Rightarrow 0 = 0
\end{aligned}$$

Thus, $x = \frac{3}{5}$ is a solution.Check $x = 2$:

$$\begin{aligned}
\sqrt[3]{5(2)^2 - 12(2) + 6} - \sqrt[3]{2} &\stackrel{?}{=} 0 \\
\sqrt[3]{20 - 24 + 6} - \sqrt[3]{2} &= 0 \\
\sqrt[3]{2} - \sqrt[3]{2} &= 0 \Rightarrow 0 = 0
\end{aligned}$$

Thus, $x = 2$ is a solution.Solution set: $\left\{\frac{3}{5}, 2\right\}$ **Classroom Example 7 (page 142)**

$$\begin{aligned}
\text{(a)} \quad x^{3/2} &= 216 \\
(x^{3/2})^{2/3} &= 216^{2/3} \\
x &= (\sqrt[3]{216})^2 = 6^2 = 36 \\
\text{Solution set: } &\{36\} \\
\text{(b)} \quad (x+7)^{2/5} &= 4 \\
[(x+7)^{2/5}]^{5/2} &= \pm 4^{5/2} \\
x+7 &= \pm (\sqrt{4})^5 \\
x+7 &= 2^5 = 32 \Rightarrow x = 25 \text{ or} \\
x+7 &= -2^5 = -32 \Rightarrow x = -39 \\
\text{Solution set: } &\{-39, 25\}
\end{aligned}$$

Classroom Example 8 (page 144)

$$\begin{aligned}
\text{(a)} \quad (x-3)^{1/2} - 6(x-3)^{1/4} + 8 &= 0 \\
\text{Let } u &= (x-3)^{1/4}, \text{ so} \\
u^2 &= [(x-3)^{1/4}]^2 = (x-3)^{1/2} \\
\text{Substituting, we have} \\
u^2 - 6u + 8 &= 0 \\
(u-4)(u-2) &= 0 \Rightarrow u = 4 \text{ or } u = 2 \\
\text{Now solve for } x, \text{ by replacing } u &\text{ with } (x-3)^{1/4}: \\
4 &= (x-3)^{1/4} \Rightarrow 4^4 = x-3 \Rightarrow 259 = x \\
2 &= (x-3)^{1/4} \Rightarrow 2^4 = x-3 \Rightarrow 19 = x \\
\text{Check } x = 259: \\
(259-3)^{1/2} - 6(259-3)^{1/4} + 8 &\stackrel{?}{=} 0 \\
\sqrt{256} - 6\sqrt[4]{256} + 8 &= 0 \\
16 - 6(4) + 8 &= 0 \\
0 &= 0
\end{aligned}$$

Thus, $x = 259$ is a solution.Check $x = 19$:

$$\begin{aligned}
(19-3)^{1/2} - 6(19-3)^{1/4} + 8 &\stackrel{?}{=} 0 \\
\sqrt{16} - 6\sqrt[4]{16} + 8 &= 0 \\
4 - 6(2) + 8 &= 0 \\
0 &= 0
\end{aligned}$$

Thus, $x = 19$ is a solution.Solution set: $\{19, 259\}$.

(b) $15x^{-2} - 4x^{-1} = 3 \Rightarrow 15x^{-2} - 4x^{-1} - 3 = 0$

Let $u = x^{-1}$. Substituting, we have

$$15u^2 - 4u - 3 = 0$$

$$(5u - 3)(3u + 1) = 0 \Rightarrow u = \frac{3}{5} \text{ or } u = -\frac{1}{3}$$

Now solve for x , by replacing u with x^{-1} :

$$x^{-1} = \frac{3}{5} \Rightarrow x = \frac{5}{3}; \quad x^{-1} = -\frac{1}{3} \Rightarrow x = -3$$

Check $x = \frac{5}{3}$:

$$15\left(\frac{5}{3}\right)^{-2} - 4\left(\frac{5}{3}\right)^{-1} \stackrel{?}{=} 3$$

$$15\left(\frac{3}{5}\right)^2 - 4\left(\frac{3}{5}\right) = 3 \Rightarrow 15\left(\frac{9}{25}\right) - \frac{12}{5} = 3$$

$$\frac{27}{5} - \frac{12}{5} = 3 \Rightarrow 3 = 3$$

Thus, $x = \frac{5}{3}$ is a solution.

Check $x = -3$:

$$15(-3)^{-2} - 4(-3)^{-1} \stackrel{?}{=} 3$$

$$15\left(-\frac{1}{3}\right)^2 - 4\left(-\frac{1}{3}\right) = 3 \Rightarrow 15\left(\frac{1}{9}\right) + \frac{4}{3} = 3$$

$$\frac{5}{3} + \frac{4}{3} = 3 \Rightarrow 3 = 3$$

Thus, $x = -3$ is a solution.

Solution set: $\left\{-3, \frac{5}{3}\right\}$

Classroom Example 9 (page 145)

$$18x^4 - 29x^2 + 3 = 0$$

Let $u = x^2$; then $u^2 = x^4$. With this substitution, the equation becomes $18u^2 - 29u + 3 = 0$. Solving, we have $18u^2 - 29u + 3 = 0$

$$(2u - 3)(9u - 1) = 0 \Rightarrow u = \frac{3}{2} \text{ or } u = \frac{1}{9}$$

Now solve for x : $x^2 = \frac{3}{2} \Rightarrow x = \pm\sqrt{\frac{3}{2}} = \pm\frac{\sqrt{6}}{2}$ or

$x^2 = \frac{1}{9} \Rightarrow x = \pm\sqrt{\frac{1}{9}} = \pm\frac{1}{3}$. Checking each proposed solution, we find that all are solutions.

Solution set: $\left\{\pm\frac{\sqrt{6}}{2}, \pm\frac{1}{3}\right\}$

Section 1.7 Inequalities

Classroom Example 1 (page 150)

$$-2x + 7 < -5$$

$$-2x < -12$$

$$x > 6$$

Solution set: $\{x \mid x > 6\}$

Classroom Example 2 (page 152)

$$3 - 4x \geq 2x + 8$$

$$-5 \geq 6x$$

$$-\frac{5}{6} \geq x \Rightarrow x \leq -\frac{5}{6}$$

Solution set: $\left(-\infty, -\frac{5}{6}\right]$

Classroom Example 3 (page 152)

$$R = 45x \text{ and } C = 30x + 5250$$

Set $R \geq C$, and solve for x :

$$45x \geq 30x + 5250$$

$$15x \geq 5250$$

$$x \geq 350$$

The break-even point is at $x = 350$. This product will at least break even only if the number of units produced and sold is in the interval $[350, \infty)$.

Classroom Example 4 (page 153)

$$1 \leq 6x - 8 \leq 4$$

$$9 \leq 6x \leq 12$$

$$\frac{3}{2} \leq x \leq 2$$

Solution set: $\left[\frac{3}{2}, 2\right]$

Classroom Example 5 (page 153)

$$x^2 - 2x - 15 \leq 0$$


Step 1: Find the values of x that satisfy

$$x^2 - 2x - 15 = 0.$$

$$x^2 - 2x - 15 = 0 \Rightarrow (x - 5)(x + 3) = 0 \Rightarrow$$

$$x = 5 \text{ or } x = -3$$

Step 2: The two numbers divide a number line into three regions.



(continued on next page)

(continued)

Step 3: Choose a test value to see if it satisfies the inequality $x^2 - 2x - 15 \leq 0$.

Interval	Test Value	Is $x^2 - 2x - 15 \leq 0$ True or False?
A: $(-\infty, -3]$	-5	$(-5)^2 - 2(-5) - 15 \stackrel{?}{\leq} 0$ $20 \leq 0$ False
B: $[-3, 5]$	0	$0^2 - 2(0) - 15 \stackrel{?}{\leq} 0$ $-15 \leq 0$ True
C: $[5, \infty)$	6	$6^2 - 2(6) - 15 \stackrel{?}{\leq} 0$ $9 \leq 0$ False

Solution set: $[-3, 5]$

Classroom Example 6 (page 154)

$$3x^2 - 11x - 4 > 0$$


Step 1: Find the values of x that satisfy

$$3x^2 - 11x - 4 = 0.$$

$$3x^2 - 11x - 4 = 0 \Rightarrow (3x + 1)(x - 4) = 0 \Rightarrow$$

$$x = -\frac{1}{3} \text{ or } x = 4$$

Step 2: The two numbers divide a number line into three regions.



Step 3: Choose a test value to see if it satisfies the inequality $3x^2 - 11x - 4 > 0$.

Interval	Test Value	Is $3x^2 - 11x - 4 > 0$ True or False?
A: $(-\infty, -\frac{1}{3})$	-1	$3(-1)^2 - 11(-1) - 4 \stackrel{?}{>} 0$ $10 > 0$ True
B: $(-\frac{1}{3}, 4)$	0	$3(0)^2 - 11(0) - 4 \stackrel{?}{>} 0$ $-4 > 0$ False
C: $(4, \infty)$	5	$3(5)^2 - 11(5) - 4 \stackrel{?}{>} 0$ $16 > 0$ True

Solution set: $(-\infty, -\frac{1}{3}) \cup (4, \infty)$

Classroom Example 7 (page 155)

$$s = -16t^2 + 144t$$

$$-16t^2 + 144t > 128$$


Step 1: Find the values of t that satisfy

$$-16t^2 + 144t = 128$$

$$-16t^2 + 144t = 128 \Rightarrow 0 = 16t^2 - 144t + 128 \Rightarrow$$

$$16(t - 8)(t - 1) = 0 \Rightarrow t = 8 \text{ or } t = 1$$

Step 2: The two numbers divide a number line into three regions. (Note that time $t \geq 0$.)



Step 3: Choose a test value to see if it satisfies the inequality $-16t^2 + 144t > 128$.

Interval	Test Value	Is $-16t^2 + 144t > 128$ True or False?
A: $[0, 1)$	0.5	$-16(0.5)^2 + 144(0.5) \stackrel{?}{>} 128$ $68 > 128$ False
B: $(1, 8)$	2	$-16(2)^2 + 144(2) \stackrel{?}{>} 128$ $224 > 128$ True
C: $(8, \infty)$	10	$-16(10)^2 + 144(10) \stackrel{?}{>} 128$ $-160 > 128$ False

Solution set: $(1, 8)$

The object will be greater than 128 ft above ground level between 1 and 8 seconds after it is launched.

Classroom Example 8 (page 156)

$$\frac{6}{x-3} \geq 4$$

Step 1: Rewrite the inequality so that 0 is on one side and there is a single fraction on the other side.

$$\frac{6}{x-3} \geq 4 \Rightarrow \frac{6}{x-3} - 4 \geq 0 \Rightarrow$$

$$\frac{6}{x-3} - \frac{4(x-3)}{x-3} \geq 0 \Rightarrow$$

$$\frac{6-4x+12}{x-3} \geq 0 \Rightarrow \frac{18-4x}{x-3} \geq 0$$


Step 2: Determine the values that will cause either the numerator or denominator to equal 0.

$$18 - 4x = 0 \Rightarrow x = \frac{9}{2} \text{ or } x - 3 = 0 \Rightarrow x = 3$$

(continued on next page)

(continued)

The values $\frac{9}{2}$ and 3 divide the number line into three regions. Use an open circle on 3 because it makes the denominator equal 0.



Step 3: Choose a test value to see if it satisfies the inequality, $\frac{6}{x-3} \geq 4$

Interval	Test Value	Is $\frac{6}{x-3} \geq 4$ True or False?
A: $(-\infty, 3)$	0	$\frac{6}{0-3} \geq 4$ $-2 \geq 4$ False
B: $(3, \frac{9}{2})$	4	$\frac{6}{4-3} \geq 4$ $6 \geq 4$ True
C: $[\frac{9}{2}, \infty)$	5	$\frac{6}{5-3} \geq 4$ $3 \geq 4$ False

Solution set: $(3, \frac{9}{2})$

Classroom Example 9 (page 157)

$$\frac{3x+1}{2x-3} < 4$$

Step 1: Rewrite the inequality so that 0 is on one side and there is a single fraction on the other side.

$$\frac{3x+1}{2x-3} < 4 \Rightarrow \frac{3x+1}{2x-3} - 4 < 0 \Rightarrow$$

$$\frac{3x+1}{2x-3} - \frac{4(2x-3)}{2x-3} < 0 \Rightarrow$$


$$\frac{3x+1-8x+12}{2x-3} < 0 \Rightarrow$$

$$\frac{-5x+13}{2x-3} < 0$$

Step 2: Determine the values that will cause either the numerator or denominator to equal 0.

$$-5x+13=0 \Rightarrow x=\frac{13}{5} \text{ or } 2x-3=0 \Rightarrow x=\frac{3}{2}. \text{ The}$$

values $\frac{3}{2}$ and $\frac{13}{5}$ divide the number line into three regions.



Step 3: Choose a test value to see if it satisfies the inequality, $\frac{3x+1}{2x-3} < 4$

Interval	Test Value	Is $\frac{3x+1}{2x-3} < 4$ True or False?
A: $(-\infty, \frac{3}{2})$	0	$\frac{3(0)+1}{2(0)-3} < 4$ $-\frac{1}{3} < 4$ True
B: $(\frac{3}{2}, \frac{13}{5})$	2	$\frac{3(2)+1}{2(2)-3} < 4$ $7 < 4$ False
C: $(\frac{13}{5}, \infty)$	5	$\frac{3(5)+1}{2(5)-3} < 4$ $\frac{16}{7} < 4$ True

Solution set: $(-\infty, \frac{3}{2}) \cup (\frac{13}{5}, \infty)$

Section 1.8 Absolute Value Equations and Inequalities**Classroom Example 1 (page 163)**

(a) $|9-4x|=7$

$$9-4x=7 \Rightarrow -4x=-2 \Rightarrow x=\frac{1}{2} \text{ or}$$

$$9-4x=-7 \Rightarrow -4x=-16 \Rightarrow x=4$$

$$\text{Solution set: } \left\{\frac{1}{2}, 4\right\}$$

(b) $|3x+2|=|x-5|$

$$3x+2=x-5 \Rightarrow 2x=-7 \Rightarrow x=-\frac{7}{2} \text{ or}$$

$$3x+2=-(x-5) \Rightarrow 3x+2=-x+5 \Rightarrow$$

$$4x=3 \Rightarrow x=\frac{3}{4}$$

$$\text{Solution set: } \left\{-\frac{7}{2}, \frac{3}{4}\right\}$$

Classroom Example 2 (page 164)

(a) $|4x-6|<10$

$$-10<4x-6<10 \Rightarrow -4<4x<16 \Rightarrow$$

$$-1<x<4$$

$$\text{Solution set: } (-1, 4)$$

(b) $|4x-6|>10$

$$4x-6<-10 \text{ or } 4x-6>10$$

$$4x<-4 \text{ or } 4x>16$$

$$x<-1 \text{ or } x>4$$

$$\text{Solution set: } (-\infty, -1) \cup (4, \infty)$$

Classroom Example 3 (page 165)

$$|5 - 8x| + 6 \geq 14$$

$$|5 - 8x| \geq 8$$

$$5 - 8x \leq -8 \quad \text{or} \quad 5 - 8x \geq 8$$

$$-8x \leq -13 \quad \text{or} \quad -8x \geq 3$$

$$x \geq \frac{13}{8} \quad \text{or} \quad x \leq -\frac{3}{8}$$

$$\text{Solution set: } \left(-\infty, -\frac{3}{8}\right] \cup \left[\frac{13}{8}, \infty\right)$$

Classroom Example 4 (page 165)

(a) $|7x + 28| = 0$

The absolute value of a number will be 0 if that number is 0. Therefore $|7x + 28| = 0$ is equivalent to $7x + 28 = 0$, which has solution set $\{-4\}$.

(b) $|6x - 9| > -2$

The absolute value of a number is always nonnegative, so the inequality $|6x - 9| > -2$ is always true. The solution set is $(-\infty, \infty)$.

(c) $|2 - 5x| \leq -5$

There is no number whose absolute value is less than any negative number. The solution set of $|2 - 5x| \leq -5$ is \emptyset .

Classroom Example 5 (page 166)

(a) “ m is no more than 9 units from 3” means that m is 9 units or less from 3. Thus the distance between m and 3 is less than or equal to 9, or $|m - 3| \leq 9$.

(b) “ t is within 0.02 unit of 5.8” means that t is less than 0.02 unit from 5.8. Thus the distance between t and 5.8 is less than 0.02, or $|t - 5.8| < 0.02$.

Classroom Example 6 (page 166)

We want y to be within 0.001 unit of 6, so we have

$$|y - 6| < 0.001 \quad \text{or} \quad |(5x - 2) - 6| < 0.001.$$

$$|5x - 8| < 0.001$$

$$-0.001 < 5x - 8 < 0.001$$

$$7.999 < 5x < 8.001$$

$$1.5998 < x < 1.6002$$

Values of x in the interval $(1.5998, 1.6002)$ will satisfy the condition.