

INSTRUCTOR'S GUIDE TO MATHEMATICS ACTIVITIES FOR ELEMENTARY SCHOOL TEACHERS

DAN DOLAN

*Project to Increase Mastery of Mathematics
and Science, Wesleyan University*

JIM WILLIAMSON

*University
of Montana*

MARI MURI

*Project to Increase Mastery of Mathematics
and Science, Wesleyan University*

to accompany

A PROBLEM SOLVING APPROACH TO MATHEMATICS FOR ELEMENTARY SCHOOL TEACHERS THIRTEENTH EDITION

Rick Billstein

University of Montana

Barbara Boschmans


Northern Arizona University

Shlomo Libeskind

University of Oregon

Johnny W. Lott

University of Montana



The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2020, 2016, 2013 by Pearson Education, Inc. 221 River Street, Hoboken, NJ 07030. All rights reserved.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-499717-9

ISBN-10: 0-13-499717-4

Table of Contents

Chapter	1	An Introduction to Problem Solving	1
Activity	1	When You Don't Know What to Do	1
Activity	2	What's the Pattern?	3
Activity	3	Ten People in a Canoe	5
Activity	4	An Ancient Game	5
Activity	5	Magic Number Tricks	6
Activity	6	Eliminate the Impossible	6
Activity	7	What's the Rule?	7
Activity	8	Fascinating Fibonacci	8
Activity	9	Paper Powers	9
Chapter	2	Introduction to Logic and Sets	11
Activity	1	What's in the Loop?	11
Activity	2	Loop de Loops	11
Chapter	3	Numeration Systems and Whole Number Operations	13
Activity	1	Regrouping Numbers	13
Activity	2	Find the Missing Numbers	13
Activity	3	A Visit to Fouria	13
Activity	4	It All Adds Up	14
Activity	5	What's the Difference?	14
Activity	6	Multiplication Arrays	15
Activity	7	Multi-digit Multiplication	15
Activity	8	Least and Greatest	15
Activity	9	Target Number	15
Activity	10	Find the Missing Factor	16
Activity	11	How Many Cookies?	16
Activity	12	Multi-digit Division	16
Chapter	4	Number Theory	17
Activity	1	Odd and Even Patterns	17
Activity	2	Great Divide Game	17
Activity	3	A Square Experiment	17
Activity	4	The Factor Game	19
Activity	5	A Sieve of Another Sort	20
Activity	6	Tiling with Squares	21
Activity	7	Pool Factors	23

Chapter	5	Integers	25
Activity	1	Charged Particles	25
Activity	2	Coin Counters	25
Activity	3	Clown on a Tightrope	26
Activity	4	Subtracting Integers	27
Activity	5	Subtraction Patterns	27
Activity	6	Multiplication and Division Patterns	28
Activity	7	Integer $\times$ and $\div$ Contig	28
Chapter	6	Rational Numbers and Proportional Reasoning	29
Activity	1	What is a Fraction?	29
Activity	2	Square Fractions	30
Activity	3	Equivalent Fractions	30
Activity	4	How Big Is It?	31
Activity	5	What Comes First?	31
Activity	6	Fraction War	31
Activity	7	Adding and Subtracting Fractions	32
Activity	8	Multiplying Fractions	33
Activity	9	Dividing Fractions	34
Activity	10	Professors Short and Tall	34
Chapter	7	Decimals, Percents, and Real Numbers	37
Activity	1	What's My Name?	37
Activity	2	Deci-Order	37
Activity	3	Race for the Flat	37
Activity	4	Empty the Board	37
Activity	5	Decimal Arrays	38
Activity	6	Decimal Multiplication	38
Activity	7	Dice and Decimals	38
Activity	8	Target Number Revisited	38
Activity	9	Repeating Decimals	38
Activity	10	What is Percent?	40
Chapter	8	Algebraic Thinking	41
Activity	1	Regular Polygons in a Row	41
Activity	2	Patterns and Expressions	42
Activity	3	What's My Function?	43
Activity	4	Graphing Rectangles	43
Activity	5	Exploring Linear Functions	46
Chapter	9	Probability	49
Activity	1	The Spinner Game	49
Activity	2	Theoretical Probability	50
Activity	3	It's in the Bag	52
Activity	4	Paper-Scissors-Rock	52
Activity	5	Pascal's Probabilities	54
Activity	6	Simulate It	55
Activity	7	How Many Arrangements?	56

Chapter 10	Data Analysis/Statistics: An Introduction	61
Activity 1	Graphing <i>m&m's</i> [®]	61
Activity 2	Grouped Data	62
Activity 3	Are Women Catching Up?	64
Activity 4	What's the Average?	68
Activity 5	Finger-Snapping Time	69
Activity 6	Populations and Samples	69
Activity 7	The Weather Report	70
Chapter 11	Introductory Geometry	73
Activity 1	What's the Angle?	73
Activity 2	Inside or Outside?	73
Activity 3	Name That Polygon	74
Activity 4	Triangle Properties—Angles	75
Activity 5	Angles on Pattern Blocks	75
Activity 6	Sum of Interior/Exterior Angles	76
Activity 7	Stars and Angles	77
Activity 8	Spatial Visualization	78
Chapter 12	Congruence and Similarity with Constructions	79
Activity 1	Triangle Properties—Sides	79
Activity 2	To Be or Not to Be Congruent?	79
Activity 3	Paper Folding Constructions	80
Activity 4	Pattern Block Similarity	81
Activity 5	Similar Triangles	82
Activity 6	Side Splitter Theorem	83
Activity 7	Outdoor Geometry	84
Chapter 13	Area, Pythagorean Theorem, and Volume	85
Activity 1	Areas of Polygons	85
Activity 2	Pythagorean Puzzles	85
Activity 3	Now You See It, Now You Don't	86
Activity 4	Right or Not?	87
Activity 5	Surface Area	87
Activity 6	Volume of a Rectangular Solid	87
Activity 7	Pyramids and Cones	88
Chapter 14	Transformations	89
Activity 1	Tessellations	89
Activity 2	Reflections	91
Activity 3	Glide Reflections	93
Activity 4	Dilations	94


Chapter 1

An Introduction to Problem Solving

Activity 1

Box 1: UNDERSTAND THE PROBLEM

1. Answers will vary.
2. Sample Responses: Shapes are congruent if they have exactly the same shape and size.
Shapes are congruent if they can be made to match exactly by placing one on top of the other.

3. a. 

b. 3 ways

4. Answers will vary.
5. Answers will vary.

Box 2: DEVISE A PLAN

1. Squares could be moved around to try to construct the staircases or they could be sketched on graph paper.
2. Simplify the problem, look for a pattern, and make a table
3. Answers will vary.

(Continued on next page.)

Activity 1 Continued**Box 4: LOOK BACK**

1. To form two congruent staircases, the number of squares must be even and you must be able to form a rectangle in which both dimensions are greater than one and one dimension is odd and the other is even using the squares.
3. a. 3 ways
- b. Since the two staircases must fit together to form a rectangle with area 120, find the dimensions of all rectangles with integral dimensions and an area of 120. The possible dimensions are 1×120 , 2×60 , 3×40 , 4×30 , 5×24 , 6×20 , 8×15 , 10×12 .

To form a staircase, both dimensions must be greater than 1, and one dimension must be even and the other odd. This leaves 3 possibilities, a 3×40 rectangle, a 5×24 rectangle, and an 8×15 rectangle.

The 3×40 rectangle can be divided into two 3-step staircases. The bottom step is 21 squares long, the middle step is 20 steps long, and the top step is 19 squares long.

The 5×24 rectangle can be divided into two 5-step staircases. The bottom step is 14 squares long, the next step is 13 squares long, the middle step is 12 steps long, the fourth step is 11 squares long and the top step is 10 squares long.

The 8×15 rectangle can be divided into two 8-step staircases. The steps are 11, 10, 9, 8, 7, 6, 5, and 4 steps long.

4. To determine whether n squares can be used to construct two congruent staircases, find the factor pairs of n . There is a pair of staircases for each factor pair in which both factors are greater than 1 and one factor is odd and the other even.

(Since the two staircases must be congruent, each must contain the same number of squares.

So if n squares can be arranged into two congruent staircases, n must be even.


Suppose a staircase has s steps and that the top step contains k squares. There are $s - 1$ steps below the top step and each step down contains one more square than the step above it, so the bottom step contains $s - 1$ more squares than the top step. That is, the bottom step contains $k + (s - 1)$ squares. When two of the staircases are put together to form a rectangle, the height of the rectangle is s and the length is $k + k + (s - 1) = 2k + (s - 1)$.

If s is odd, $s - 1$ is even and $2k + (s - 1)$ is even since the sum of two even numbers is even.

If s is even, $s - 1$ is odd and $2k + (s - 1)$ is odd since the sum of an odd and an even number is odd.




So if n squares can be arranged into two congruent staircases, you must be able to make a rectangle with the squares in which one dimension is odd and the other dimension is even.

Suppose one dimension of a rectangle is odd and the other is even. Orient the rectangle so the height is odd and the length is even. Starting at the center of the middle row of squares, form steps going up to the right and steps going down to the left. Since the number of rows above the middle row and the number of rows below it are equal, this process will result in two congruent staircases.)




Activity 2

Box 1

1. 
2. 
3. 
4. 6, 3, 4, 5, 6, 7
5. 14, 17, 20, 23, 26, 29
6. 25, 18, 11, 4, -3, -10
7. 15, 21, 28, 36, 45, 55
8. 28, 39, 52, 67, 84, 103
9. 32, 64, 128, 256, 512, 1024
10. 9, 3, 1, $\frac{1}{3}$, $\frac{1}{9}$, $\frac{1}{27}$
11. 34, 55, 89, 144, 233, 377

12. Each term of the sequence in Exercise 5 is the number of squares in the corresponding term of the sequence in Exercise 1. Similarly, each term of the sequence in Exercise 7 is the number of triangles in the corresponding term of the sequence in Exercise 3.
13. a. The number of triangles in each term of the sequence in Exercise 3 is half the number of squares in the corresponding term of the sequence in Exercise 2.
b. Sample Response:



(Continued on next page.)

Activity 2 Continued**Box 2**

1. 7, 12, 17, 22, 27, 32, 37, 42, 47
2. 2, 5, 8, 11, 14, 17, 20, 23, 26

Each term of the sequence in Exercise 1 on page 5 is a rectangular array 3 squares tall and n squares long (where n is the term number) with the top right-hand square removed. Thus the number of squares used to form each term is 1 less than 3 times n .

3. 44, 41, 38, 35, 32, 29, 26, 23, 20
4. 7, 13, 23, 37, 55, 77, 103, 133, 167
5. 2, 6, 12, 20, 30, 42, 56, 72, 90

Each term of the sequence in Exercise 2 on page 5 is a rectangular array of squares. The height of the rectangle is equal to the term number and its length is 1 more than the term number. Since the area of the rectangle is the length times the width, the number of squares in the array is *the term number times the term number plus 1*. That is, *the term number times the next term number*.

6. In Exercises 1–3, the difference between successive terms is a constant. In Exercise 1 the difference is always 5, in Exercise 2 it is always 3, and in Exercise 3 it is -3 . In each case, the difference is the constant by which the term number was multiplied.

The differences between successive terms in Exercises 4 and 5 are not constant. Note that, in these cases, the rule for generating the sequence does not involve multiplying the term number by a constant.

7. Answers will vary. One possibility is to look for a pattern in the differences between successive terms and to continue that pattern to find the next five terms in the sequence. For example, in Exercise 5 the differences are the sequence of even numbers 4, 6, 8, 10, 12, 14, 16, 18... Thus, the next five terms would be $90 + 20 = 110$, $110 + 22 = 132$, $132 + 24 = 156$, $156 + 26 = 182$, and $182 + 28 = 210$.
8. 2, 4, 8, 16, 30, **52, 84, 128**, 186, **260**
9. The first four terms of the sequences are the same, but from there on they are different. So no matter how many of the initial terms of a sequence you may know, there may be more than one way to extend the sequence.

Activity 3

2.	Number of People	2	4	6	8	10	$2N$
	Number of Pairs	1	2	3	4	5	N
	Min Number of Moves	3	8	15	24	35	$N^2 + 2N$
	Sequence of Moves	RLR	RLLRRL LR	RLLRRR LLLRRR LLR	RLLRRR LLLRRR RRLLLL RRRLLR	RLLRRR LLLRRR RRRLLL LLRRRR RLLLLR RRLLR	1R, 2L, 3R, ... , $(N-1)L$, NR, ML, NR, $(N-1)L$, ... , 3R, 2L, 1R if N is odd 1R, 2L, 3R, ... , $(N-1)R$, ML, NR, ML, $(N-1)R$, ... , 3R, 2L, 1R if N is even

4. 12 people

5. 255 moves

6.	Number of People	1	3	5	7	9	N
	Min Number of Moves	1	5	11	19	29	$(N^2 + 4N - 1)/4$
	Sequence of Moves	R	RLRRL	RLLRRR LLRRL	RLLRRR LLLRRR RLLLLR L	RLLRRR LLLRRR RRRLLL LRRRRL LLRRL	

The Legend of the Tower of Brahma

- | | | | | | | | |
|----|-----------------|---|---|---|----|----|-----------|
| 1. | Number of Disks | 1 | 2 | 3 | 4 | 5 | n |
| | Number of Moves | 1 | 3 | 7 | 15 | 31 | $2^n - 1$ |
2. a. $2^{10} - 1 = 17 \text{ min } 3 \text{ sec}$
 - b. $2^{30} - 1 \approx 34 \text{ years}$
 - c. $2^{50} - 1 \approx 36 \text{ million years}$
 - d. $2^{64} - 1 \approx 585 \text{ billion years}$

Activity 4

2. a. 1 Subtracting 1 leaves 4 on the display.
If Player A subtracts 1, that leaves 3 on the display and B wins by subtracting 3.
If A subtracts 2, that leaves 2 on the display and B wins by subtracting 2.
If A subtracts 3, it leaves 1 on the display and B wins by subtracting 1.
- b. To be certain to win, you want to be the first player. At the end of your turn you want to leave 16, 12, 8, or 4 on the display.
3. a. In this game you want to be the second player. At the end of your turn you want to leave 20, 15, 10, or 5 on the display.
- b. In this game you want to be the second player. At the end of your turn you want to leave 40, 30, 20, or 10 on the display.

Activity 5**Box 1**

1. The results for Steps 1-5 will vary depending on the number selected by each person.
2. Sample Response: 10 because that was the answer I got for Persons 1 through 5.
3. Step 1: n
 Step 2: $3n$
 Step 3: $3n + 30$
 Step 4: $(3n + 30) \div 3 = n + 10$
 Step 5: $n + 10 - n$
 The result is 10.

Box 3

1. Answers will vary.
2. The display on the calculator shows the month and day of a person's birthday. For example, 1021 for October 21.
3. Step 1: m
 Step 2: $5m$
 Step 3: $5m + 20$
 Step 4: $20m + 80$
 Step 5: $20m + 73$
 Step 6: $100m + 365$
 Step 7: $100m + 365 + d$
 Step 8: $100m + d$

In Step 8, multiplying the month, m , by 100 moves the month to the hundreds place in the final answer. If m is a single digit number, that digit will be in the hundreds place of the answer. If m is a two-digit number, its tens digit will be in the thousands place of the answer and its units digit will be in the hundreds place. When the day, d , is added, the digits of d will appear in the tens and units places of the final answer.

Activity 6**Box 1**

1. 153 (or 1 if it isn't a multi-digit number)
2. Sample Response: Since $7^3 = 343$, from the last two clues I knew the digits in the number had to be less than 7. From the second clue, I knew the digits could only be 1, 3, or 5.
3. 25 cards or 85 cards

Box 2

1. No. If there were green gumballs in the jar, it would be labeled correctly, but no jar has a correct label.
2. Red
3. No. If so, the jar labeled GREEN would have GREEN gumballs, but all labels are incorrect.
4. The correct label for the jar labeled Red - Green is Red, the correct label for the jar labeled Red is Green, and the correct label for the jar labeled Green is Red - Green.

(Continued on next page.)

Activity 6 Continued**Box 3**

State	Alabama	Alaska	Oklahoma	Minnesota
Flower	Camellia	Forget Me Not	Mistletoe	Lady's Slipper
Bird	Yellowhammer	Willow ptarmigan	Flycatcher	Loon

Clues b, d, and c are the keys to solving the problem.

From b, we know that Alaska and forget me not go together.

From d, we know that Alabama and yellowhammer go together.

From c, either camellia or lady's slipper goes with Minnesota. If camellia and Minnesota go together, then the state bird can't be the willow ptarmigan (clue e) or the loon (clue c) so it must be the flycatcher. But this would contradict clue a, so Minnesota, lady's slipper, and loon must go together.

Box 4

Two answers are possible. Freddie or Susie can be in either First or Fourth place.

First	Second	Third	Fourth
Yellow	Purple	Green	Red
Freddie or Susie	Liz	Joe	Susie or Freddie

Activity 7**Box 1**

Term Number	1	2	3	4	5	6	7	8
Term	4	11	18	25	32	39	46	53
Difference		<u>7</u>	<u>7</u>	<u>7</u>	<u>7</u>	<u>7</u>	<u>7</u>	<u>7</u>

Constant difference = 7

Term Number	Constant Difference	What Was Done?	To Get	
1	$\square$ 7	$\rightarrow$ 7	$\underline{-3}$	= 4 First Term
2	$\square$ 7	$\rightarrow$ <u>14</u>	$\underline{-3}$	= 11 Second Term
3	$\square$ <u>7</u>	$\rightarrow$ <u>21</u>	$\underline{-3}$	= <u>18</u> Third Term
10	$\square$ <u>7</u>	$\rightarrow$ <u>70</u>	$\underline{-3}$	= <u>67</u> Tenth Term
50	$\square$ <u>7</u>	$\rightarrow$ <u>350</u>	$\underline{-3}$	= <u>347</u> Fiftieth Term

Each term is 3 less than 7 times the term number,

Rule: n th term = $7n - 3$

Box 2

	Rule	25 th Term	100 th Term
1. 25, 29, 33, 37	$4T + 5$	105	405
2. 30, 37, 44, 51	$7T - 5$	170	695
3. 5, 7, 9, 11	$2T - 5$	45	195
4. 90, 88, 86, 84	$-2T + 100$	50	-100

$$5. \quad 65, \quad 62, \quad 59, \quad 56 \quad -3T + 80 \quad 5 \quad -220$$

Activity 8**Box 1**

1.

Beginning of month	Number of newborn pairs	Number of 1-month-old pairs	Number of pairs 2 months old or older	Total number of pairs
1	1	0	0	1
2	0	1	0	1
3	1	0	1	2
4	1	1	1	3
5	2	1	2	5
6	3	2	3	8
7	5	3	5	13

2. a. The number of 1-month old pairs in month N ($N \geq 2$) is equal to the number of newborn pairs in month $N - 1$.
- b. The number of pairs 2 months old or older in month N ($N \geq 2$) is equal to the sum of the number of 1-month old pairs and the number of pairs 2-months old or older in month $N - 1$.
- c. From the second month on, the number of newborn pairs each month is equal to the number of pairs 2 months old or older.

Box 2: CREATING A SPREADSHEET

2. a. 1 was added to the 1 in cell A2, so a 2 appears in cell A3.
 - b. The formula “= A3 + 1” is copied into cell A4, the formula “= A4 + 1” is copied into cell A5, and so on. Thus for each cell A4 through A14, 1 is added to the number in the preceding cell. The result is that the numbers 3 through 13 appear in cells A4 through A14 respectively.
 - c. To find the number of pairs after one year, you must extend the table to the beginning of the 13th month.
3. a. The content of cell B2 was copied into cell C3, so a 1 appears in cell C3.

Month	Newborn	1 Month Old	2 Months Old	Total Pairs
1	1	0	0	1
2	0	1	0	1
3	1	0	1	2
4	1	1	1	3
5	2	1	2	5
6	3	2	3	8
7	5	3	5	13
8	8	5	8	21
9	13	8	13	34
10	21	13	21	55
11	34	21	34	89
12	55	34	55	144
13	89	55	89	233

7. 233 pairs

8. From the third term on, each term equals the sum of the two preceding terms.

(Continued on next page.)

Activity 8 Continued**Box 3: LOOKING FOR PATTERNS**

Fibonacci #s	Sums	Fib. # – 1	Squares	Sum of Squares	Products	Quotients
1	1	0	1	1	1	1
1	2	0	1	2	2	2
2	4	1	4	6	6	1.5
3	7	2	9	15	15	1.666666667
5	12	4	25	40	40	1.6
8	20	7	64	104	104	1.625
13	33	12	169	273	273	1.615384615
21	54	20	441	714	714	1.619047619
34	88	33	1156	1870	1870	1.617647059
55	143	54	3025	4895	4895	1.618181818
89	232	88	7921	12816	12816	1.617977528
144	376	143	20736	33552	33552	1.618055556
233	609	232	54289	87841	87841	1.618025751
377	986	376	142129	229970	229970	1.618037135
610	1596	609	372100	602070	602070	1.618032787
987	2583	986	974169	1576239	1576239	1.618034448
1597	4180	1596	2550409	4126648	4126648	1.618033813
2584	6764	2583	6677056	10803704	10803704	1.618034056
4181	10945	4180	17480761	28284465	28284465	1.618033963
6765	17710	6764	45765225	74049690	0	0

1. c. The first sum is 1 less than the third Fibonacci number, the second sum is 1 less than the fourth Fibonacci number, and so on—the n th sum is 1 less than the $n + 2$ Fibonacci number.
2. c. The numbers in the two columns are the same.
d. The sum of the squares of the first n Fibonacci numbers equals the product of the n th Fibonacci number and the $n + 1$ Fibonacci number.
3. The quotients approach 1.61803.

Activity 9

- Answers will vary.
- See table. The number of layers doubles.

3.	Folds	0	1	2	3	4	5	6	7	8
	No. Layers (Std. Form)	1	2	4	8	16	32	64	128	256
	No. Layers (Fact. Form)	1	2	$2 \cdot 2$	$2 \cdot 2 \cdot 2$	$2 \cdot 2 \cdot 2 \cdot 2$	$2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$	$2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$	$2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$	$2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$
	No. Layers (Exp. Form)	2^0	2^1	2^2	2^3	2^4	2^5	2^6	2^7	2^8
	Approximate Height (cm)	0.008	0.016	0.031	0.063	0.125	0.25	0.5	1.0	2

- Answers will vary. The paper can be folded 5-6 times.
- 1 fold, 2^1 ; 0 folds 2^0
- See table.
- After 14 folds, the stack is 128 cm tall and after 15 folds it is 256 cm tall, so depending on the student's height, their answer will be 14 or 15 folds.
- The height of the stack after 30 folds would be about 8.4 km. This is higher than the Willis Tower (0.442 km) and less than the altitude of an orbiting satellite (altitudes range from about 150 km to 1000 km at the point nearest to Earth) and the distance to the moon (about 385,000 km).

Chapter 2

Introduction to Logic and Sets

Activity 1

Box 1

2. a. SRS, SRT, SRC, SRR (small, red, rhombus)
LRS, LRT, LRC, LRR
- b. 8
- c. 32
3. a. SGS, SGT, SGC, SGR, LGS, LGT, LGC, LGR,
SYS, SYT, SYC, SYR, LYS, LYT, LYC, LYR,
SBS, SBT, SBC, SBR, LBS, LBT, LBC, LBR
- b. The pieces are outside the loop.
4. a. Sample Responses: the set of blue pieces; the set of circles
- b. You could pair up the pieces in set A with the pieces in set B to make sure there are no pieces left over in either set. That is, you could create a one-to-one correspondence between the elements of the two sets.
- c. The set of pieces that are not large.

Activity 2

1. b. If the loops did not overlap, there would be no way to place the pieces that are Large AND Red.
2. a. (either) **RED or LARGE**
- b. 20
3. a. **RED and LARGE**
- b. 4
4. Sample Response: **Square and Circle**
5. Sample Response: **Triangle and Not Square** (The triangles are a subset of the pieces that are not square.)
6. a. 12
- b. A
7. **Not Square and Blue**

Since both loops contain pieces that are large and pieces that are small, the labels do not involve size. There are two possibilities for the label for the loop on the left, Not Yellow (since it contains pieces with all the other colors) or Not Square (since it contains only circles, squares, and rhombi). Suppose the correct label is Not Yellow. Then the small red square would go in that loop. Since it isn't, the label for the left loop cannot be Not Yellow. Thus the label must be Not Square. Similarly, the label for the right loop could be Blue or Not Rhombus. However, if the label was Not Rhombus, then the small red square would be in the loop. Since it isn't, the label must be Blue.

Chapter 3

Numeration Systems and Whole Number Operations

Activity 1

Box 1

14 tens

23 ones

13 ones

13 ones

1 hundreds

Sample Response: Writing a number in several different ways develops understanding of place value and of the regrouping (renaming) process used in computational algorithms.

Activity 2

1. 22 or 13
2. 53
3. 33 or 24
4. 222 or 141
5. 200, 110, 101, 20, 11 or 2
6. 3, 12, 21, or 30
7. 4, 13, 22, 31, or 40
8. 130, 121, 112, or 103

Activity 3

Box 2

1. A blue coin may be traded for four red coins. Then one of the red coins can be traded for four white coins.

Box 3

1. a. 2 blue, 3 red, 1 white b. 1 blue, 2 white c. 1 red, 3 white d. 2 red
2. a. 121_{four} b. 23_{four} c. 200_{four} d. 203_{four}
3.

1_{four}	2_{four}	3_{four}	10_{four}	11_{four}	12_{four}	13_{four}	20_{four}	21_{four}	22_{four}
23_{four}	30_{four}	31_{four}	32_{four}	33_{four}	100_{four}	101_{four}	102_{four}	103_{four}	110_{four}
111_{four}	112_{four}	113_{four}	120_{four}	121_{four}	122_{four}	123_{four}	130_{four}	...	
331_{four}	332_{four}	333_{four}	1000_{four}	1001_{four}	1002_{four}	1003_{four}	1010_{four}	1011_{four}	1012_{four}

Box 4

1. \$25
2. a. \$118
- b. 1102_{four}

(Continued on next page.)

Activity 3 Continued**Box 5**

- When you combine the three white coins used to make 23_{four} with the three white coins used to make 33_{four} , you can trade four of the coins for a red coin and have two white coins left over. The 1 represents this red coin. That is, $3_{\text{four}} + 3_{\text{four}} = 12_{\text{four}}$, so when adding, the 2 is recorded in the units place of the sum and the 1 (representing the trade for one red coin) is “carried” to the 10_{four} -column to be combined with (added to) the other red coins.
- 332_{four}
 - 1010_{four}
- To subtract the units digits, 1 four must be exchanged for units. This regrouping leaves 2 fours and 11_{four} (five) units.
 - After the regrouping in A, the units digits can be subtracted: $11_{\text{four}} - 2_{\text{four}} = 3_{\text{four}}$. The result is recorded in the units-place of the difference.
 - Finally, subtract the 4s-digits ($20_{\text{four}} - 10_{\text{four}} = 10_{\text{four}}$) and record the result (1) in the 4s-place.
- 112_{four}
 - 23_{four}

EXTENSIONS

- In both systems, the value of each digit in a numeral is determined by its position. In the decimal (base-ten) system, the value is a power of 10, and in the Fourian (base-four) system, it is a power of 4.
- The regrouping in Fourian addition and subtraction is identical to the regrouping in base-ten addition and subtraction except that the exchanges are for units, fours, sixteens, sixty-fours, and so on instead of units, tens, hundreds, etc.

Activity 4

- Sample Response: The word carry gives the impression that something is being moved from one place to another, but none of the numbers are moved in the addition algorithm. Instead, one of the addends is rewritten (renamed) in an equivalent form and the digits are regrouped according to their place-values before combining.

Box 5

The strategy used in each problem may vary.

- Compensation: $78 + 93 = (78 + 2) - (93 - 2) = 80 + 91 = 171$
- Decomposition: $413 + 289 = (400 + 200) + (10 + 80) + (3 + 9) = 600 + 90 + 12 = 702$
- Making Nice Numbers: $57 + 248 =$
 $57 + 250 = 307$ (2 was added to get 250, so subtract 2.)
 $307 - 2 = 305$
- Compensation: $544 + 58 = (544 + 6) + (58 - 6) = 550 + 52 = 602$
- Decomposition: $69 + 43 = (60 + 40) + (9 + 3) = 100 + 12 = 112$
- Making Nice Numbers: $2006 + 483 =$
 $2000 + 483 = 2483$ (6 was subtracted to get 2000, so add 6.)
 $2483 + 6 = 2489$

Activity 5

Answers will vary.