

Instructor's Solutions Manual
to accompany

Modern Physics, 4th Edition

Kenneth S. Krane
Department of Physics
Oregon State University

Copyright © 2020 John Wiley & Sons, Inc. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning or otherwise, except as permitted under Sections 107 or 108 of the 1976 United States Copyright Act, without prior written permission of the Publisher.

Chapter 1

1. (a) Conservation of momentum gives $p_{x,\text{initial}} = p_{x,\text{final}}$, or

$$m_{\text{H}} v_{\text{H},\text{initial}} + m_{\text{He}} v_{\text{He},\text{initial}} = m_{\text{H}} v_{\text{H},\text{final}} + m_{\text{He}} v_{\text{He},\text{final}}$$

Solving for $v_{\text{He},\text{final}}$ with $v_{\text{He},\text{initial}} = 0$, we obtain

$$\begin{aligned} v_{\text{He},\text{final}} &= \frac{m_{\text{H}}(v_{\text{H},\text{initial}} - v_{\text{H},\text{final}})}{m_{\text{He}}} \\ &= \frac{(1.674 \times 10^{-27} \text{ kg})[1.1250 \times 10^7 \text{ m/s} - (-6.724 \times 10^6 \text{ m/s})]}{6.646 \times 10^{-27} \text{ kg}} = 4.527 \times 10^6 \text{ m/s} \end{aligned}$$

(b) Kinetic energy is the only form of energy we need to consider in this elastic collision. Conservation of energy then gives $K_{\text{initial}} = K_{\text{final}}$, or

$$\frac{1}{2} m_{\text{H}} v_{\text{H},\text{initial}}^2 + \frac{1}{2} m_{\text{He}} v_{\text{He},\text{initial}}^2 = \frac{1}{2} m_{\text{H}} v_{\text{H},\text{final}}^2 + \frac{1}{2} m_{\text{He}} v_{\text{He},\text{final}}^2$$

Solving for $v_{\text{He},\text{final}}$ with $v_{\text{He},\text{initial}} = 0$, we obtain

$$\begin{aligned} v_{\text{He},\text{final}} &= \sqrt{\frac{m_{\text{H}}(v_{\text{H},\text{initial}}^2 - v_{\text{H},\text{final}}^2)}{m_{\text{He}}}} \\ &= \sqrt{\frac{(1.674 \times 10^{-27} \text{ kg})[(1.1250 \times 10^7 \text{ m/s})^2 - (-6.724 \times 10^6 \text{ m/s})^2]}{6.646 \times 10^{-27} \text{ kg}}} = 4.527 \times 10^6 \text{ m/s} \end{aligned}$$

2. (a) Let the helium initially move in the x direction. Then conservation of momentum gives:

$$\begin{aligned} p_{x,\text{initial}} &= p_{x,\text{final}} : & m_{\text{He}} v_{\text{He},\text{initial}} &= m_{\text{He}} v_{\text{He},\text{final}} \cos \theta_{\text{He}} + m_{\text{O}} v_{\text{O},\text{final}} \cos \theta_{\text{O}} \\ p_{y,\text{initial}} &= p_{y,\text{final}} : & 0 &= m_{\text{He}} v_{\text{He},\text{final}} \sin \theta_{\text{He}} + m_{\text{O}} v_{\text{O},\text{final}} \sin \theta_{\text{O}} \end{aligned}$$

From the second equation,

$$v_{\text{O},\text{final}} = -\frac{m_{\text{He}} v_{\text{He},\text{final}} \sin \theta_{\text{He}}}{m_{\text{O}} \sin \theta_{\text{O}}} = -\frac{(6.6465 \times 10^{-27} \text{ kg})(6.636 \times 10^6 \text{ m/s})(\sin 84.7^\circ)}{(2.6560 \times 10^{-26} \text{ kg})[\sin(-40.4^\circ)]} = 2.551 \times 10^6 \text{ m/s}$$

(b) From the first momentum equation,

$$\begin{aligned}
v_{\text{He,initial}} &= \frac{m_{\text{He}} v_{\text{He,final}} \cos \theta_{\text{He}} + m_{\text{O}} v_{\text{O,final}} \cos \theta_{\text{O}}}{m_{\text{He}}} \\
&= \frac{(6.6465 \times 10^{-27} \text{ kg})(6.636 \times 10^6 \text{ m/s})(\cos 84.7^\circ) + (2.6560 \times 10^{-26} \text{ kg})(2.551 \times 10^6 \text{ m/s})[\cos(-40.4^\circ)]}{6.6465 \times 10^{-27} \text{ kg}} \\
&= 8.376 \times 10^6 \text{ m/s}
\end{aligned}$$

3. (a) Using conservation of momentum for this one-dimensional situation, we have

$$p_{x,\text{initial}} = p_{x,\text{final}}, \text{ or}$$

$$m_{\text{He}} v_{\text{He}} + m_{\text{N}} v_{\text{N}} = m_{\text{D}} v_{\text{D}} + m_{\text{O}} v_{\text{O}}$$

Solving for v_{O} with $v_{\text{N}} = 0$, we obtain

$$v_{\text{O}} = \frac{m_{\text{He}} v_{\text{He}} - m_{\text{D}} v_{\text{D}}}{m_{\text{O}}} = \frac{(3.016 \text{ u})(6.346 \times 10^6 \text{ m/s}) - (2.014 \text{ u})(1.531 \times 10^7 \text{ m/s})}{15.003 \text{ u}} = -7.79 \times 10^5 \text{ m/s}$$

- (b) The kinetic energies are:

$$\begin{aligned}
K_{\text{initial}} &= \frac{1}{2} m_{\text{He}} v_{\text{He}}^2 + \frac{1}{2} m_{\text{N}} v_{\text{N}}^2 = \frac{1}{2} (3.016 \text{ u})(1.6605 \times 10^{-27} \text{ kg/u})(6.346 \times 10^6 \text{ m/s})^2 = 1.008 \times 10^{-13} \text{ J} \\
K_{\text{final}} &= \frac{1}{2} m_{\text{D}} v_{\text{D}}^2 + \frac{1}{2} m_{\text{O}} v_{\text{O}}^2 = \frac{1}{2} (2.014 \text{ u})(1.6605 \times 10^{-27} \text{ kg/u})(1.531 \times 10^7 \text{ m/s})^2 \\
&\quad + \frac{1}{2} (15.003 \text{ u})(1.6605 \times 10^{-27} \text{ kg/u})(7.79 \times 10^5 \text{ m/s})^2 = 3.995 \times 10^{-13} \text{ J}
\end{aligned}$$

As in Example 1.2, this is also a case in which nuclear energy turns into kinetic energy. The gain in kinetic energy is exactly equal to the loss in nuclear energy.

4. Let the two helium atoms move in opposite directions along the x axis with speeds v_1 and v_2 . Conservation of momentum along the x direction ($p_{x,\text{initial}} = p_{x,\text{final}}$) gives

$$0 = m_1 v_1 - m_2 v_2 \quad \text{or} \quad v_1 = v_2$$

The energy released is in the form of the total kinetic energy of the two helium atoms:

$$K_1 + K_2 = 92.2 \text{ keV}$$

Because $v_1 = v_2$, it follows that $K_1 = K_2 = 46.1 \text{ keV}$, so

$$\begin{aligned}
v &= \sqrt{\frac{2K_1}{m_1}} = \sqrt{\frac{2(46.1 \times 10^3 \text{ eV})(1.602 \times 10^{-19} \text{ J/eV})}{(4.00 \text{ u})(1.6605 \times 10^{-27} \text{ kg/u})}} = 1.49 \times 10^6 \text{ m/s} \\
v_2 &= v_1 = 1.49 \times 10^6 \text{ m/s}
\end{aligned}$$

5. (a) The kinetic energy of the electrons is

$$K_i = \frac{1}{2}mv_i^2 = \frac{1}{2}(9.11 \times 10^{-31} \text{ kg})(1.76 \times 10^6 \text{ m/s}) = 14.11 \times 10^{-19} \text{ J}$$

In passing through a potential difference of $\Delta V = V_f - V_i = +4.15 \text{ volts}$, the potential energy of the electrons changes by

$$\Delta U = q\Delta V = (-1.602 \times 10^{-19} \text{ C})(+4.15 \text{ V}) = -6.65 \times 10^{-19} \text{ J}$$

Conservation of energy gives $K_i + U_i = K_f + U_f$, so

$$K_f = K_i + (U_i - U_f) = K_i - \Delta U = 14.11 \times 10^{-19} \text{ J} + 6.65 \times 10^{-19} \text{ J} = 20.76 \times 10^{-19} \text{ J}$$

$$v_f = \sqrt{\frac{2K_f}{m}} = \sqrt{\frac{2(20.76 \times 10^{-19} \text{ J})}{9.11 \times 10^{-31} \text{ kg}}} = 2.13 \times 10^6 \text{ m/s}$$

- (b) In this case $\Delta V = -4.15 \text{ volts}$, so $\Delta U = +6.65 \times 10^{-19} \text{ J}$ and thus

$$K_f = K_i - \Delta U = 14.11 \times 10^{-19} \text{ J} - 6.65 \times 10^{-19} \text{ J} = 7.46 \times 10^{-19} \text{ J}$$

$$v_f = \sqrt{\frac{2K_f}{m}} = \sqrt{\frac{2(7.46 \times 10^{-19} \text{ J})}{9.11 \times 10^{-31} \text{ kg}}} = 1.28 \times 10^6 \text{ m/s}$$

6. (a) $\Delta x_A = v\Delta t_A = (0.624)(2.997 \times 10^8 \text{ m/s})(124 \times 10^{-9} \text{ s}) = 23.2 \text{ m}$

(b) $\Delta x_B = v\Delta t_B = (0.624)(2.997 \times 10^8 \text{ m/s})(159 \times 10^{-9} \text{ s}) = 29.7 \text{ m}$

7. With $T = 35^\circ\text{C} = 308 \text{ K}$ and $P = 1.22 \text{ atm} = 1.23 \times 10^5 \text{ Pa}$,

$$\frac{N}{V} = \frac{P}{kT} = \frac{1.23 \times 10^5 \text{ Pa}}{(1.38 \times 10^{-23} \text{ J/K})(308 \text{ K})} = 2.89 \times 10^{25} \text{ atoms/m}^3$$

so the volume available to each atom is $(2.89 \times 10^{25} \text{ m}^{-3})^{-1} = 3.46 \times 10^{-26} \text{ m}^3$. For a spherical atom, the volume would be

$$\frac{4}{3}\pi R^3 = \frac{4}{3}\pi(0.710 \times 10^{-10} \text{ m})^3 = 1.50 \times 10^{-30} \text{ m}^3$$

The fraction is then

$$\frac{1.50 \times 10^{-30}}{3.46 \times 10^{-26}} = 4.34 \times 10^{-5}$$

8. Differentiating $N(E)$ from Equation 1.23, we obtain

$$\frac{dN}{dE} = \frac{2N}{\sqrt{\pi}} \frac{1}{(kT)^{3/2}} \left[\frac{1}{2} E^{-1/2} e^{-E/kT} + E^{1/2} \left(-\frac{1}{kT} \right) e^{-E/kT} \right]$$

To find the maximum, we set this function equal to zero:

$$\frac{2N}{\sqrt{\pi}} \frac{1}{(kT)^{3/2}} E^{-1/2} e^{-E/kT} \left(\frac{1}{2} - \frac{E}{kT} \right) = 0$$

Solving, we find the maximum occurs at $E = \frac{1}{2} kT$. Note that $E = 0$ and $E = \infty$ also satisfy the equation, but these solutions give minima rather than maxima.

9. For this case $kT = (280 \text{ K})(8.617 \times 10^{-5} \text{ eV/K}) = 0.0241 \text{ eV}$. We take dE as the width of the interval (0.0012 eV) and E as its midpoint (0.0306 eV). Then

$$dN = N(E) dE = \frac{2N}{\sqrt{\pi}} \frac{1}{(0.0241 \text{ eV})^{3/2}} (0.0306 \text{ eV})^{1/2} e^{-(0.0306 \text{ eV})/(0.0241 \text{ eV})} (0.0012 \text{ eV}) = 1.8 \times 10^{-2} N$$

10. (a) From Eq. 1.33,

$$\Delta E_{\text{int}} = \frac{5}{2} nR \Delta T = \frac{5}{2} (2.37 \text{ moles})(8.315 \text{ J/mol} \cdot \text{K})(65.2 \text{ K}) = 3.21 \times 10^3 \text{ J}$$

(b) From Eq. 1.34,

$$\Delta E_{\text{int}} = \frac{7}{2} nR \Delta T = \frac{7}{2} (2.37 \text{ moles})(8.315 \text{ J/mol} \cdot \text{K})(65.2 \text{ K}) = 4.50 \times 10^3 \text{ J}$$

(c) For both cases, the change in the translational part of the kinetic energy is given by Eq. 1.31:

$$\Delta E_{\text{int}} = \frac{3}{2} nR \Delta T = \frac{3}{2} (2.37 \text{ moles})(8.315 \text{ J/mol} \cdot \text{K})(65.2 \text{ K}) = 1.93 \times 10^3 \text{ J}$$

11. After the collision, m_1 moves with speed v'_1 (in the y direction) and m_2 with speed v'_2 (at an angle θ with the x axis). Conservation of energy then gives $E_{\text{initial}} = E_{\text{final}}$:

$$\frac{1}{2} m_1 v_1^2 = \frac{1}{2} m_1 v_1'^2 + \frac{1}{2} m_2 v_2'^2 \quad \text{or} \quad v^2 = v_1'^2 + 3v_2'^2$$

Conservation of momentum gives:

$$p_{x,\text{initial}} = p_{x,\text{final}} : m_1 v_1 = m_2 v'_2 \cos \theta \quad \text{or} \quad v = 3v'_2 \cos \theta$$

$$p_{y,\text{initial}} = p_{y,\text{final}} : 0 = m_1 v'_1 - m_2 v'_2 \sin \theta \quad \text{or} \quad v'_1 = 3v'_2 \sin \theta$$

We first solve for the speeds by eliminating θ from these equations. Squaring the two momentum equations and adding them, we obtain $v^2 + v_1'^2 = 9v_2'^2$, and combining this result with the energy equation allows us to solve for the speeds:

$$v'_1 = v / \sqrt{2} \quad \text{and} \quad v'_2 = v / \sqrt{6}$$

By substituting this value of v'_2 into the first momentum equation, we obtain

$$\cos \theta = \sqrt{2/3} \quad \text{or} \quad \theta = 35.3^\circ$$

12. The combined particle, with mass $m' = m_1 + m_2 = 3m$, moves with speed v' at an angle θ with respect to the x axis. Conservation of momentum then gives:

$$p_{x,\text{initial}} = p_{x,\text{final}} : m_1 v_1 = m' v' \cos \theta \quad \text{or} \quad v = 3v' \cos \theta$$

$$p_{y,\text{initial}} = p_{y,\text{final}} : m_2 v_2 = m' v' \sin \theta \quad \text{or} \quad \frac{4}{3} v = 3v' \sin \theta$$

We can first solve for θ by dividing these two equations to eliminate the unknown v' :

$$\tan \theta = \frac{4}{3} \quad \text{or} \quad \theta = 53.1^\circ$$

Now we can substitute this result into either of the momentum equations to find

$$v' = 5v / 9$$

The kinetic energy lost is the difference between the initial and final kinetic energies:

$$K_{\text{initial}} - K_{\text{final}} = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2 - \frac{1}{2} m' v'^2 = \frac{1}{2} m v^2 + \frac{1}{2} (2m) \left(\frac{2}{3} v\right)^2 - \frac{1}{2} (3m) \left(\frac{5}{9} v\right)^2 = \frac{26}{27} \left(\frac{1}{2} m v^2\right)$$

The total initial kinetic energy is $\frac{1}{2} m v^2 + \frac{1}{2} (2m) \left(\frac{2}{3} v\right)^2 = \frac{17}{9} \left(\frac{1}{2} m v^2\right)$. The loss in kinetic energy is then $\frac{26}{51} = 51\%$ of the initial kinetic energy.

13. (a) Let v_1 represent the helium atom that moves in the $+x$ direction, and let v_2 represent the other helium atom (which might move either in the positive or negative x direction). Then conservation of momentum ($p_{x,\text{initial}} = p_{x,\text{final}}$) gives

$$m v = m_1 v_1 + m_2 v_2 \quad \text{or} \quad 2v = v_1 + v_2$$

where v_2 may be positive or negative. The initial velocity v is

$$v = \sqrt{\frac{2K}{m}} = \sqrt{\frac{2(40.0 \times 10^3 \text{ eV})(1.602 \times 10^{-19} \text{ J/eV})}{(8.00 \text{ u})(1.6605 \times 10^{-27} \text{ kg/u})}} = 9.822 \times 10^5 \text{ m/s}$$

The energy available to the two helium atoms after the decay is the initial kinetic energy of the beryllium atom plus the energy released in its decay:

$$K + 92.2 \text{ keV} = \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2v_2^2 = \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2(2v - v_1)^2$$

where the last substitution is made from the momentum equation. Solving this quadratic equation for v_1 , we obtain $v_1 = 2.47 \times 10^6 \text{ m/s}$ or $-0.508 \times 10^6 \text{ m/s}$.

Because we identified m_1 as the helium moving in the positive x direction, it is identified with the positive root and thus (because the two heliums are interchangeable in the equation) the second value represents the velocity of m_2 :

$$v_1 = 2.47 \times 10^6 \text{ m/s}, v_2 = -0.508 \times 10^6 \text{ m/s}$$

(b) Suppose we were to travel in the positive x direction at a speed of $v = 9.822 \times 10^5 \text{ m/s}$, which is the original speed of the beryllium from part (a). If we travel at the same speed as the beryllium, it appears to be at rest, so its initial momentum is zero in this frame of reference. The two heliums then travel with equal speeds in opposite directions along the x axis. Because they share the available energy equally, each helium has a kinetic energy of 46.1 keV and a speed of $\sqrt{2K/m} = 1.49 \times 10^6 \text{ m/s}$, as we found in Problem 4. Let's represent these velocities in this frame of reference as $v'_1 = +1.49 \times 10^6 \text{ m/s}$ and $v'_2 = -1.49 \times 10^6 \text{ m/s}$. Transforming back to the original frame, we find

$$v_1 = v'_1 + v = 1.49 \times 10^6 \text{ m/s} + 9.822 \times 10^5 \text{ m/s} = 2.47 \times 10^6 \text{ m/s}$$

$$v_2 = v'_2 + v = -1.49 \times 10^6 \text{ m/s} + 9.822 \times 10^5 \text{ m/s} = -0.508 \times 10^6 \text{ m/s}$$

14. (a) Let the second helium move in a direction at an angle θ with the x axis. (We'll assume that the 30° angle for m_1 is measured above the x axis, while the angle θ for m_2 is measured below the x axis. Then conservation of momentum gives:

$$p_{x,\text{initial}} = p_{x,\text{final}} : mv = m_1v_1 \cos 30^\circ + m_2v_2 \cos \theta \quad \text{or} \quad 2v - \frac{\sqrt{3}}{2}v_1 = v_2 \cos \theta$$

$$p_{y,\text{initial}} = p_{y,\text{final}} : 0 = m_1v_1 \sin 30^\circ - m_2v_2 \sin \theta \quad \text{or} \quad \frac{1}{2}v_1 = v_2 \sin \theta$$

We can eliminate the angle θ by squaring and adding the two momentum equations:

$$4v^2 + v_1^2 - 2\sqrt{3}vv_1 = v_2^2$$

The kinetic energy given to the two heliums is equal to the original kinetic energy $\frac{1}{2}mv^2$ of the beryllium plus the energy released in the decay:

$$\begin{aligned}\frac{1}{2}mv^2 + 92.2 \text{ keV} &= \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2v_2^2 = \frac{1}{2}m_1v_1^2 + \frac{1}{2}m_2(4v^2 + v_1^2 - 2\sqrt{3}vv_1) \\ \frac{1}{2}(m_1 + m_2)v_1^2 - \sqrt{3}m_2vv_1 + (2m_2v^2 - \frac{1}{2}mv^2 - 92.2 \text{ keV}) &= 0\end{aligned}$$

Solving this quadratic equation gives

$$v_1 = 2.405 \times 10^6 \text{ m/s}, -0.321 \times 10^6 \text{ m/s}$$

Based on the directions assumed in writing the momentum equations, only the positive root is meaningful. We can substitute this value for v_1 into either the momentum or the energy equations to find v_2 and so our solution is:

$$v_1 = 2.41 \times 10^6 \text{ m/s}, v_2 = 1.25 \times 10^6 \text{ m/s}$$

The angle θ can be found by substituting these values into either of the momentum equations, for example

$$\theta = \sin^{-1} \frac{v_1}{2v_2} = \sin^{-1} \frac{2.41 \times 10^6 \text{ m/s}}{2(1.25 \times 10^6 \text{ m/s})} = 74.9^\circ$$

(b) The original speed of the beryllium atom is $v = \sqrt{2K/m} = 1.203 \times 10^6 \text{ m/s}$. If we were to view the experiment from a frame of reference moving at this velocity, the original beryllium atom would appear to be at rest. In this frame of reference, in which the initial momentum is zero, the two helium atoms are emitted in opposite directions with equal speeds. Each helium has a kinetic energy of 46.1 keV and a speed of $v'_1 = v'_2 = 1.49 \times 10^6 \text{ m/s}$. Let ϕ represent the angle that each of the helium atoms makes with the x axis in this frame of reference. Then the relationship between the x components of the velocity of m_1 in this frame of reference and the original frame of reference is

$$v_1 \cos 30^\circ = v'_1 \cos \phi + v$$

and similarly for the y components

$$v_1 \sin 30^\circ = v'_1 \sin \phi$$

We can divide these two equations to get

$$\cot 30^\circ = \frac{v'_1 \cos \phi + v}{v'_1 \sin \phi}$$

which can be solved to give $\phi = 53.8^\circ$. Using this value of ϕ , we can then find $v_1 = 2.41 \times 10^6$ m/s. We can also write the velocity addition equations for m_2 :

$$v_2 \cos \theta = -v'_2 \cos \phi + v \quad \text{and} \quad v_2 \sin \theta = -v'_2 \sin \phi$$

which describe respectively the x and y components. Solving as we did for m_1 , we find $v_2 = 1.25 \times 10^6$ m/s and $\theta = 74.9^\circ$.

15. (a) With $K = \frac{3}{2}kT$,

$$\Delta K = \frac{3}{2}k\Delta T = \frac{3}{2}(1.38 \times 10^{-23} \text{ J/K})(80 \text{ K}) = 1.66 \times 10^{-21} \text{ J} = 0.0104 \text{ eV}$$

- (b) With $U = mgh$,

$$h = \frac{U}{mg} = \frac{1.66 \times 10^{-21} \text{ J}}{(40.0 \text{ u})(1.66 \times 10^{-27} \text{ kg/u})(9.80 \text{ m/s}^2)} = 2550 \text{ m}$$

16. We take dE to be the width of this small interval: $dE = 0.04kT - 0.02kT = 0.02kT$, and we evaluate the distribution function at an energy equal to the midpoint of the interval ($E = 0.03kT$):

$$\frac{dN}{N} = \frac{N(E)dE}{N} = \frac{2}{\sqrt{\pi}} \frac{1}{(kT)^{3/2}} (0.03kT)^{1/2} e^{-(0.03kT)/kT} (0.02kT) = 3.79 \times 10^{-3}$$

17. If we represent the molecule as two atoms considered as point masses m separated by a distance $2R$, the rotational inertia about one of the axes is $I_{x'} = mR^2 + mR^2 = 2mR^2$.

On average, the rotational kinetic energy about any one axis is $\frac{1}{2}kT$, so

$$\frac{1}{2}I_{x'}\omega_{x'}^2 = \frac{1}{2}kT \text{ and}$$

$$\omega_{x'} = \sqrt{\frac{kT}{I_{x'}}} = \sqrt{\frac{kT}{2mR^2}} = \sqrt{\frac{(1.38 \times 10^{-23} \text{ J/K})(300 \text{ K})}{2(15.995 \text{ u})(1.6605 \times 10^{-27} \text{ kg/u})(0.0605 \times 10^{-9} \text{ m})^2}} = 4.61 \times 10^{12} \text{ rad/s}$$

Chapter 2

1. Your air speed in still air is $(750 \text{ km})/(3.14 \text{ h}) = 238.8 \text{ km/h}$. With the nose of the plane pointed 22° west of north, you would be traveling at this speed in that direction if there were no wind. With the wind blowing, you are actually traveling due north at an effective speed of $(750 \text{ km})/(4.32 \text{ h}) = 173.6 \text{ km/h}$. The wind must therefore have a north-south component of $(238.8 \text{ km/h})(\cos 22^\circ) - 173.6 \text{ km/h} = 47.8 \text{ km/h}$ (toward the south) and an east-west component of $(238.8 \text{ km/h})(\sin 22^\circ) = 89.5 \text{ km/h}$ (toward the east). The wind speed is thus

$$v = \sqrt{(47.8 \text{ km/h})^2 + (89.5 \text{ km/h})^2} = 101 \text{ km/h}$$

in a direction that makes an angle of

$$\theta = \tan^{-1} \frac{89.5 \text{ km/h}}{47.8 \text{ km/h}} = 62^\circ \text{ east of south}$$

2. (a) $\frac{95 \text{ m}}{0.53 \text{ m/s}} = 179 \text{ s}$

(b) $\frac{95 \text{ m}}{1.24 \text{ m/s} + 0.53 \text{ m/s}} = 54 \text{ s}$

(c) $\frac{95 \text{ m}}{2.48 \text{ m/s} - 0.53 \text{ m/s}} = 49 \text{ s}$

3. $\Delta t = t_{\text{up}} + t_{\text{down}} - 2t_{\text{across}} = \frac{2L}{c} \left[\frac{1}{1 - u^2/c^2} - \frac{1}{\sqrt{1 - u^2/c^2}} \right]$

Assuming $u \ll c$,

$$\frac{1}{1 - u^2/c^2} \cong 1 + \frac{u^2}{c^2} \quad \text{and} \quad \frac{1}{\sqrt{1 - u^2/c^2}} \cong 1 + \frac{1}{2} \frac{u^2}{c^2}$$

$$\Delta t \cong \frac{2L}{c} \left[1 + \frac{u^2}{c^2} - \left(1 + \frac{1}{2} \frac{u^2}{c^2} \right) \right] = \frac{Lu^2}{c^3}$$

$$u = \sqrt{\frac{c^3 \Delta t}{L}} = \sqrt{\frac{(3 \times 10^8 \text{ m/s})^3 (2 \times 10^{-15} \text{ s})}{11 \text{ m}}} = 7 \times 10^4 \text{ m/s}$$

4. (a) $u = 100 \text{ km/h} = 28 \text{ m/s} \ll c$

$$\sqrt{1 - u^2 / c^2} \cong 1 - \frac{1}{2} \frac{u^2}{c^2} = 1 - \frac{1}{2} \frac{(28 \text{ m/s})^2}{(3 \times 10^8 \text{ m/s})^2} = 1 - 4.3 \times 10^{-15}$$

$$L = L_0 \sqrt{1 - u^2 / c^2} = L_0 (1 - 4.3 \times 10^{-15})$$

$$L_0 - L = (4.3 \times 10^{-15}) L_0 = (4.3 \times 10^{-15})(4 \times 10^6 \text{ m}) = 1.7 \times 10^{-8} \text{ m}$$

This is less than the wavelength of light.

(b) $\Delta t = \frac{\Delta t_0}{\sqrt{1 - u^2 / c^2}} \cong \frac{\Delta t_0}{1 - 4.3 \times 10^{-15}} \cong \Delta t_0 (1 + 4.3 \times 10^{-15})$

$$\Delta t - \Delta t_0 = (4.3 \times 10^{-15}) \Delta t_0 = (4.3 \times 10^{-15})(40 \text{ h})(3600 \text{ s/h}) = 6.2 \times 10^{-10} \text{ s}$$

5. With $L = \frac{1}{2} L_0$, the length contraction formula gives $\frac{1}{2} L_0 = L_0 \sqrt{1 - u^2 / c^2}$, so

$$u = \sqrt{3/4} c = 2.6 \times 10^8 \text{ m/s}$$

6. The astronaut must travel 600 light-years at a speed close to the speed of light and must age only 12 years. To an Earth-bound observer, the trip takes about $\Delta t = 600$ years, but this is a dilated time interval; in the astronaut's frame of reference, the elapsed time is the

proper time interval Δt_0 of 12 years. Thus, with $\Delta t = \Delta t_0 / \sqrt{1 - u^2 / c^2}$,

$$600 \text{ years} = \frac{12 \text{ years}}{\sqrt{1 - u^2 / c^2}} \quad \text{or} \quad 1 - \frac{u^2}{c^2} = \left(\frac{12}{600} \right)^2$$

$$u = \sqrt{1 - (12 / 600)^2} c = 0.9998c$$

7. (a) $\Delta t = \frac{\Delta t_0}{\sqrt{1 - u^2 / c^2}} = \frac{120.0 \text{ ns}}{\sqrt{1 - (0.950)^2}} = 384 \text{ ns}$

(b) $d = v \Delta t = 0.950(3.00 \times 10^8 \text{ m/s})(384 \times 10^{-9} \text{ s}) = 109 \text{ m}$

(c) $d_0 = v \Delta t_0 = 0.950(3.00 \times 10^8 \text{ m/s})(120.0 \times 10^{-9} \text{ s}) = 34.2 \text{ m}$

8. In the laboratory reference frame, the lifetime is

$$\Delta t = \frac{d}{u} = \frac{1.15 \text{ mm}}{0.993(3.00 \times 10^8 \text{ m/s})} = 0.386 \times 10^{-11} \text{ s}$$

$$\Delta t_0 = \Delta t \sqrt{1 - u^2 / c^2} = (0.386 \times 10^{-11} \text{ s}) \sqrt{1 - (0.993)^2} = 4.56 \times 10^{-13} \text{ s}$$

9. From Equation 2.15, $\Delta t_1 = L / (v - u)$, and from Equation 2.16, $\Delta t_2 = L / (c + u)$.

$$\Delta t = \Delta t_1 + \Delta t_2 = \frac{L}{v - u} + \frac{L}{c + u} = \frac{\Delta t_0}{\sqrt{1 - u^2 / c^2}}$$

With $\Delta t_0 = L_0 / v' + L_0 / c$ and $L = L_0 \sqrt{1 - u^2 / c^2}$, this becomes

$$\frac{1}{v - u} + \frac{1}{c + u} = \frac{1}{1 - u^2 / c^2} \left(\frac{1}{v'} + \frac{1}{c} \right)$$

Solving for v , we obtain

$$v = \frac{v' + u}{1 + v'u / c^2}$$

10. Let ship A represent observer O , and let observer O' be on Earth. Then $v' = 0.831c$ and $u = -0.743c$, and so

$$v = \frac{v' + u}{1 + v'u / c^2} = \frac{0.831c + 0.743c}{1 + (0.831)(0.743)} = 0.973c$$

If now ship B represents observer O , then $v' = -0.743c$ and $u = -0.831c$.

$$v = \frac{v' + u}{1 + v'u / c^2} = \frac{-0.743c - 0.831c}{1 + (-0.743)(-0.831)} = -0.973c$$

11. Let O' be the observer on the space station, and let O be the observer on ship B . Then $v' = 0.811c$ and $u = -0.665c$.

$$v = \frac{v' + u}{1 + v'u / c^2} = \frac{0.811c - 0.665c}{1 + (0.811)(-0.665)} = 0.317c$$

12. (a) With $f' = f \sqrt{(1-u/c)/(1+u/c)}$ and $\lambda = c/f$, we obtain

$$\lambda' = \lambda \sqrt{\frac{1+u/c}{1-u/c}} \quad \text{or} \quad 366 \text{ nm} = 122 \text{ nm} \sqrt{\frac{1+u/c}{1-u/c}}$$

Solving, we get $u/c = 0.800$ or $u = 2.40 \times 10^8 \text{ m/s}$.

$$(b) \lambda' = \lambda \sqrt{\frac{1-u/c}{1+u/c}} = 122 \text{ nm} \sqrt{\frac{1-0.800}{1+0.800}} = 40.7 \text{ nm}$$

13. With $f' = f \sqrt{(1-u/c)/(1+u/c)}$ and $\lambda = c/f$, we obtain

$$\frac{1-u/c}{1+u/c} = \left(\frac{f'}{f} \right)^2 = \left(\frac{\lambda}{\lambda'} \right)^2 = \left(\frac{650 \text{ nm}}{550 \text{ nm}} \right)^2 = 1.397$$

Solving, $u/c = 0.166$ or $u = 5.0 \times 10^7 \text{ m/s}$.

14. According to O , the rest length L_0 of the hypotenuse is shortened to

$$L = L_0 \sqrt{1-u^2/c^2} = L_0 \sqrt{1-(0.92)^2} = 0.392L_0$$

The height of the triangle, which is $0.5L_0$, is not affected by the length contraction. The base angles of the triangle are then

$$\theta = \tan^{-1} \left(\frac{0.5L_0}{\frac{1}{2} \times 0.392L_0} \right) = \tan^{-1}(2.55) = 68.6^\circ$$

The apex angle is then $180^\circ - 2(68.6^\circ) = 42.8^\circ$.

15. In the direction of motion, the radius of the gold nucleus is reduced to

$$(7.0 \text{ fm}) \sqrt{1-(0.99995)^2} = 0.070 \text{ fm}$$

At this speed the formerly spherical gold nucleus looks like a pancake whose thickness is only 1% that of the original sphere and thus whose density has been increased by a factor of 100. The collision of two such compressed, high-energy nuclei is thought to create for a fraction of a second the conditions of density and temperature that occurred in the very early universe.

$$16. \quad dx' = \frac{dx - u dt}{\sqrt{1 - u^2/c^2}} \quad \text{and} \quad dt' = \frac{dt - u dx/c^2}{\sqrt{1 - u^2/c^2}}$$

$$v'_x = \frac{dx'}{dt'} = \frac{dx - u dt}{dt - u dx/c^2} = \frac{dx/dt - u}{1 - u(dx/dt)/c^2} = \frac{v_x - u}{1 - uv_x/c^2}$$

With $dz' = dz$, we obtain

$$v'_z = \frac{dz'}{dt'} = \frac{dz}{(dt - u dx/c^2)/\sqrt{1 - u^2/c^2}} = \frac{(dz/dt)\sqrt{1 - u^2/c^2}}{1 - uv_x/c^2} = \frac{v_z\sqrt{1 - u^2/c^2}}{1 - uv_x/c^2}$$

17. For the light beam, observer O measures $v_x = 0$, $v_y = c$. Observer O' measures

$$v'_x = \frac{v_x - u}{1 - uv_x/c^2} = 0 - u = -u \quad \text{and} \quad v'_y = \frac{v_y\sqrt{1 - u^2/c^2}}{1 - uv_x/c^2} = c\sqrt{1 - u^2/c^2}$$

According to O' , the speed of the light beam is

$$v' = \sqrt{(v'_x)^2 + (v'_y)^2} = \sqrt{u^2 + c^2(1 - u^2/c^2)} = c$$

18. O measures times t_1 and t_2 for the beginning and end of the interval, while O' measures t'_1 and t'_2 . Using Equation 2.23d,

$$t'_1 = \frac{t_1 - ux/c^2}{\sqrt{1 - u^2/c^2}} \quad \text{and} \quad t'_2 = \frac{t_2 - ux/c^2}{\sqrt{1 - u^2/c^2}}$$

The same coordinate x appears in both expressions, because the bulb is at rest according to O (so Δt is the proper time interval). Subtracting these two equations, we obtain

$$t'_2 - t'_1 = \frac{t_2 - t_1}{\sqrt{1 - u^2/c^2}} \quad \text{or} \quad \Delta t' = \frac{\Delta t}{\sqrt{1 - u^2/c^2}}$$

19. Suppose observer O is moving with the K meson; to this observer, the K meson appears to be at rest, and so O measures $v_1 = +0.815c$ and $v_2 = -0.815c$ for the two π mesons. Observer O' is moving relative to O with a velocity $u = -0.453c$; in the reference frame of O' , observer O and the K meson are moving in the positive x direction with a velocity of $0.453c$. We can use the Lorentz velocity transformation (Equation 2.28a) to find the velocities of the two π mesons according to O' :

$$v'_1 = \frac{v_1 - u}{1 - v_1 u / c^2} = \frac{+0.815c - (-0.453c)}{1 - (0.815)(-0.453)} = +0.926c$$

$$v'_2 = \frac{v_2 - u}{1 - v_2 u / c^2} = \frac{-0.815c - (-0.453c)}{1 - (-0.815)(-0.453)} = -0.575c$$

20. Imagine the rod to be the hypotenuse of a right triangle having sides L_x along the x axis and L_y in the y direction. According to O' , the length L_x is shortened by the length contraction, but the length L_y is unaffected because it is perpendicular to the direction of motion. For O , $L_y = L_x \tan 34^\circ$, while for O' , $L'_y = L'_x \tan 52^\circ$ where $L'_x = L_x \sqrt{1 - u^2 / c^2}$. Because $L_y = L'_y$, we have

$$L_x \tan 34^\circ = L'_x \tan 52^\circ = L_x \sqrt{1 - u^2 / c^2} \tan 52^\circ$$

or

$$u = c \sqrt{1 - (\tan 34^\circ)^2 / (\tan 52^\circ)^2} = 0.850c$$

21. (a) Changing the coordinates in Equation 2.23d to intervals, we have

$$\Delta t' = \frac{\Delta t - u \Delta x / c^2}{\sqrt{1 - u^2 / c^2}} = \frac{0.528 \mu s - (0.685)(49.5 \text{ m}) / (300 \text{ m}/\mu s)}{\sqrt{1 - (0.685)^2}} = +0.570 \mu s$$

- (b) Changing coordinates to intervals in Equation 2.23a,

$$\Delta x' = \frac{\Delta x - u \Delta t}{\sqrt{1 - u^2 / c^2}} = \frac{49.5 \text{ m} - (0.685 \times 300 \text{ m}/\mu s)(0.528 \mu s)}{\sqrt{1 - (0.685)^2}} = -81.0 \text{ m}$$

The negative sign of $\Delta x'$ indicates that O' finds the two events in inverted locations compared with O ; for example, if O finds that event 1 occurs at a smaller x coordinate than event 2, then O' finds that event 1 occurs at a *larger* x' coordinate than event 2. That is, O sees event 1 to the left of event 2, while O' sees event 1 to the right of event 2. Note that both observers find the time interval to be positive – event 2 occurs after event 1 to both observers.

22. From Equation 2.23d written in terms of intervals, for O' to find $\Delta t' = 0$, it must be true that $\Delta t - (u / c^2) \Delta x = 0$. Thus

$$u = \frac{c^2 \Delta t}{\Delta x} = c \frac{(300 \text{ m}/\mu\text{s})(0.138 \mu\text{s} - 0.124 \mu\text{s})}{(23.6 \text{ m} - 10.4 \text{ m})} = +0.32c$$

23. (a) In your frame of reference on the plane, the distance between Los Angeles and Boston is contracted to $(3000 \text{ mi})\sqrt{1 - [(600 \text{ mi/h})/(1000 \text{ mi/h})]^2} = 2400 \text{ mi}$, and you measure the time for the trip to be $(2400 \text{ mi})/(600 \text{ mi/h}) = 4 \text{ h}$. So your watch reads 2:00 pm.
 (b) According to an observer on the ground, the trip takes $(3000 \text{ mi})/(600 \text{ mi/h}) = 5 \text{ h}$. So the airport clock reads 3:00 pm.
 (c) The return trip again takes 4 h as measured in your frame of reference and 5 h from the frame of reference of the ground. When you depart Boston, the airport clock reads 10:00 am but your watch reads 9:00 am, so when you land in Los Angeles, your watch reads 1:00 pm and the airport clock reads 3:00 pm.

This analysis duplicates the result of the twin paradox: If your twin wearing an identical watch had remained at the Los Angeles airport, all observers would agree that you were 2 hours younger than your twin.

24. (a) On the outward journey at $0.60c$, the rate at which signals are received is

$$f' = f \sqrt{\frac{1 - u/c}{1 + u/c}} = (1/\text{year}) \sqrt{\frac{1 - 0.60}{1 + 0.60}} = 0.5/\text{year}$$

- (b) During the return journey,

$$f' = f \sqrt{\frac{1 + u/c}{1 - u/c}} = (1/\text{year}) \sqrt{\frac{1 + 0.60}{1 - 0.60}} = 2/\text{year}$$


- (c) According to Casper, Amelia's outward journey lasts 16 years (it is 16 years before he sees her arrive at the planet), during which he receives 8 signals ($0.5/\text{year} \times 16 \text{ years}$). Her total journey lasts 20 Earth years, so the return journey lasts 4 Earth years, during which he receives 8 signals ($2/\text{year} \times 4 \text{ years}$). Thus Casper receives 16 signals (8 during the outward trip and 8 during the return) and concludes that his sister has aged 16 years.

25. According to Amelia, the distance to the star is shortened to

$$L = L_0 \sqrt{1 - v^2/c^2} = (8.0 \text{ l-y}) \sqrt{1 - (0.80)^2} = 4.8 \text{ l-y}$$

and at a speed of $0.80c$ Amelia's travel time to the star is $(4.8 \text{ l-y})/(0.80c) = 6.0 \text{ y}$. The total round-trip time in Amelia's frame of reference is 12 years, so she is 8 years younger than her brother when she returns.

26. (a)



(b) 16 years

(c) 4 years

27. (a) To an Earth-bound observer Alice's round trip takes 20 years each way ($20 \text{ years} \times 0.6c = 12 \text{ light-years}$) for a total time of 40 years. Bob's travel time is 15 years each way ($15 \text{ years} \times 0.8c = 12 \text{ light-years}$) for a total travel time of 30 years. With Bob's 10-year delay in departing, the two arrive on Earth simultaneously.
 (b) To Alice, the distance to the star is contracted to

$$L = L_0 \sqrt{1 - v^2 / c^2} = 12 \text{ light-years} \sqrt{1 - (0.6)^2} = 9.6 \text{ light-years}$$

So in Alice's frame of reference the trip takes a time of $(9.6 \text{ light years}) / 0.6c = 16 \text{ years}$ each way. To Bob, the distance to the star is

$$L = L_0 \sqrt{1 - v^2 / c^2} = 12 \text{ light-years} \sqrt{1 - (0.8)^2} = 7.2 \text{ light-years}$$

and in Bob's frame the travel time is $(7.2 \text{ light-years})/0.8c = 9 \text{ years}$ each way. Relative to Alice's original departure time, Alice has aged 32 years while Bob has aged $10 + 18 = 28 \text{ years}$. So Bob is younger by 4 years.

28. (a) Suppose Agnes travels at speed v . Then in her reference frame the distance to the star is shortened to $L = L_0 \sqrt{1 - v^2 / c^2}$, so the time for her one-way trip is L/v and thus

$$\frac{16 \text{ light-years} \sqrt{1 - v^2 / c^2}}{v} = 10 \text{ y} \quad \text{or} \quad \sqrt{\frac{c^2}{v^2} - 1} = \frac{10}{16}$$

Solving, we find $v = 0.848c$.

(b) According to Bert, Agnes traveled on a journey of 32 light-years at a speed of $0.848c$ which corresponds to a time of $(32 \text{ light-years})/0.848c = 37.7 \text{ years}$.

29. (a)

$$\begin{aligned} K'_i &= K'_{li} + K'_{2i} = \frac{m_1 c^2}{\sqrt{1 - v_{li}^2 / c^2}} - m_1 c^2 + \frac{m_2 c^2}{\sqrt{1 - v_{2i}^2 / c^2}} - m_2 c^2 \\ &= \frac{(2m)c^2}{\sqrt{1 - 0^2}} - (2m)c^2 + \frac{mc^2}{\sqrt{1 - (0.750)^2}} - mc^2 = 0.512mc^2 \\ K'_f &= K'_{lf} + K'_{2f} = \frac{m_1 c^2}{\sqrt{1 - v_{lf}^2 / c^2}} - m_1 c^2 + \frac{m_2 c^2}{\sqrt{1 - v_{2f}^2 / c^2}} - m_2 c^2 \\ &= \frac{(2m)c^2}{\sqrt{1 - (-0.585)^2}} - (2m)c^2 + \frac{mc^2}{\sqrt{1 - (0.294)^2}} - mc^2 = 0.512mc^2 \end{aligned}$$

- (b)

$$\begin{aligned} K_i &= K_{li} + K_{2i} = \frac{m_1 c^2}{\sqrt{1 - v_{li}^2 / c^2}} - m_1 c^2 + \frac{m_2 c^2}{\sqrt{1 - v_{2i}^2 / c^2}} - m_2 c^2 \\ &= \frac{(2m)c^2}{\sqrt{1 - (0.550)^2}} - (2m)c^2 + \frac{mc^2}{\sqrt{1 - (-0.340)^2}} - mc^2 = 0.458mc^2 \\ K_f &= K_{lf} + K_{2f} = \frac{m_1 c^2}{\sqrt{1 - v_{lf}^2 / c^2}} - m_1 c^2 + \frac{m_2 c^2}{\sqrt{1 - v_{2f}^2 / c^2}} - m_2 c^2 \\ &= \frac{(2m)c^2}{\sqrt{1 - (-0.051)^2}} - (2m)c^2 + \frac{mc^2}{\sqrt{1 - (0.727)^2}} - mc^2 = 0.458mc^2 \end{aligned}$$

30.
$$p = \frac{mv}{\sqrt{1 - v^2 / c^2}} = \frac{1}{c} \frac{(mc^2)(v/c)}{\sqrt{1 - v^2 / c^2}} = \frac{1}{c} \frac{(938.3 \text{ MeV})(0.835)}{\sqrt{1 - (0.835)^2}} = 1424 \text{ MeV}/c$$

$$K = \frac{mc^2}{\sqrt{1-v^2/c^2}} - mc^2 = \frac{938.3 \text{ MeV}}{\sqrt{1-(0.835)^2}} - 938.3 \text{ MeV} = 767 \text{ MeV}$$

$$E = K + mc^2 = 767 \text{ MeV} + 938.3 \text{ MeV} = 1705 \text{ MeV}$$

31. $E = K + mc^2 = 0.923 \text{ MeV} + 0.511 \text{ MeV} = 1.434 \text{ MeV}$

Solving Equation 2.36 for v , we obtain

$$v = c \sqrt{1 - \left(\frac{mc^2}{E} \right)^2} = c \sqrt{1 - \left(\frac{0.511 \text{ MeV}}{1.434 \text{ MeV}} \right)^2} = 0.934c$$

32.
$$W = \int F dx = \int \frac{dp}{dt} dx = \int dp \frac{dx}{dt} = \int v dp$$

$$K = \int_0^v v dp = pv - \int_0^v p dv = \frac{mv^2}{\sqrt{1-v^2/c^2}} - \int_0^v \frac{mv}{\sqrt{1-v^2/c^2}} dv$$

$$= \frac{mv^2}{\sqrt{1-v^2/c^2}} + mc^2 \sqrt{1-v^2/c^2} - mc^2 = \frac{mc^2}{\sqrt{1-v^2/c^2}} - mc^2$$

33. For what range of velocities is $K - \frac{1}{2}mv^2 \leq 0.01K$? At the upper limit of this range, where $K - \frac{1}{2}mv^2 = 0.01K$, we have

$$0.99K = 0.99 \left(\frac{mc^2}{\sqrt{1-v^2/c^2}} - mc^2 \right) = \frac{1}{2}mv^2$$

With $x = v^2/c^2$, $0.99 \left(\frac{1}{\sqrt{1-x}} - 1 \right) = \frac{1}{2}x$ which gives $\frac{1}{1-x} = \left(1 + \frac{0.5}{0.99}x \right)^2$

$$1 = (1-x)(1 + 1.0101x + 0.2551x^2) \quad \text{or} \quad 0.2551x^2 + 0.7550x - 0.0101 = 0$$

Solving using the quadratic formula, we find $x = 0.0133$ or -2.97 . Only the positive solution is physically meaningful, so

$$v = \sqrt{0.0133}c = 0.115c$$

That is, for speeds smaller than $0.115c$, the classical kinetic energy is accurate to within 1%. For a different approach to that same type of calculation, see Problem 36.

34. As in Problem 33, let us now find the *lower* limit on the momentum such that

$$\sqrt{(pc)^2 + (mc^2)^2} - pc \leq 0.01\sqrt{(pc)^2 + (mc^2)^2}$$

From the lower limit, we obtain $0.99\sqrt{(pc)^2 + (mc^2)^2} = pc$, which can be written as

$$(pc)^2 = \frac{m^2 c^4}{1/(0.99)^2 - 1} \quad \text{or} \quad pc = 7.02mc^2$$

With $mvc / \sqrt{1 - v^2 / c^2} = 7.02mc^2$, we obtain

$$\frac{v^2}{c^2} = 49.25 \left(1 - \frac{v^2}{c^2} \right) \quad \text{or} \quad v/c = 0.990$$

Whenever $v/c \geq 0.990$, the expression $E = pc$ will be accurate to within 1%.

$$35. \quad E^2 = \frac{(mc^2)^2}{1 - v^2 / c^2} = (mc^2)^2 \left(\frac{1 - v^2 / c^2 + v^2 / c^2}{1 - v^2 / c^2} \right) = (mc^2)^2 + \frac{m^2 c^2 v^2}{1 - v^2 / c^2} = (mc^2)^2 + (pc)^2$$

$$E = \sqrt{(mc^2)^2 + (pc)^2}$$

36. With $(1+x)^n = 1 + nx + n(n-1)x^2/2!$ we have

$$\frac{1}{\sqrt{1 - v^2 / c^2}} = 1 + \frac{1}{2} \frac{v^2}{c^2} + \frac{(-1/2)(-3/2)}{2} \left(\frac{v^2}{c^2} \right)^2 + \dots$$

and so

$$K = mc^2 \left(\frac{1}{\sqrt{1 - v^2 / c^2}} - 1 \right) = mc^2 \left(1 + \frac{1}{2} \frac{v^2}{c^2} + \frac{3}{8} \frac{v^4}{c^4} + \dots - 1 \right) = \frac{1}{2} mv^2 \left(1 + \frac{3}{4} \frac{v^2}{c^2} + \dots \right)$$

The correction term is $3v^2/4c^2$, which has the value 0.1% when $3v^2/4c^2 = 0.001$, or

$$v = \sqrt{0.001(4/3)} c = 0.0365c$$

37. (a) With $E = 1351$ MeV and $pc = 1256$ MeV, Equation 2.39 gives

$$m = \frac{1}{c^2} \sqrt{E^2 - (pc)^2} = \frac{1}{c^2} \sqrt{(1351 \text{ MeV})^2 - (1256 \text{ MeV})^2} = 498 \text{ MeV}/c^2$$

$$(b) \quad E = \sqrt{(pc)^2 + (mc^2)^2} = \sqrt{(857 \text{ MeV})^2 + (498 \text{ MeV})^2} = 991 \text{ MeV}$$

38.

$$\begin{aligned} K_f - K_i &= (E_f - mc^2) - (E_i - mc^2) = E_f - E_i \\ &= \frac{mc^2}{\sqrt{1 - v_f^2/c^2}} - \frac{mc^2}{\sqrt{1 - v_i^2/c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1 - (0.91)^2}} - \frac{0.511 \text{ MeV}}{\sqrt{1 - (0.85)^2}} = 0.262 \text{ MeV} \end{aligned}$$

39.

$$\Delta E = mc \Delta T = (1 \text{ g})(0.40 \text{ J/g} \cdot \text{K})(100 \text{ K}) = 40 \text{ J}$$

$$\Delta m = \frac{\Delta E}{c^2} = \frac{40 \text{ J}}{9 \times 10^{16} \text{ m}^2/\text{s}^2} = 4.4 \times 10^{-16} \text{ kg}$$

40. (a) At such low speed, the classical approximation is valid.

$$K = \frac{1}{2}mv^2 = \frac{1}{2}mc^2 \left(\frac{v^2}{c^2} \right) = \frac{1}{2}(0.511 \text{ MeV})(1.00 \times 10^{-4})^2 = 2.56 \times 10^{-3} \text{ eV}$$

(b) The relativistic expression gives

$$K = mc^2 \left(\frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right) = 0.511 \text{ MeV} \left(\frac{1}{\sqrt{1 - (0.01)^2}} - 1 \right) = 25.6 \text{ eV}$$

For this speed, the classical expression $\frac{1}{2}mv^2$ also gives 25.6 eV, so the two calculations agree to at least three significant figures. (Actually they agree to four significant figures, but not to five.)

(c) The relativistic expression gives

$$K = mc^2 \left(\frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right) = 0.511 \text{ MeV} \left(\frac{1}{\sqrt{1 - (0.3)^2}} - 1 \right) = 24.7 \text{ keV}$$

For this speed, the classical expression gives 23.0 keV, which is incorrect by about 7%.

$$(d) \quad K = mc^2 \left(\frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right) = 0.511 \text{ MeV} \left(\frac{1}{\sqrt{1 - (0.999)^2}} - 1 \right) = 10.9 \text{ MeV}$$

41. Because the electrons and the protons have charges of the same magnitude e , after acceleration through a potential difference of magnitude $\Delta V = 12.0$ million volts (a positive difference for the electron, a negative difference for the proton), each loses potential energy of $\Delta U = -e\Delta V = -12.0$ MeV and thus each acquires a kinetic energy of $K = +12.0$ MeV. For the electron, $E = K + mc^2 = 12.0$ MeV + 0.511 MeV = 12.5 MeV. The momentum is then

$$p = \frac{1}{c} \sqrt{E^2 - (mc^2)^2} = \frac{1}{c} \sqrt{(12.5 \text{ MeV})^2 - (0.511 \text{ MeV})^2} = 12.5 \text{ MeV}/c$$

The classical formula $K = p^2 / 2m$ gives

$$p = \sqrt{2mK} = \sqrt{2(0.511 \text{ MeV}/c^2)(12.0 \text{ MeV})} = 3.50 \text{ MeV}/c$$

which is far from the correct result (a discrepancy we would expect for such highly relativistic electrons). For the protons, $E = K + mc^2 = 12.0$ MeV + 938.3 MeV = 950.3 MeV, and the momentum is

$$p = \frac{1}{c} \sqrt{E^2 - (mc^2)^2} = \frac{1}{c} \sqrt{(950.3 \text{ MeV})^2 - (938.3 \text{ MeV})^2} = 150.5 \text{ MeV}/c$$

The classical formula gives

$$p = \sqrt{2mK} = \sqrt{2(938.3 \text{ MeV}/c^2)(12.0 \text{ MeV})} = 150.1 \text{ MeV}/c$$

The difference between the classical and relativistic formulas appears only in the fourth significant figure.

42. The mass of a uranium atom is about $(235 \text{ u})(1.66 \times 10^{-27} \text{ kg/u}) = 3.90 \times 10^{-25} \text{ kg}$, so 1.50 kg contains $1.50 \text{ kg} / 3.90 \times 10^{-25} \text{ kg} = 3.84 \times 10^{24}$ atoms. The total energy released is

$$\Delta E = (210 \text{ MeV/atom})(3.84 \times 10^{24} \text{ atoms}) = 8.06 \times 10^{26} \text{ MeV}$$

and the change in mass is

$$\Delta m = \frac{\Delta E}{c^2} = \frac{(8.06 \times 10^{26} \text{ MeV})(1.602 \times 10^{-13} \text{ J/MeV})}{(2.998 \times 10^8 \text{ MeV})^2} = 1.44 \times 10^{-3} \text{ kg}$$

About one gram of matter vanishes for each kilogram that fissions!

43. (a) The change in mass is

$$\begin{aligned}\Delta m &= m(^2\text{H}) + m(\text{n}) - m(^3\text{H}) \\ &= 2.014102 \text{ u} + 1.008665 \text{ u} - 3.016049 \text{ u} = 0.006718 \text{ u}\end{aligned}$$

The corresponding energy change is $\Delta E = \Delta m c^2 = (931.5 \text{ MeV/u})(0.006718 \text{ u}) = 6.26 \text{ MeV}$.

(b)

$$\begin{aligned}\Delta m &= m(^3\text{He}) + m(\text{n}) - m(^4\text{He}) \\ &= 3.016029 \text{ u} + 1.008665 \text{ u} - 4.002603 \text{ u} = 0.022352 \text{ u} \\ \Delta E &= \Delta m c^2 = (931.5 \text{ MeV/u})(0.022352 \text{ u}) = 20.8 \text{ MeV}\end{aligned}$$

44. (a)

$$E = E_\pi + E_p = \frac{m_\pi c^2}{\sqrt{1-v^2/c^2}} + m_p c^2 = \frac{139.6 \text{ MeV}}{\sqrt{1-(0.921)^2}} + 938.3 \text{ MeV} = 1296.7 \text{ MeV}$$

(b)

$$p = p_\pi + p_p = \frac{m_\pi v}{\sqrt{1-v^2/c^2}} = \frac{1}{c} \frac{m_\pi c^2 (v/c)}{\sqrt{1-v^2/c^2}} = \frac{1}{c} \frac{(139.6 \text{ MeV})(0.921)}{\sqrt{1-(0.921)^2}} = 330.0 \text{ MeV}/c$$

(c)

$$mc^2 = \sqrt{E^2 - (pc)^2} = \sqrt{(1296.7 \text{ MeV})^2 - (330.0 \text{ MeV})^2} = 1254 \text{ MeV}$$

45. Before the collision, the total relativistic energy of each electron is

$$E_e = \frac{m_e c^2}{\sqrt{1-v^2/c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1-(0.99999)^2}} = 114.3 \text{ MeV}$$

The total energy in the collision is therefore $2 \times 114.3 \text{ MeV} = 228.6 \text{ MeV}$. The total momentum is zero before the collision, because the two particles move with equal and opposite velocities and have equal masses. After the collision, the total momentum is still zero, so we know that the two muons must move with equal speeds and thus have equal energies. The total energy of each muon is then 114.3 MeV and its kinetic energy is

$$K_\mu = E_\mu - m_\mu c^2 = 114.3 \text{ MeV} - 105.7 \text{ MeV} = 8.6 \text{ MeV}$$

46. The two protons have equal (and opposite) momenta and thus equal energies E_1 and E_2 . The new particle is created with zero momentum (at rest), so its total energy is equal to its rest energy $Mc^2 = 9460 \text{ MeV}$. Conservation of energy then gives $E_1 + E_2 = Mc^2$, so $E_1 = E_2 = Mc^2/2$.

$$E_1 = \frac{m_p c^2}{\sqrt{1-v^2/c^2}} = \frac{Mc^2}{2}$$

$$1 - \frac{v^2}{c^2} = \left[\frac{2(m_p c^2)}{Mc^2} \right]^2 = \left[\frac{2(938.3 \text{ MeV})}{9460 \text{ MeV}} \right]^2 = 0.0394 \quad \text{so} \quad v = 0.980c$$

47. For particle 1, moving in the positive x direction,

$$E_1 = K_1 + mc^2 = 282 \text{ MeV} + 140 \text{ MeV} = 422 \text{ MeV}$$

$$cp_1 = \sqrt{E_1^2 - (mc^2)^2} = \sqrt{(422 \text{ MeV})^2 - (140 \text{ MeV})^2} = +398 \text{ MeV}$$

For particle 2, moving in the negative x direction,

$$E_2 = K_2 + mc^2 = 25 \text{ MeV} + 140 \text{ MeV} = 165 \text{ MeV}$$

$$cp_2 = -\sqrt{E_2^2 - (mc^2)^2} = -\sqrt{(165 \text{ MeV})^2 - (140 \text{ MeV})^2} = -87 \text{ MeV}$$

The net final momentum is $p_f = p_1 + p_2 = 398 \text{ MeV}/c - 87 \text{ MeV}/c = 311 \text{ MeV}/c$, and the net final energy is $E_f = E_1 + E_2 = 422 \text{ MeV} + 165 \text{ MeV} = 587 \text{ MeV}$. Because of the conservation laws, these must be equal to the momentum and the energy of the initial particle, so that its rest energy is then

$$m_i c^2 = \sqrt{E_i^2 - (cp_i)^2} = \sqrt{(587 \text{ MeV})^2 - (311 \text{ MeV})^2} = 498 \text{ MeV}$$

Solving Equation 2.36 for v , we obtain

$$v = c \sqrt{1 - \left(\frac{mc^2}{E} \right)^2} = c \sqrt{1 - \left(\frac{498 \text{ MeV}}{587 \text{ MeV}} \right)^2} = 0.529c$$

48. (a) In this frame of reference, the total momentum is zero before and after the collision, and at threshold the four product particles are formed at rest. So the total final energy is $4mc^2$, and since this must be equal to the total initial energy each initial colliding proton must have an energy of $2mc^2$.

(b) With $E = mc^2 / \sqrt{1-v^2/c^2}$, we have for each of the initial protons

$$2mc^2 = \frac{mc^2}{\sqrt{1-v^2/c^2}} \quad \text{or} \quad \sqrt{1-v^2/c^2} = \frac{1}{2}$$

so $v = 0.866c$.

(c) With the transformation speed $u = 0.866c$, we have

$$v' = \frac{v - u}{1 - vu/c^2} = \frac{-0.866c - 0.866c}{1 - (-0.866)(0.866)} = 0.9897c$$

The proton energy in this frame is then

$$E' = \frac{mc^2}{\sqrt{1 - v'^2/c^2}} = \frac{mc^2}{\sqrt{1 - (0.9897)^2}} = 7mc^2$$

49.
$$pc = \frac{(mc^2)(v/c)}{\sqrt{1 - v^2/c^2}}$$

$$\frac{v}{c} = \frac{\sqrt{(pc/mc^2)^2}}{\sqrt{1 + (pc/mc^2)^2}} = \frac{\sqrt{(3094 \text{ MeV}/105.7 \text{ MeV})^2}}{\sqrt{1 + (3094 \text{ MeV}/105.7 \text{ MeV})^2}} = 0.99942$$

$$\Delta t = \frac{\Delta t_0}{\sqrt{1 - v^2/c^2}} = \frac{2.198 \mu s}{\sqrt{1 - (0.99942)^2}} = 64.38 \mu s$$

50.
$$(pc)^2 = E^2 - (mc^2)^2 = (K + mc^2)^2 - (mc^2)^2 = K^2 + 2Kmc^2$$

$$p^2 = 2Km + \frac{K^2}{c^2} \quad \text{so} \quad \frac{p^2}{2K} = m + \frac{K}{2c^2}$$

51. (a) If the astronaut travels at speed v for a distance $d = 200$ ly in the reference frame of Earth, then in the spacecraft reference frame the distance to the star is $d\sqrt{1 - v^2/c^2}$ and the time T for the astronaut to reach the star (which must be 10 years in the spacecraft frame of reference) is

$$\frac{d\sqrt{1 - v^2/c^2}}{v} = T \quad \text{or} \quad \sqrt{1 - v^2/c^2} = v \frac{T}{d} = \frac{v}{c} \frac{cT}{d}$$

Solving, we find

$$v = c \sqrt{\frac{1}{1 + (Tc/d)^2}} = c \sqrt{\frac{1}{1 + (10/200)^2}} = 0.99875c$$

- (b) According to an observer on Earth, the astronaut travels a total distance of 400 ly at a speed of $0.99875c$, so the total time for the round trip is $(400 \text{ ly})/0.99875c = 400.5 \text{ y}$.

52.
$$t'_1 = \frac{t_1 - (u/c^2)x_1}{\sqrt{1 - u^2/c^2}} \quad \text{and} \quad t'_2 = \frac{t_2 - (u/c^2)x_2}{\sqrt{1 - u^2/c^2}}$$

$$t'_2 - t'_1 = \frac{t_2 - t_1}{\sqrt{1 - u^2/c^2}} - \frac{(u/c^2)(x_2 - x_1)}{\sqrt{1 - u^2/c^2}}$$

In the O frame of reference, the “cause” can travel to the “effect” with a speed that can be no greater than c ; that is, $(x_2 - x_1)/(t_2 - t_1) \leq c$, or $(x_2 - x_1) \leq c(t_2 - t_1)$.

Substituting for $(x_2 - x_1)$ in the second term, we obtain

$$t'_2 - t'_1 \geq \frac{t_2 - t_1}{\sqrt{1 - u^2/c^2}} - \frac{(u/c)(t_2 - t_1)}{\sqrt{1 - u^2/c^2}} = (t_2 - t_1) \frac{1 - u/c}{\sqrt{1 - u^2/c^2}} \geq 0$$

53. For the red flash, $x_1 = 0$ at $t_1 = 0$, so $x'_1 = 0$ and $t'_1 = 0$. For the blue flash, which occurs at $x_2 = 3.65 \text{ km}$ and $t_2 = 8.24 \mu\text{s}$,

$$x'_2 = \frac{x_2 - ut_2}{\sqrt{1 - u^2/c^2}} = \frac{3.65 \text{ km} - (0.534c)(8.24 \mu\text{s})}{\sqrt{1 - (0.534)^2}} = -0.89 \text{ km}$$

$$t'_2 = \frac{t_2 - (u/c^2)x_2}{\sqrt{1 - u^2/c^2}} = \frac{8.24 \mu\text{s} - (0.534/0.300 \text{ km}/\mu\text{s})(3.65 \text{ km})}{\sqrt{1 - (0.534)^2}} = 2.06 \mu\text{s}$$

54. Let O' be the observer on ship A , traveling at $u = 0.65c$ relative to the space station. Observer O on the space station measures $v_{Bx} = 0$, $v_{By} = 0.50c$ for ship B . According to O' ,

$$v'_{Bx} = \frac{v_{Bx} - u}{1 - v_{Bx}u/c^2} = \frac{0 - 0.65c}{1 - 0} = -0.65c$$

$$v'_{By} = \frac{v_{By}\sqrt{1 - u^2/c^2}}{1 - v_{Bx}u/c^2} = \frac{0.50c\sqrt{1 - (0.65)^2}}{1 - 0} = 0.38c$$

$$v'_B = \sqrt{(v'_{Bx})^2 + (v'_{By})^2} = \sqrt{(-0.65c)^2 + (0.38c)^2} = 0.75c$$

$$\theta_B = \tan^{-1} \frac{v'_{By}}{v'_{Bx}} = \tan^{-1} \frac{0.38c}{-0.65c} = 150^\circ$$

For ship C , O measures $v_{Cx} = -0.50c$, $v_{Cy} = 0$.

$$v'_{Cx} = \frac{-0.50c - 0.65c}{1 - (-0.50)(0.65)} = -0.87c, \quad v'_{Cy} = 0$$

$$v'_C = 0.87c \text{ at } \theta_C = 180^\circ \text{ (negative } x \text{ direction)}$$

For ship D , O measures $v_{Dx} = (-0.50c) \sin 45^\circ = -0.35c$, $v_{Dy} = +0.35c$.

$$v'_{Dx} = \frac{-0.35c - 0.65c}{1 - (-0.35)(0.65)} = -0.81c, \quad v'_{Dy} = \frac{0.35c \sqrt{1 - (0.65)^2}}{1 - (-0.35)(0.65)} = 0.22c$$

$$v'_D = \sqrt{(-0.81c)^2 + (0.22c)^2} = 0.84c \quad \text{at} \quad \theta_D = \tan^{-1} \frac{0.22c}{-0.81c} = 165^\circ$$

$$55. \quad (a) \quad t' = \frac{t - (u/c^2)x}{\sqrt{1 - u^2/c^2}} = \frac{1.52 \mu s - (0.563/300 \text{ m}/\mu s)(524 \text{ m})}{\sqrt{1 - (0.563)^2}} = 0.648 \mu s$$

$$(b) \quad x' = \frac{x - ut}{\sqrt{1 - u^2/c^2}} = \frac{524 \text{ m} - (0.563 \times 300 \text{ m}/\mu s)(1.52 \mu s)}{\sqrt{1 - (0.563)^2}} = 335 \text{ m}$$

56. (a) O' measures $v' = (v - u)/(1 - uv/c^2)$, and according to O' the energy is

$$E' = \frac{mc^2}{\sqrt{1 - v'^2/c^2}} = \frac{mc^2}{\sqrt{1 - [(v - u)/(1 - uv/c^2)]^2/c^2}}$$

and the momentum is

$$p' = \frac{mv'}{\sqrt{1 - v'^2/c^2}} = \frac{m(v - u)/(1 - uv/c^2)}{\sqrt{1 - [(v - u)/(1 - uv/c^2)]^2/c^2}}$$

$$(b) \quad E'^2 - (p'c)^2 = \frac{m^2c^4 - m^2[(v - u)/(1 - uv/c^2)]^2}{1 - [(v - u)/(1 - uv/c^2)]^2/c^2} = m^2c^4$$

The quantity $E'^2 - (p'c)^2$ is equal to m^2c^4 in *every* frame of reference, no matter what its relative speed u . In other words, every observer measures the same value for the rest energy or mass.

$$57. \quad (a) \quad O' \text{ measures } v'_x = -u, v'_y = v\sqrt{1 - u^2/c^2}.$$


$$v' = \sqrt{(v'_x)^2 + (v'_y)^2} = \sqrt{u^2 + v^2 - u^2 v^2 / c^2}$$

$$E' = \frac{mc^2}{\sqrt{1 - v'^2 / c^2}} = \frac{mc^2}{\sqrt{1 - u^2 / c^2 - v^2 / c^2 + u^2 v^2 / c^4}}$$

$$p' = \frac{mv'}{\sqrt{1 - v'^2 / c^2}} = \frac{m\sqrt{u^2 + v^2 - u^2 v^2 / c^2}}{\sqrt{1 - u^2 / c^2 - v^2 / c^2 + u^2 v^2 / c^4}}$$

$$(b) \quad E'^2 - (p'c)^2 = \frac{m^2 c^4 - m^2 c^2 (u^2 + v^2 - u^2 v^2 / c^2)}{1 - u^2 / c^2 - v^2 / c^2 + u^2 v^2 / c^4} = m^2 c^4$$

58.



(b) 40 years

(c) During Amelia's outbound journey, her worldline and Bernice's overlap. This part of her journey takes 8 years, as measured in the clocks in Bernice's frame of reference.

(d) Let $O = \text{Casper}$ and $O' = \text{Bernice}$. Then $u = 0.60c$ and $v_x = -0.60c$.

$$v'_x = \frac{v_x - u}{1 - v_x u / c^2} = \frac{-0.60c - 0.60c}{1 - (-0.60)(0.60)} = -0.882c$$

(e) On his 4th birthday.

(f) On his 16th birthday.

59. (a) Before the first acceleration, $E = E_0 = mc^2$. After the acceleration, the energy is

$$E_1 = \frac{mc^2}{\sqrt{1 - v^2 / c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1 - (0.99)^2}} = 3.6 \text{ MeV}$$

The change in energy is $\Delta E = E_1 - E_0 = 3.6 \text{ MeV} - 0.5 \text{ MeV} = 3.1 \text{ MeV}$, so the first stage adds 3.1 MeV to the energy of the electron.

$$(b) \quad E_2 = \frac{mc^2}{\sqrt{1 - v^2 / c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1 - (0.999)^2}} = 11.4 \text{ MeV}$$

The change in energy is $\Delta E = E_2 - E_1 = 11.4 \text{ MeV} - 3.6 \text{ MeV} = 7.8 \text{ MeV}$, so the second stage adds about 2.5 times as much energy as the first stage, even though the second stage increases the velocity by only 0.9%.

$$60. \quad K = \frac{mc^2}{\sqrt{1 - v^2 / c^2}} - mc^2 = \frac{0.511 \text{ MeV}}{\sqrt{1 - (0.813)^2}} - 0.511 \text{ MeV} = 0.367 \text{ MeV per particle}$$

$$E_{\text{beam}} = (0.367 \text{ MeV/particle})(2.14 \times 10^{11} \text{ particles/s})(3600 \text{ s}) = 2.82 \times 10^{14} \text{ MeV} = 45.2 \text{ J}$$

Copper has a density of $\rho = 8.92 \text{ g/cm}^3$ and a specific heat capacity of $c_p = 0.385 \text{ J/g}\cdot\text{K}$. The mass M of the copper is $M = \rho V = (8.92 \text{ g/cm}^3)(2.54 \text{ cm})^3 = 146 \text{ g}$. Assuming all of the kinetic energy carried by the particles in the beam acts to produce a change in temperature of the copper ($E_{\text{beam}} = Mc_p \Delta T$), the temperature increase is

$$\Delta T = \frac{E_{\text{beam}}}{Mc_p} = \frac{45.2 \text{ J}}{(146 \text{ g})(0.385 \text{ J/g}\cdot\text{K})} = 0.80 \text{ K} = 0.80 \text{ C}^\circ$$

$$61. \quad (a) \quad p_i = \frac{mv_i}{\sqrt{1 - v_i^2 / c^2}} = \frac{(0.511 \text{ MeV}/c^2)(0.960c)}{\sqrt{1 - (0.960)^2}} = 1.752 \text{ MeV}/c$$

$$p_{1f} = \frac{mv_{1f}}{\sqrt{1-v_{1f}^2/c^2}} = \frac{(0.511 \text{ MeV}/c^2)(0.956c)}{\sqrt{1-(0.956)^2}} = 1.665 \text{ MeV}/c$$

$$p_{xf} = p_{1f} \cos \theta_1 + p_{2f} \cos \theta_2 = (1.665 \text{ MeV}/c)(\cos 9.7^\circ) + p_{2f} \cos \theta_2 = 1.642 \text{ MeV}/c + p_{2f} \cos \theta_2$$

$$p_{yf} = p_{1f} \sin \theta_1 + p_{2f} \sin \theta_2 = (1.665 \text{ MeV}/c)(\sin 9.7^\circ) + p_{2f} \sin \theta_2 = 0.281 \text{ MeV}/c + p_{2f} \sin \theta_2$$

Conservation of momentum gives $p_{xf} = p_i$ and $p_{yf} = 0$. Thus

$$p_{2f} \cos \theta_2 = 1.752 \text{ MeV}/c - 1.642 \text{ MeV}/c = 0.110 \text{ MeV}/c$$

$$p_{2f} \sin \theta_2 = 0 - 0.281 \text{ MeV}/c = -0.281 \text{ MeV}/c$$

Dividing the second result by the first gives

$$\tan \theta_2 = \frac{-0.281 \text{ MeV}/c}{0.110 \text{ MeV}/c} = -2.55 \quad \text{or} \quad \theta_2 = -68.6^\circ$$

With $p_{2f} = (0.110 \text{ MeV}/c)/[\cos(-68.6^\circ)] = 0.302 \text{ MeV}/c$, we have

$$p_{2f} = 0.302 \text{ MeV}/c = \frac{mv_{2f}}{\sqrt{1-v_{2f}^2/c^2}}$$

and solving, we find $v_{2f} = 0.508c$.

$$(b) \quad E_i = \frac{mc^2}{\sqrt{1-v_i^2/c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1-(0.960)^2}} = 2.336 \text{ MeV}$$

$$E_{1f} = \frac{mc^2}{\sqrt{1-v_{1f}^2/c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1-(0.956)^2}} = 1.743 \text{ MeV}$$

Conservation of energy gives $E_i = E_{1f} + E_{2f}$, so $E_{2f} = 2.336 \text{ MeV} - 1.743 \text{ MeV} = 0.593 \text{ MeV}$ and

$$E_{2f} = 0.593 \text{ MeV} = \frac{mc^2}{\sqrt{1-v_{2f}^2/c^2}} = \frac{0.511 \text{ MeV}}{\sqrt{1-v_{2f}^2/c^2}}$$

and solving we find $v_{2f} = 0.508c$, in agreement with part (a).

62. The initial energy is

$$E_\pi = \frac{mc^2}{\sqrt{1-v^2/c^2}} = \frac{135 \text{ MeV}}{\sqrt{1-(0.98)^2}} = 678 \text{ MeV}$$