


Chapter 1

Functions and Graphs


Section 1.1

Check Point Exercises

1.



2.



$$x = -3, y = 7$$

$$x = -2, y = 6$$

$$x = -1, y = 5$$


$$x = 0, y = 4$$

$$x = 1, y = 3$$

$$x = 2, y = 2$$

$$x = 3, y = 1$$

3.



$$x = -4, y = 3$$

$$x = -3, y = 2$$

$$x = -2, y = 1$$

$$x = -1, y = 0$$

$$x = 0, y = 1$$


$$x = 1, y = 2$$

$$x = 2, y = 3$$

4. The meaning of a $[-100, 100, 50]$ by $[-100, 100, 10]$ viewing rectangle is as follows:

distance
between
x-axis
tick
marks
 minimum x-value maximum x-value
 $[-100, 100, 50]$
 by

distance
between
y-axis
tick
marks
 minimum y-value maximum y-value
 $[-100, 100, 10]$













5. a. The graph crosses the x -axis at $(-3, 0)$. Thus, the x -intercept is -3 . The graph crosses the y -axis at $(0, 5)$. Thus, the y -intercept is 5 .
- b. The graph does not cross the x -axis. Thus, there is no x -intercept. The graph crosses the y -axis at $(0, 4)$. Thus, the y -intercept is 4 .
- c. The graph crosses the x - and y -axes at the origin $(0, 0)$. Thus, the x -intercept is 0 and the y -intercept is 0 .
6. a. $d = 4n + 5$
 $d = 4(15) + 5 = 65$
 65% of marriages end in divorce after 15 years when the wife is under 18 at the time of marriage.
- b. According to the line graph, 60% of marriages end in divorce after 15 years when the wife is under 18 at the time of marriage.
- c. The mathematical model overestimates the actual percentage shown in the graph by 5%.


Concept and Vocabulary Check 1.1

1. x -axis
2. y -axis
3. origin
4. quadrants; four
5. x -coordinate; y -coordinate
6. solution; satisfies
7. x -intercept; zero
8. y -intercept; zero


Exercise Set 1.1




11.



12.




13.




$x = -3, y = 7$
 $x = -2, y = 2$
 $x = -1, y = -1$
 $x = 0, y = -2$
 $x = 1, y = -1$
 $x = 2, y = 2$
 $x = 3, y = 7$

14.




$x = -3, y = 11$
 $x = -2, y = 6$
 $x = -1, y = 3$
 $x = 0, y = 2$
 $x = 1, y = 3$
 $x = 2, y = 6$
 $x = 3, y = 11$

15.




$x = -3, y = -5$
 $x = -2, y = -4$
 $x = -1, y = -3$
 $x = 0, y = -2$
 $x = 1, y = -1$
 $x = 2, y = 0$
 $x = 3, y = 1$

16.




$x = -3, y = -1$
 $x = -2, y = 0$
 $x = -1, y = 1$
 $x = 0, y = 2$
 $x = 1, y = 3$
 $x = 2, y = 4$
 $x = 3, y = 5$

17.




$x = -3, y = -5$
 $x = -2, y = -3$
 $x = -1, y = -1$
 $x = 0, y = 1$
 $x = 1, y = 3$
 $x = 2, y = 5$
 $x = 3, y = 7$

18.




$x = -3, y = -10$
 $x = -2, y = -8$
 $x = -1, y = -6$
 $x = 0, y = -4$
 $x = 1, y = -2$
 $x = 2, y = 0$
 $x = 3, y = 2$

19.




$x = -3, y = \frac{3}{2}$
 $x = -2, y = 1$
 $x = -1, y = \frac{1}{2}$
 $x = 0, y = 0$
 $x = 1, y = -\frac{1}{2}$
 $x = 2, y = -1$
 $x = 3, y = -\frac{3}{2}$

20.




$x = -3, y = \frac{7}{2}$
 $x = -2, y = 3$
 $x = -1, y = \frac{5}{2}$
 $x = 0, y = 2$
 $x = 1, y = \frac{3}{2}$
 $x = 2, y = 1$
 $x = 3, y = \frac{1}{2}$

21.




$x = -3, y = 6$
 $x = -2, y = 4$
 $x = -1, y = 2$
 $x = 0, y = 0$
 $x = 1, y = 2$
 $x = 2, y = 4$
 $x = 3, y = 6$


22.




$x = -3, y = -6$
 $x = -2, y = -4$
 $x = -1, y = -2$
 $x = 0, y = 0$
 $x = 1, y = -2$
 $x = 2, y = -4$
 $x = 3, y = -6$




- $x = -3, y = 4$
- $x = -2, y = 3$
- $x = -1, y = 2$
- $x = 0, y = 1$
- $x = 1, y = 2$
- $x = 2, y = 3$
- $x = 3, y = 4$




- $x = -3, y = 2$
- $x = -2, y = 1$
- $x = -1, y = 0$
- $x = 0, y = -1$
- $x = 1, y = 0$
- $x = 2, y = 1$
- $x = 3, y = 2$




- $x = -3, y = 0$
- $x = -2, y = 5$
- $x = -1, y = 8$
- $x = 0, y = 9$
- $x = 1, y = 8$
- $x = 2, y = 5$
- $x = 3, y = 0$



- $x = -3, y = -9$
- $x = -2, y = -4$
- $x = -1, y = -1$
- $x = 0, y = 0$
- $x = 1, y = -1$
- $x = 2, y = -4$
- $x = 3, y = -9$



- $x = -3, y = -27$
- $x = -2, y = -8$
- $x = -1, y = -1$
- $x = 0, y = 0$
- $x = 1, y = 1$
- $x = 2, y = 8$
- $x = 3, y = 27$



- $x = -3, y = -28$
- $x = -2, y = -9$
- $x = -1, y = -2$
- $x = 0, y = -1$
- $x = 1, y = 0$
- $x = 2, y = 7$
- $x = 3, y = 26$

29. (c) x-axis tick marks $-5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5$; y-axis tick marks are the same.

30. (d) x -axis tick marks $-10, -8, -6, -4, -2, 0, 2, 4, 6, 8, 10$; y -axis tick marks $-4, -2, 0, 2, 4$

31. (b); x -axis tick marks $-20, -10, 0, 10, 20, 30, 40, 50, 60, 70, 80$; y -axis tick marks $-30, -20, -10, 0, 10, 20, 30, 40, 50, 60, 70$

32. (a) x -axis tick marks $-40, -20, 0, 20, 40$; y -axis tick marks $-1000, -900, -800, -700, \dots, 700, 800, 900, 1000$

33. The equation that corresponds to Y_2 in the table is (c), $y_2 = 2 - x$. We can tell because all of the points $(-3, 5)$, $(-2, 4)$, $(-1, 3)$, $(0, 2)$, $(1, 1)$, $(2, 0)$, and $(3, -1)$ are on the line $y = 2 - x$, but all are not on any of the others.

34. The equation that corresponds to Y_1 in the table is (b), $y_1 = x^2$. We can tell because all of the points $(-3, 9)$, $(-2, 4)$, $(-1, 1)$, $(0, 0)$, $(1, 1)$, $(2, 4)$, and $(3, 9)$ are on the graph $y = x^2$, but all are not on any of the others.

35. No. It passes through the point $(0, 2)$.

36. Yes. It passes through the point $(0, 0)$.

37. $(2, 0)$

38. $(0, 2)$

39. The graphs of Y_1 and Y_2 intersect at the points $(-2, 4)$ and $(1, 1)$.

40. The values of Y_1 and Y_2 are the same when $x = -2$ and $x = 1$.

41. a. 2; The graph intersects the x -axis at $(2, 0)$.

b. -4 ; The graph intersects the y -axis at $(0, -4)$.

42. a. 1; The graph intersects the x -axis at $(1, 0)$.

b. 2; The graph intersects the y -axis at $(0, 2)$.

43. a. 1, -2 ; The graph intersects the x -axis at $(1, 0)$ and $(-2, 0)$.

b. 2; The graph intersects the y -axis at $(0, 2)$.

44. a. 1, -1 ; The graph intersects the x -axis at $(1, 0)$ and $(-1, 0)$.

b. 1; The graph intersects the y -axis at $(0, 1)$.


45. a. -1 ; The graph intersects the x -axis at $(-1, 0)$.

b. none; The graph does not intersect the y -axis.


46. a. none; The graph does not intersect the x -axis.

b. 2; The graph intersects the y -axis at $(0, 2)$.


47.




48.



49.




50.




51.

x	(x, y)
-3	$(-3, 5)$
-2	$(-2, 5)$
-1	$(-1, 5)$
0	$(0, 5)$
1	$(1, 5)$
2	$(2, 5)$
3	$(3, 5)$




52.

x	(x, y)
-3	$(-3, -1)$
-2	$(-2, -1)$
-1	$(-1, -1)$
0	$(0, -1)$
1	$(1, -1)$
2	$(2, -1)$
3	$(3, -1)$




53.

x	(x, y)
-2	$\left(-2, -\frac{1}{2}\right)$
-1	$(-1, -1)$
$-\frac{1}{2}$	$\left(-\frac{1}{2}, -2\right)$
$-\frac{1}{3}$	$\left(-\frac{1}{3}, -3\right)$
$\frac{1}{3}$	$\left(\frac{1}{3}, 3\right)$
$\frac{1}{2}$	$\left(\frac{1}{2}, 2\right)$
1	$(1, 1)$
2	$\left(2, \frac{1}{2}\right)$



54.

x	(x, y)
-2	$\left(-2, \frac{1}{2}\right)$
-1	$(-1, 1)$
$-\frac{1}{2}$	$\left(-\frac{1}{2}, 2\right)$
$-\frac{1}{3}$	$\left(-\frac{1}{3}, 3\right)$
$\frac{1}{3}$	$\left(\frac{1}{3}, -3\right)$
$\frac{1}{2}$	$\left(\frac{1}{2}, -2\right)$
1	$(1, -1)$
2	$\left(2, -\frac{1}{2}\right)$



55. a. According to the line graph, 20% of seniors used marijuana in 2005.
- b. 2005 is 25 years after 1980.
 $M = -0.3n + 27$
 $M = -0.3(25) + 27 = 19.5$
 According to formula, 19.5% of seniors used marijuana in 2005. This underestimates the value in the graph by 0.5%.
- c. According to the line graph, about 47% of seniors used alcohol in 2005.
- d. 2005 is 25 years after 1980.
 $A = -0.9n + 69$
 $A = -0.9(25) + 69 = 46.5$
 According to formula, 46.5% of seniors used alcohol in 2005. It is less than the estimate, although answers may vary.
- e. The minimum for marijuana was reached in 1990.
 According to the line graph, about 14% of seniors used marijuana in 1990.
56. a. According to the line graph, 50% of seniors used alcohol in 2000.
- b. 2000 is 20 years after 1980.
 $A = -0.9n + 69$
 $A = -0.9(20) + 69 = 51$
 According to formula, 51% of seniors used alcohol in 2000. This overestimates the value in the graph by 1%.
- c. According to the line graph, about 22% of seniors used marijuana in 2000.
- d. 2000 is 20 years after 1980.
 $M = -0.3n + 27$
 $M = -0.3(20) + 27 = 21$
 According to formula, 21% of seniors used marijuana in 2000. It is less than the estimate, although answers may vary.

- e. The maximum for alcohol was reached in 1980. According to the line graph, about 72% of seniors used alcohol in 1980.
57. At age 8, women have the least number of awakenings, averaging about 1 awakening per night.
58. At age 65, men have the greatest number of awakenings, averaging about 8 awakenings per night.
59. The difference between the number of awakenings for 25-year-old men and women is about 1.9.
60. The difference between the number of awakenings for 18-year-old men and women is about 1.1.
61. – 66. Answers will vary.
67. makes sense
68. does not make sense; Explanations will vary. Sample explanation: Most graphing utilities do not display numbers on the axes.
69. does not make sense; Explanations will vary. Sample explanation: These three points are not collinear.
70. does not make sense; Explanations will vary. Sample explanation: As the time of day goes up, the total calories burned will also go up.
71. false; Changes to make the statement true will vary. A sample change is: The product of the coordinates of a point in quadrant III is also positive.
72. false; Changes to make the statement true will vary. A sample change is: A point on the x -axis will have $y = 0$.
73. true
74. false; Changes to make the statement true will vary. A sample change is: $3(5) - 2(2) \neq -4$.
75. I, III
76. II, IV
77. IV
78. II
79. (a)
80. (d)
81. (b)

82. (c)

83. (b)

84. (a)

85. (c)


86. (b)

87. Set 1 has each x -coordinate paired with only one y -coordinate.

88.

x	$y = 2x$	(x, y)
-2	$y = 2(-2) = -4$	$(-2, -4)$
-1	$y = 2(-1) + 4 = 2$	$(-1, 2)$
0	$y = 2(0) = 0$	$(0, 0)$
1	$y = 2(1) = 2$	$(1, 2)$
2	$y = 2(2) = 4$	$(2, 4)$

x	$y = 2x + 4$	(x, y)
-2	$y = 2(-2) + 4 = 0$	$(-2, 0)$
-1	$y = 2(-1) + 4 = 2$	$(-1, 2)$
0	$y = 2(0) + 4 = 4$	$(0, 4)$
1	$y = 2(1) + 4 = 6$	$(1, 6)$
2	$y = 2(2) + 4 = 8$	$(2, 8)$



89. a. When the x -coordinate is 2, the y -coordinate is 3.

b. When the y -coordinate is 4, the x -coordinates are -3 and 3.

c. The x -coordinates are all real numbers.

d. The y -coordinates are all real numbers greater than or equal to 1.

Section 1.2

Check Point Exercises

1. The domain is the set of all first components: $\{0, 10, 20, 30, 40\}$. The range is the set of all second components: $\{9.1, 6.7, 10.7, 13.2, 21.2\}$.

2. a. The relation is not a function since the two ordered pairs $(5, 6)$ and $(5, 8)$ have the same first component but different second components.

b. The relation is a function since no two ordered pairs have the same first component and different second components.

3. a. $2x + y = 6$
 $y = 6 - 2x$

For each value of x , there is one and only one value for y , so the equation defines y as a function of x .

b. $x^2 + y^2 = 1$
 $y^2 = 1 - x^2$
 $y = \pm\sqrt{1 - x^2}$

Since there are values of x (all values between -1 and 1 exclusive) that give more than one value for y (for example, if $x = 0$, then

$y = \pm\sqrt{1 - 0^2} = \pm 1$), the equation does not define y as a function of x .

4. a. $f(-5) = (-5)^2 - 2(-5) + 7$
 $= 25 - (-10) + 7$
 $= 42$


b. $f(x+4) = (x+4)^2 - 2(x+4) + 7$
 $= x^2 + 8x + 16 - 2x - 8 + 7$
 $= x^2 + 6x + 15$

c. $f(-x) = (-x)^2 - 2(-x) + 7$
 $= x^2 - (-2x) + 7$
 $= x^2 + 2x + 7$

5.

x	$f(x) = 2x$	(x, y)
-2	-4	$(-2, -4)$
-1	-2	$(-1, -2)$
0	0	$(0, 0)$
1	2	$(1, 2)$
2	4	$(2, 4)$

x	$g(x) = 2x - 3$	(x, y)
-2	$g(-2) = 2(-2) - 3 = -7$	$(-2, -7)$
-1	$g(-1) = 2(-1) - 3 = -5$	$(-1, -5)$
0	$g(0) = 2(0) - 3 = -3$	$(0, -3)$
1	$g(1) = 2(1) - 3 = -1$	$(1, -1)$
2	$g(2) = 2(2) - 3 = 1$	$(2, 1)$



The graph of g is the graph of f shifted down 3 units.

6. The graph (a) passes the vertical line test and is therefore is a function.

The graph (b) fails the vertical line test and is therefore not a function.

The graph (c) passes the vertical line test and is therefore is a function.

The graph (d) fails the vertical line test and is therefore not a function.

7. a. $f(5) = 400$

b. $x = 9$, $f(9) = 100$

c. The minimum T cell count in the asymptomatic stage is approximately 425.

8. a. domain: $\{x | -2 \leq x \leq 1\}$ or $[-2, 1]$.
range: $\{y | 0 \leq y \leq 3\}$ or $[0, 3]$.

b. domain: $\{x | -2 < x \leq 1\}$ or $(-2, 1]$.
range: $\{y | -1 \leq y < 2\}$ or $[-1, 2)$.

c. domain: $\{x | -3 \leq x < 0\}$ or $[-3, 0)$.
range: $\{y | y = -3, -2, -1\}$.

Concept and Vocabulary Check 1.2

1. relation; domain; range

2. function

3. f ; x

4. true

5. false

6. x ; $x + 6$

7. ordered pairs

8. more than once; function

9. $[0, 3)$; domain

10. $[1, \infty)$; range

11. 0; 0; zeros

12. false

Exercise Set 1.2

1. The relation is a function since no two ordered pairs have the same first component and different second components. The domain is $\{1, 3, 5\}$ and the range is $\{2, 4, 5\}$.

2. The relation is a function because no two ordered pairs have the same first component and different second components. The domain is $\{4, 6, 8\}$ and the range is $\{5, 7, 8\}$.

3. The relation is not a function since the two ordered pairs $(3, 4)$ and $(3, 5)$ have the same first component but different second components (the same could be said for the ordered pairs $(4, 4)$ and $(4, 5)$). The domain is $\{3, 4\}$ and the range is $\{4, 5\}$.

4. The relation is not a function since the two ordered pairs (5, 6) and (5, 7) have the same first component but different second components (the same could be said for the ordered pairs (6, 6) and (6, 7)). The domain is {5, 6} and the range is {6, 7}.

5. The relation is a function because no two ordered pairs have the same first component and different second components. The domain is {3, 4, 5, 7} and the range is {-2, 1, 9}.

6. The relation is a function because no two ordered pairs have the same first component and different second components. The domain is {-2, -1, 5, 10} and the range is {1, 4, 6}.

7. The relation is a function since there are no same first components with different second components. The domain is {-3, -2, -1, 0} and the range is {-3, -2, -1, 0}.

8. The relation is a function since there are no ordered pairs that have the same first component but different second components. The domain is {-7, -5, -3, 0} and the range is {-7, -5, -3, 0}.

9. The relation is not a function since there are ordered pairs with the same first component and different second components. The domain is {1} and the range is {4, 5, 6}.

10. The relation is a function since there are no two ordered pairs that have the same first component and different second components. The domain is {4, 5, 6} and the range is {1}.

11. $x + y = 16$
 $y = 16 - x$
 Since only one value of y can be obtained for each value of x , y is a function of x .

12. $x + y = 25$
 $y = 25 - x$
 Since only one value of y can be obtained for each value of x , y is a function of x .

13. $x^2 + y = 16$
 $y = 16 - x^2$
 Since only one value of y can be obtained for each value of x , y is a function of x .

14. $x^2 + y = 25$
 $y = 25 - x^2$
 Since only one value of y can be obtained for each value of x , y is a function of x .

15. $x^2 + y^2 = 16$
 $y^2 = 16 - x^2$
 $y = \pm\sqrt{16 - x^2}$
 If $x = 0$, $y = \pm 4$.
 Since two values, $y = 4$ and $y = -4$, can be obtained for one value of x , y is not a function of x .

16. $x^2 + y^2 = 25$
 $y^2 = 25 - x^2$
 $y = \pm\sqrt{25 - x^2}$
 If $x = 0$, $y = \pm 5$.
 Since two values, $y = 5$ and $y = -5$, can be obtained for one value of x , y is not a function of x .

17. $x = y^2$
 $y = \pm\sqrt{x}$
 If $x = 1$, $y = \pm 1$.
 Since two values, $y = 1$ and $y = -1$, can be obtained for $x = 1$, y is not a function of x .

18. $4x = y^2$
 $y = \pm\sqrt{4x} = \pm 2\sqrt{x}$
 If $x = 1$, then $y = \pm 2$.
 Since two values, $y = 2$ and $y = -2$, can be obtained for $x = 1$, y is not a function of x .

19. $y = \sqrt{x + 4}$
 Since only one value of y can be obtained for each value of x , y is a function of x .

20. $y = -\sqrt{x + 4}$
 Since only one value of y can be obtained for each value of x , y is a function of x .

21. $x + y^3 = 8$
 $y^3 = 8 - x$
 $y = \sqrt[3]{8 - x}$
 Since only one value of y can be obtained for each value of x , y is a function of x .

22. $x + y^3 = 27$

$$y^3 = 27 - x$$

$$y = \sqrt[3]{27 - x}$$

Since only one value of y can be obtained for each value of x , y is a function of x .

23. $xy + 2y = 1$

$$y(x + 2) = 1$$

$$y = \frac{1}{x + 2}$$

Since only one value of y can be obtained for each value of x , y is a function of x .

24. $xy - 5y = 1$

$$y(x - 5) = 1$$

$$y = \frac{1}{x - 5}$$

Since only one value of y can be obtained for each value of x , y is a function of x .

25. $|x| - y = 2$

$$-y = -|x| + 2$$

$$y = |x| - 2$$

Since only one value of y can be obtained for each value of x , y is a function of x .

26. $|x| - y = 5$

$$-y = -|x| + 5$$

$$y = |x| - 5$$

Since only one value of y can be obtained for each value of x , y is a function of x .

27. a. $f(6) = 4(6) + 5 = 29$

b. $f(x + 1) = 4(x + 1) + 5 = 4x + 9$

c. $f(-x) = 4(-x) + 5 = -4x + 5$

28. a. $f(4) = 3(4) + 7 = 19$

b. $f(x + 1) = 3(x + 1) + 7 = 3x + 10$

c. $f(-x) = 3(-x) + 7 = -3x + 7$

29. a. $g(-1) = (-1)^2 + 2(-1) + 3$
 $= 1 - 2 + 3$
 $= 2$

b. $g(x + 5) = (x + 5)^2 + 2(x + 5) + 3$
 $= x^2 + 10x + 25 + 2x + 10 + 3$
 $= x^2 + 12x + 38$

c. $g(-x) = (-x)^2 + 2(-x) + 3$
 $= x^2 - 2x + 3$

30. a. $g(-1) = (-1)^2 - 10(-1) - 3$
 $= 1 + 10 - 3$
 $= 8$

b. $g(x + 2) = (x + 2)^2 - 10(8 + 2) - 3$
 $= x^2 + 4x + 4 - 10x - 20 - 3$
 $= x^2 - 6x - 19$

c. $g(-x) = (-x)^2 - 10(-x) - 3$
 $= x^2 + 10x - 3$

31. a. $h(2) = 2^4 - 2^2 + 1$
 $= 16 - 4 + 1$
 $= 13$

b. $h(-1) = (-1)^4 - (-1)^2 + 1$
 $= 1 - 1 + 1$
 $= 1$

c. $h(-x) = (-x)^4 - (-x)^2 + 1 = x^4 - x^2 + 1$

d. $h(3a) = (3a)^4 - (3a)^2 + 1$
 $= 81a^4 - 9a^2 + 1$

32. a. $h(3) = 3^3 - 3 + 1 = 25$

b. $h(-2) = (-2)^3 - (-2) + 1$
 $= -8 + 2 + 1$
 $= -5$

c. $h(-x) = (-x)^3 - (-x) + 1 = -x^3 + x + 1$

d. $h(3a) = (3a)^3 - (3a) + 1$
 $= 27a^3 - 3a + 1$

$$33. \text{ a. } f(-6) = \sqrt{-6+6} + 3 = \sqrt{0} + 3 = 3$$

$$\begin{aligned} \text{b. } f(10) &= \sqrt{10+6} + 3 \\ &= \sqrt{16} + 3 \\ &= 4 + 3 \\ &= 7 \end{aligned}$$

$$\text{c. } f(x-6) = \sqrt{x-6+6} + 3 = \sqrt{x} + 3$$

$$34. \text{ a. } f(16) = \sqrt{25-16} - 6 = \sqrt{9} - 6 = 3 - 6 = -3$$

$$\begin{aligned} \text{b. } f(-24) &= \sqrt{25-(-24)} - 6 \\ &= \sqrt{49} - 6 \\ &= 7 - 6 = 1 \end{aligned}$$

$$\begin{aligned} \text{c. } f(25-2x) &= \sqrt{25-(25-2x)} - 6 \\ &= \sqrt{2x} - 6 \end{aligned}$$

$$35. \text{ a. } f(2) = \frac{4(2)^2 - 1}{2^2} = \frac{15}{4}$$

$$\text{b. } f(-2) = \frac{4(-2)^2 - 1}{(-2)^2} = \frac{15}{4}$$

$$\text{c. } f(-x) = \frac{4(-x)^2 - 1}{(-x)^2} = \frac{4x^2 - 1}{x^2}$$

$$36. \text{ a. } f(2) = \frac{4(2)^3 + 1}{2^3} = \frac{33}{8}$$

$$\text{b. } f(-2) = \frac{4(-2)^3 + 1}{(-2)^3} = \frac{-31}{-8} = \frac{31}{8}$$

$$\begin{aligned} \text{c. } f(-x) &= \frac{4(-x)^3 + 1}{(-x)^3} = \frac{-4x^3 + 1}{-x^3} \\ &\text{or } \frac{4x^3 - 1}{x^3} \end{aligned}$$

$$37. \text{ a. } f(6) = \frac{6}{|6|} = 1$$

$$\text{b. } f(-6) = \frac{-6}{|-6|} = \frac{-6}{6} = -1$$

$$\text{c. } f(r^2) = \frac{r^2}{|r^2|} = \frac{r^2}{r^2} = 1$$

$$38. \text{ a. } f(5) = \frac{|5+3|}{5+3} = \frac{|8|}{8} = 1$$


$$\text{b. } f(-5) = \frac{|-5+3|}{-5+3} = \frac{|-2|}{-2} = \frac{2}{-2} = -1$$

$$\begin{aligned} \text{c. } f(-9-x) &= \frac{|-9-x+3|}{-9-x+3} \\ &= \frac{|-x-6|}{-x-6} = \begin{cases} 1, & \text{if } x < -6 \\ -1, & \text{if } x > -6 \end{cases} \end{aligned}$$

39.

x	$f(x) = x$	(x, y)
-2	$f(-2) = -2$	$(-2, -2)$
-1	$f(-1) = -1$	$(-1, -1)$
0	$f(0) = 0$	$(0, 0)$
1	$f(1) = 1$	$(1, 1)$
2	$f(2) = 2$	$(2, 2)$


x	$g(x) = x + 3$	(x, y)
-2	$g(-2) = -2 + 3 = 1$	$(-2, 1)$
-1	$g(-1) = -1 + 3 = 2$	$(-1, 2)$
0	$g(0) = 0 + 3 = 3$	$(0, 3)$
1	$g(1) = 1 + 3 = 4$	$(1, 4)$
2	$g(2) = 2 + 3 = 5$	$(2, 5)$


 The graph of g is the graph of f shifted up 3 units.

40.

x	$f(x) = x$	(x, y)
-2	$f(-2) = -2$	$(-2, -2)$
-1	$f(-1) = -1$	$(-1, -1)$
0	$f(0) = 0$	$(0, 0)$
1	$f(1) = 1$	$(1, 1)$
2	$f(2) = 2$	$(2, 2)$

x	$g(x) = x - 4$	(x, y)
-2	$g(-2) = -2 - 4 = -6$	$(-2, -6)$
-1	$g(-1) = -1 - 4 = -5$	$(-1, -5)$
0	$g(0) = 0 - 4 = -4$	$(0, -4)$
1	$g(1) = 1 - 4 = -3$	$(1, -3)$
2	$g(2) = 2 - 4 = -2$	$(2, -2)$




The graph of g is the graph of f shifted down 4 units.

41.

x	$f(x) = -2x$	(x, y)
-2	$f(-2) = -2(-2) = 4$	$(-2, 4)$
-1	$f(-1) = -2(-1) = 2$	$(-1, 2)$
0	$f(0) = -2(0) = 0$	$(0, 0)$
1	$f(1) = -2(1) = -2$	$(1, -2)$
2	$f(2) = -2(2) = -4$	$(2, -4)$

x	$g(x) = -2x - 1$	(x, y)
-2	$g(-2) = -2(-2) - 1 = 3$	$(-2, 3)$
-1	$g(-1) = -2(-1) - 1 = 1$	$(-1, 1)$
0	$g(0) = -2(0) - 1 = -1$	$(0, -1)$
1	$g(1) = -2(1) - 1 = -3$	$(1, -3)$
2	$g(2) = -2(2) - 1 = -5$	$(2, -5)$




The graph of g is the graph of f shifted down 1 unit.

42.

x	$f(x) = -2x$	(x, y)
-2	$f(-2) = -2(-2) = 4$	$(-2, 4)$
-1	$f(-1) = -2(-1) = 2$	$(-1, 2)$
0	$f(0) = -2(0) = 0$	$(0, 0)$
1	$f(1) = -2(1) = -2$	$(1, -2)$
2	$f(2) = -2(2) = -4$	$(2, -4)$

x	$g(x) = -2x + 3$	(x, y)
-2	$g(-2) = -2(-2) + 3 = 7$	$(-2, 7)$
-1	$g(-1) = -2(-1) + 3 = 5$	$(-1, 5)$
0	$g(0) = -2(0) + 3 = 3$	$(0, 3)$
1	$g(1) = -2(1) + 3 = 1$	$(1, 1)$
2	$g(2) = -2(2) + 3 = -1$	$(2, -1)$




The graph of g is the graph of f shifted up 3 units.

43.

x	$f(x) = x^2$	(x, y)
-2	$f(-2) = (-2)^2 = 4$	$(-2, 4)$
-1	$f(-1) = (-1)^2 = 1$	$(-1, 1)$
0	$f(0) = (0)^2 = 0$	$(0, 0)$
1	$f(1) = (1)^2 = 1$	$(1, 1)$
2	$f(2) = (2)^2 = 4$	$(2, 4)$

x	$g(x) = x^2 + 1$	(x, y)
-2	$g(-2) = (-2)^2 + 1 = 5$	$(-2, 5)$
-1	$g(-1) = (-1)^2 + 1 = 2$	$(-1, 2)$
0	$g(0) = (0)^2 + 1 = 1$	$(0, 1)$
1	$g(1) = (1)^2 + 1 = 2$	$(1, 2)$
2	$g(2) = (2)^2 + 1 = 5$	$(2, 5)$




The graph of g is the graph of f shifted up 1 unit.

44.

x	$f(x) = x^2$	(x, y)
-2	$f(-2) = (-2)^2 = 4$	$(-2, 4)$
-1	$f(-1) = (-1)^2 = 1$	$(-1, 1)$
0	$f(0) = (0)^2 = 0$	$(0, 0)$
1	$f(1) = (1)^2 = 1$	$(1, 1)$
2	$f(2) = (2)^2 = 4$	$(2, 4)$

x	$g(x) = x^2 - 2$	(x, y)
-2	$g(-2) = (-2)^2 - 2 = 2$	$(-2, 2)$
-1	$g(-1) = (-1)^2 - 2 = -1$	$(-1, -1)$
0	$g(0) = (0)^2 - 2 = -2$	$(0, -2)$
1	$g(1) = (1)^2 - 2 = -1$	$(1, -1)$
2	$g(2) = (2)^2 - 2 = 2$	$(2, 2)$




The graph of g is the graph of f shifted down 2 units.

45.

x	$f(x) = x $	(x, y)
-2	$f(-2) = -2 = 2$	$(-2, 2)$
-1	$f(-1) = -1 = 1$	$(-1, 1)$
0	$f(0) = 0 = 0$	$(0, 0)$
1	$f(1) = 1 = 1$	$(1, 1)$
2	$f(2) = 2 = 2$	$(2, 2)$

x	$g(x) = x - 2$	(x, y)
-2	$g(-2) = -2 - 2 = 0$	$(-2, 0)$
-1	$g(-1) = -1 - 2 = -1$	$(-1, -1)$
0	$g(0) = 0 - 2 = -2$	$(0, -2)$
1	$g(1) = 1 - 2 = -1$	$(1, -1)$
2	$g(2) = 2 - 2 = 0$	$(2, 0)$




The graph of g is the graph of f shifted down 2 units.

46.

x	$f(x) = x $	(x, y)
-2	$f(-2) = -2 = 2$	$(-2, 2)$
-1	$f(-1) = -1 = 1$	$(-1, 1)$
0	$f(0) = 0 = 0$	$(0, 0)$
1	$f(1) = 1 = 1$	$(1, 1)$
2	$f(2) = 2 = 2$	$(2, 2)$

x	$g(x) = x + 1$	(x, y)
-2	$g(-2) = -2 + 1 = 3$	$(-2, 3)$
-1	$g(-1) = -1 + 1 = 2$	$(-1, 2)$
0	$g(0) = 0 + 1 = 1$	$(0, 1)$
1	$g(1) = 1 + 1 = 2$	$(1, 2)$
2	$g(2) = 2 + 1 = 3$	$(2, 3)$




The graph of g is the graph of f shifted up 1 unit.

47.

x	$f(x) = x^3$	(x, y)
-2	$f(-2) = (-2)^3 = -8$	$(-2, -8)$
-1	$f(-1) = (-1)^3 = -1$	$(-1, -1)$
0	$f(0) = (0)^3 = 0$	$(0, 0)$
1	$f(1) = (1)^3 = 1$	$(1, 1)$
2	$f(2) = (2)^3 = 8$	$(2, 8)$

x	$g(x) = x^3 + 2$	(x, y)
-2	$g(-2) = (-2)^3 + 2 = -6$	$(-2, -6)$
-1	$g(-1) = (-1)^3 + 2 = 1$	$(-1, 1)$
0	$g(0) = (0)^3 + 2 = 2$	$(0, 2)$
1	$g(1) = (1)^3 + 2 = 3$	$(1, 3)$
2	$g(2) = (2)^3 + 2 = 10$	$(2, 10)$




The graph of g is the graph of f shifted up 2 units.

48.

x	$f(x) = x^3$	(x, y)
-2	$f(-2) = (-2)^3 = -8$	$(-2, -8)$
-1	$f(-1) = (-1)^3 = -1$	$(-1, -1)$
0	$f(0) = (0)^3 = 0$	$(0, 0)$
1	$f(1) = (1)^3 = 1$	$(1, 1)$
2	$f(2) = (2)^3 = 8$	$(2, 8)$

x	$g(x) = x^3 - 1$	(x, y)
-2	$g(-2) = (-2)^3 - 1 = -9$	$(-2, -9)$
-1	$g(-1) = (-1)^3 - 1 = -2$	$(-1, -2)$
0	$g(0) = (0)^3 - 1 = -1$	$(0, -1)$
1	$g(1) = (1)^3 - 1 = 0$	$(1, 0)$
2	$g(2) = (2)^3 - 1 = 7$	$(2, 7)$




The graph of g is the graph of f shifted down 1 unit.

49.

x	$f(x) = 3$	(x, y)
-2	$f(-2) = 3$	$(-2, 3)$
-1	$f(-1) = 3$	$(-1, 3)$
0	$f(0) = 3$	$(0, 3)$
1	$f(1) = 3$	$(1, 3)$
2	$f(2) = 3$	$(2, 3)$

x	$g(x) = 5$	(x, y)
-2	$g(-2) = 5$	$(-2, 5)$
-1	$g(-1) = 5$	$(-1, 5)$
0	$g(0) = 5$	$(0, 5)$
1	$g(1) = 5$	$(1, 5)$
2	$g(2) = 5$	$(2, 5)$




The graph of g is the graph of f shifted up 2 units.

50.

x	$f(x) = -1$	(x, y)
-2	$f(-2) = -1$	$(-2, -1)$
-1	$f(-1) = -1$	$(-1, -1)$
0	$f(0) = -1$	$(0, -1)$
1	$f(1) = -1$	$(1, -1)$
2	$f(2) = -1$	$(2, -1)$

x	$g(x) = 4$	(x, y)
-2	$g(-2) = 4$	$(-2, 4)$
-1	$g(-1) = 4$	$(-1, 4)$
0	$g(0) = 4$	$(0, 4)$
1	$g(1) = 4$	$(1, 4)$
2	$g(2) = 4$	$(2, 4)$




The graph of g is the graph of f shifted up 5 units.

51.

x	$f(x) = \sqrt{x}$	(x, y)
0	$f(0) = \sqrt{0} = 0$	$(0, 0)$
1	$f(1) = \sqrt{1} = 1$	$(1, 1)$
4	$f(4) = \sqrt{4} = 2$	$(4, 2)$
9	$f(9) = \sqrt{9} = 3$	$(9, 3)$

x	$g(x) = \sqrt{x} - 1$	(x, y)
0	$g(0) = \sqrt{0} - 1 = -1$	$(0, -1)$
1	$g(1) = \sqrt{1} - 1 = 0$	$(1, 0)$
4	$g(4) = \sqrt{4} - 1 = 1$	$(4, 1)$
9	$g(9) = \sqrt{9} - 1 = 2$	$(9, 2)$




The graph of g is the graph of f shifted down 1 unit.

52.

x	$f(x) = \sqrt{x}$	(x, y)
0	$f(0) = \sqrt{0} = 0$	$(0, 0)$
1	$f(1) = \sqrt{1} = 1$	$(1, 1)$
4	$f(4) = \sqrt{4} = 2$	$(4, 2)$
9	$f(9) = \sqrt{9} = 3$	$(9, 3)$

x	$g(x) = \sqrt{x} + 2$	(x, y)
0	$g(0) = \sqrt{0} + 2 = 2$	$(0, 2)$
1	$g(1) = \sqrt{1} + 2 = 3$	$(1, 3)$
4	$g(4) = \sqrt{4} + 2 = 4$	$(4, 4)$
9	$g(9) = \sqrt{9} + 2 = 5$	$(9, 5)$




The graph of g is the graph of f shifted up 2 units.

53.

x	$f(x) = \sqrt{x}$	(x, y)
0	$f(0) = \sqrt{0} = 0$	(0, 0)
1	$f(1) = \sqrt{1} = 1$	(1, 1)
4	$f(4) = \sqrt{4} = 2$	(4, 2)
9	$f(9) = \sqrt{9} = 3$	(9, 3)

x	$g(x) = \sqrt{x-1}$	(x, y)
1	$g(1) = \sqrt{1-1} = 0$	(1, 0)
2	$g(2) = \sqrt{2-1} = 1$	(2, 1)
5	$g(5) = \sqrt{5-1} = 2$	(5, 2)
10	$g(10) = \sqrt{10-1} = 3$	(10, 3)




The graph of g is the graph of f shifted right 1 unit.

54.

x	$f(x) = \sqrt{x}$	(x, y)
0	$f(0) = \sqrt{0} = 0$	(0, 0)
1	$f(1) = \sqrt{1} = 1$	(1, 1)
4	$f(4) = \sqrt{4} = 2$	(4, 2)
9	$f(9) = \sqrt{9} = 3$	(9, 3)

x	$g(x) = \sqrt{x+2}$	(x, y)
-2	$g(-2) = \sqrt{-2+2} = 0$	(-2, 0)
-1	$g(-1) = \sqrt{-1+2} = 1$	(-1, 1)
2	$g(2) = \sqrt{2+2} = 2$	(2, 2)
7	$g(7) = \sqrt{7+2} = 3$	(7, 3)



The graph of g is the graph of f shifted left 2 units.

55. function

56. function

57. function

58. not a function

59. not a function

60. not a function

61. function

62. not a function

63. function

64. function

65. $f(-2) = -4$

66. $f(2) = -4$

67. $f(4) = 4$

68. $f(-4) = 4$

69. $f(-3) = 0$

70. $f(-1) = 0$

71. $g(-4) = 2$

72. $g(2) = -2$

73. $g(-10) = 2$

74. $g(10) = -2$

75. When $x = -2$, $g(x) = 1$.

76. When $x = 1$, $g(x) = -1$.

77. a. domain: $(-\infty, \infty)$

b. range: $[-4, \infty)$

c. x -intercepts: -3 and 1

d. y -intercept: -3

e. $f(-2) = -3$ and $f(2) = 5$

78. a. domain: $(-\infty, \infty)$
 b. range: $(-\infty, 4]$
 c. x -intercepts: -3 and 1
 d. y -intercept: 3
 e. $f(-2) = 3$ and $f(2) = -5$
79. a. domain: $(-\infty, \infty)$
 b. range: $[1, \infty)$
 c. x -intercept: none
 d. y -intercept: 1
 e. $f(-1) = 2$ and $f(3) = 4$
80. a. domain: $(-\infty, \infty)$
 b. range: $[0, \infty)$
 c. x -intercept: -1
 d. y -intercept: 1
 e. $f(-4) = 3$ and $f(3) = 4$
81. a. domain: $[0, 5)$
 b. range: $[-1, 5)$
 c. x -intercept: 2
 d. y -intercept: -1
 e. $f(3) = 1$
82. a. domain: $(-6, 0]$
 b. range: $[-3, 4)$
 c. x -intercept: -3.75
 d. y -intercept: -3
 e. $f(-5) = 2$
83. a. domain: $[0, \infty)$
 b. range: $[1, \infty)$
 c. x -intercept: none
 d. y -intercept: 1
 e. $f(4) = 3$
84. a. domain: $[-1, \infty)$
 b. range: $[0, \infty)$
 c. x -intercept: -1
 d. y -intercept: 1
 e. $f(3) = 2$
85. a. domain: $[-2, 6]$
 b. range: $[-2, 6]$
 c. x -intercept: 4
 d. y -intercept: 4
 e. $f(-1) = 5$
86. a. domain: $[-3, 2]$
 b. range: $[-5, 5]$
 c. x -intercept: $-\frac{1}{2}$
 d. y -intercept: 1
 e. $f(-2) = -3$
87. a. domain: $(-\infty, \infty)$
 b. range: $(-\infty, -2]$
 c. x -intercept: none
 d. y -intercept: -2
 e. $f(-4) = -5$ and $f(4) = -2$
88. a. domain: $(-\infty, \infty)$
 b. range: $[0, \infty)$
 c. x -intercept: $\{x \mid x \leq 0\}$
 d. y -intercept: 0
 e. $f(-2) = 0$ and $f(2) = 4$
89. a. domain: $(-\infty, \infty)$
 b. range: $(0, \infty)$
 c. x -intercept: none
 d. y -intercept: 1.5
 e. $f(4) = 6$

90. a. domain: $(-\infty, 1) \cup (1, \infty)$

b. range: $(-\infty, 0) \cup (0, \infty)$

c. x -intercept: none

d. y -intercept: -1

e. $f(2) = 1$

91. a. domain: $\{-5, -2, 0, 1, 3\}$

b. range: $\{2\}$

c. x -intercept: none

d. y -intercept: 2

e. $f(-5) + f(3) = 2 + 2 = 4$

92. a. domain: $\{-5, -2, 0, 1, 4\}$

b. range: $\{-2\}$

c. x -intercept: none

d. y -intercept: -2

e. $f(-5) + f(4) = -2 + (-2) = -4$

93. $g(1) = 3(1) - 5 = 3 - 5 = -2$

$$f(g(1)) = f(-2) = (-2)^2 - (-2) + 4 \\ = 4 + 2 + 4 = 10$$

94. $g(-1) = 3(-1) - 5 = -3 - 5 = -8$

$$f(g(-1)) = f(-8) = (-8)^2 - (-8) + 4 \\ = 64 + 8 + 4 = 76$$

95. $\sqrt{3 - (-1)} - (-6)^2 + 6 \div (-6) \cdot 4$

$$= \sqrt{3 + 1} - 36 + 6 \div (-6) \cdot 4$$

$$= \sqrt{4} - 36 + -1 \cdot 4$$

$$= 2 - 36 + -4$$

$$= -34 + -4$$

$$= -38$$

96. $|-4 - (-1)| - (-3)^2 + -3 \div 3 \cdot -6$

$$= |-4 + 1| - 9 + -3 \div 3 \cdot -6$$

$$= |-3| - 9 + -1 \cdot -6$$

$$= 3 - 9 + 6 = -6 + 6 = 0$$

97. $f(-x) - f(x)$

$$= (-x)^3 + (-x) - 5 - (x^3 + x - 5)$$

$$= -x^3 - x - 5 - x^3 - x + 5 = -2x^3 - 2x$$

98. $f(-x) - f(x)$

$$= (-x)^2 - 3(-x) + 7 - (x^2 - 3x + 7)$$

$$= x^2 + 3x + 7 - x^2 + 3x - 7$$

$$= 6x$$

99. a. $\{(Philippines, 12), (Spain, 13),$
 $(Italy, 14), (Germany, 14),$
 $(Russia, 16)\}$

b. Yes, the relation is a function. Each country (element in the domain) corresponds to only one age (element in the range).

c. $\{(12, Philippines), (13, Spain),$
 $(14, Italy), (14, Germany),$
 $(16, Russia)\}$

d. No, the relation is not a function. 14 in the domain corresponds to two members in the range, Italy and Germany.

100. a. $\{(Philippines, 18), (Spain, 18),$
 $(Italy, 16), (Germany, 16),$
 $(Russia, 16)\}$

b. Yes, the relation is a function. Each country (element in the domain) corresponds to only one age (element in the range).

c. $\{(18, Philippines), (18, Spain),$
 $(16, Italy), (16, Germany),$
 $(16, Russia)\}$

d. No, the relation is not a function. 18 in the domain corresponds to two members of the range, Philippines and Spain, and 16 in the domain corresponds to three members of the range, Italy, Germany, and Russia.

101. a. $f(70) = 83$ which means the chance that a 60-year old will survive to age 70 is 83%.

b. $g(70) = 76$ which means the chance that a 60-year old will survive to age 70 is 76%.

c. Function f is the better model.


- 102.** a. $f(90) = 25$ which means the chance that a 60-year old will survive to age 90 is 25%.
 b. $g(90) = 10$ which means the chance that a 60-year old will survive to age 90 is 10%.
 c. Function f is the better model.
- 103.** a. $G(30) = -0.01(30)^2 + (30) + 60 = 81$
 In 2010, the wage gap was 81%. This is represented as $(30, 81)$ on the graph.
 b. $G(30)$ underestimates the actual data shown by the bar graph by 2%.
- 104.** a. $G(10) = -0.01(10)^2 + (10) + 60 = 69$
 In 1990, the wage gap was 69%. This is represented as $(10, 69)$ on the graph.
 b. $G(10)$ underestimates the actual data shown by the bar graph by 2%.
- 105.** $C(x) = 100,000 + 100x$
 $C(90) = 100,000 + 100(90) = \$109,000$
 It will cost \$109,000 to produce 90 bicycles.
- 106.** $V(x) = 22,500 - 3200x$
 $V(3) = 22,500 - 3200(3) = \$12,900$
 After 3 years, the car will be worth \$12,900.
- 107.** $T(x) = \frac{40}{x} + \frac{40}{x+30}$
 $T(30) = \frac{40}{30} + \frac{40}{30+30}$
 $= \frac{80}{60} + \frac{40}{60}$
 $= \frac{120}{60}$
 $= 2$
 If you travel 30 mph going and 60 mph returning, your total trip will take 2 hours.
- 108.** $S(x) = 0.10x + 0.60(50 - x)$
 $S(30) = 0.10(30) + 0.60(50 - 30) = 15$
 When 30 mL of the 10% mixture is mixed with 20 mL of the 60% mixture, there will be 15 mL of sodium-iodine in the vaccine.
- 109. – 117.** Answers will vary.
- 118.** makes sense
- 119.** does not make sense; Explanations will vary. Sample explanation: The parentheses used in function notation, such as $f(x)$, do not imply multiplication.
- 120.** does not make sense; Explanations will vary. Sample explanation: The domain is the number of years worked for the company.
- 121.** does not make sense; Explanations will vary. Sample explanation: This would not be a function because some elements in the domain would correspond to more than one age in the range.
- 122.** false; Changes to make the statement true will vary. A sample change is: The domain is $[-4, 4]$.
- 123.** false; Changes to make the statement true will vary. A sample change is: The range is $[-2, 2]$.
- 124.** true
- 125.** false; Changes to make the statement true will vary. A sample change is: $f(0) = 0.8$
- 126.** $f(a+h) = 3(a+h) + 7 = 3a + 3h + 7$
 $f(a) = 3a + 7$
 $\frac{f(a+h) - f(a)}{h}$
 $= \frac{(3a + 3h + 7) - (3a + 7)}{h}$
 $= \frac{3a + 3h + 7 - 3a - 7}{h} = \frac{3h}{h} = 3$
- 127.** Answers will vary. An example is $\{(1, 1), (2, 1)\}$
- 128.** It is given that $f(x+y) = f(x) + f(y)$ and $f(1) = 3$.
 To find $f(2)$, rewrite 2 as $1 + 1$.
 $f(2) = f(1+1) = f(1) + f(1)$
 $= 3 + 3 = 6$
 Similarly:
 $f(3) = f(2+1) = f(2) + f(1)$
 $= 6 + 3 = 9$
 $f(4) = f(3+1) = f(3) + f(1)$
 $= 9 + 3 = 12$
 While $f(x+y) = f(x) + f(y)$ is true for this function, it is not true for all functions. It is not true for $f(x) = x^2$, for example.

129. $C(t) = 20 + 0.40(t - 60)$

$$\begin{aligned} C(100) &= 20 + 0.40(100 - 60) \\ &= 20 + 0.40(40) \\ &= 20 + 16 \\ &= 36 \end{aligned}$$

For 100 calling minutes, the monthly cost is \$36.

130. $f(x) = x + 2, x \leq 1$



131. $2(x+h)^2 + 3(x+h) + 5 - (2x^2 + 3x + 5)$

$$\begin{aligned} &= 2(x^2 + 2xh + h^2) + 3x + 3h + 5 - 2x^2 - 3x - 5 \\ &= 2x^2 + 4xh + 2h^2 + 3x + 3h + 5 - 2x^2 - 3x - 5 \\ &= 2x^2 - 2x^2 + 4xh + 2h^2 + 3x - 3x + 3h + 5 - 5 \\ &= 4xh + 2h^2 + 3h \end{aligned}$$

Section 1.3

Check Point Exercises

1. The function is increasing on the interval $(-\infty, -1)$, decreasing on the interval $(-1, 1)$, and increasing on the interval $(1, \infty)$.

2. a. $f(-x) = (-x)^2 + 6 = x^2 + 6 = f(x)$
The function is even.

b. $g(-x) = 7(-x)^3 - (-x) = -7x^3 + x = -f(x)$
The function is odd.


c. $h(-x) = (-x)^5 + 1 = -x^5 + 1$
The function is neither even nor odd.

3. $C(t) = \begin{cases} 20 & \text{if } 0 \leq t \leq 60 \\ 20 + 0.40(t - 60) & \text{if } t > 60 \end{cases}$

b. Since $0 \leq 40 \leq 60$, $C(40) = 20$
With 40 calling minutes, the cost is \$20.
This is represented by $(40, 20)$.

c. Since $80 > 60$,
 $C(80) = 20 + 0.40(80 - 60) = 28$
With 80 calling minutes, the cost is \$28.
This is represented by $(80, 28)$.

4.



$$f(x) = \begin{cases} 3 & \text{if } x \leq -1 \\ x - 2 & \text{if } x > -1 \end{cases}$$

5. a. $f(x) = -2x^2 + x + 5$

$$\begin{aligned} f(x+h) &= -2(x+h)^2 + (x+h) + 5 \\ &= -2(x^2 + 2xh + h^2) + x + h + 5 \\ &= -2x^2 - 4xh - 2h^2 + x + h + 5 \end{aligned}$$

b. $\frac{f(x+h) - f(x)}{h}$

$$\begin{aligned} &= \frac{-2x^2 - 4xh - 2h^2 + x + h + 5 - (-2x^2 + x + 5)}{h} \\ &= \frac{-2x^2 - 4xh - 2h^2 + x + h + 5 + 2x^2 - x - 5}{h} \\ &= \frac{-4xh - 2h^2 + h}{h} \\ &= \frac{h(-4x - 2h + 1)}{h} \\ &= -4x - 2h + 1, \quad h \neq 0 \end{aligned}$$

Concept and Vocabulary Check 1.3

- $< f(x_2)$; $> f(x_2)$; $= f(x_2)$
- maximum; minimum
- $f(x)$; y-axis
- $-f(x)$; origin
- piecewise
- less than or equal to x ; 2; -3 ; 0
- difference quotient; $x + h$; $f(x)$; h ; h
- false
- false

Exercise Set 1.3

1.
 - a. increasing: $(-1, \infty)$
 - b. decreasing: $(-\infty, -1)$
 - c. constant: none
2.
 - a. increasing: $(-\infty, -1)$
 - b. decreasing: $(-1, \infty)$
 - c. constant: none
3.
 - a. increasing: $(0, \infty)$
 - b. decreasing: none
 - c. constant: none
4.
 - a. increasing: $(-1, \infty)$
 - b. decreasing: none
 - c. constant: none
5.
 - a. increasing: none
 - b. decreasing: $(-2, 6)$
 - c. constant: none
6.
 - a. increasing: $(-3, 2)$
 - b. decreasing: none
 - c. constant: none
7.
 - a. increasing: $(-\infty, -1)$
 - b. decreasing: none
 - c. constant: $(-1, \infty)$
8.
 - a. increasing: $(0, \infty)$
 - b. decreasing: none
 - c. constant: $(-\infty, 0)$
9.
 - a. increasing: $(-\infty, 0)$ or $(1.5, 3)$
 - b. decreasing: $(0, 1.5)$ or $(3, \infty)$
 - c. constant: none
10.
 - a. increasing: $(-5, -4)$ or $(-2, 0)$ or $(2, 4)$
 - b. decreasing: $(-4, -2)$ or $(0, 2)$ or $(4, 5)$
 - c. constant: none
11.
 - a. increasing: $(-2, 4)$
 - b. decreasing: none
 - c. constant: $(-\infty, -2)$ or $(4, \infty)$
12.
 - a. increasing: none
 - b. decreasing: $(-4, 2)$
 - c. constant: $(-\infty, -4)$ or $(2, \infty)$
13.
 - a. $x = 0$, relative maximum = 4
 - b. $x = -3$, 3, relative minimum = 0
14.
 - a. $x = 0$, relative maximum = 2
 - b. $x = -3$, 3, relative minimum = -1
15.
 - a. $x = -2$, relative maximum = 21
 - b. $x = 1$, relative minimum = -6
16.
 - a. $x = 1$, relative maximum = 30
 - b. $x = 4$, relative minimum = 3
17.

$$f(x) = x^3 + x$$

$$f(-x) = (-x)^3 + (-x)$$

$$f(-x) = -x^3 - x = -(x^3 + x)$$

$$f(-x) = -f(x), \text{ odd function}$$
18.

$$f(x) = x^3 - x$$

$$f(-x) = (-x)^3 - (-x)$$

$$f(-x) = -x^3 + x = -(x^3 - x)$$

$$f(-x) = -f(x), \text{ odd function}$$
19.

$$g(x) = x^2 + x$$

$$g(-x) = (-x)^2 + (-x)$$

$$g(-x) = x^2 - x, \text{ neither}$$
20.

$$g(x) = x^2 - x$$

$$g(-x) = (-x)^2 - (-x)$$

$$g(-x) = x^2 + x, \text{ neither}$$

21. $h(x) = x^2 - x^4$
 $h(-x) = (-x)^2 - (-x)^4$
 $h(-x) = x^2 - x^4$
 $h(-x) = h(x)$, even function

22. $h(x) = 2x^2 + x^4$
 $h(-x) = 2(-x)^2 + (-x)^4$
 $h(-x) = 2x^2 + x^4$
 $h(-x) = h(x)$, even function

23. $f(x) = x^2 - x^4 + 1$
 $f(-x) = (-x)^2 - (-x)^4 + 1$
 $f(-x) = x^2 - x^4 + 1$
 $f(-x) = f(x)$, even function

24. $f(x) = 2x^2 + x^4 + 1$
 $f(-x) = 2(-x)^2 + (-x)^4 + 1$
 $f(-x) = 2x^2 + x^4 + 1$
 $f(-x) = f(x)$, even function

25. $f(x) = \frac{1}{5}x^6 - 3x^2$
 $f(-x) = \frac{1}{5}(-x)^6 - 3(-x)^2$
 $f(-x) = \frac{1}{5}x^6 - 3x^2$
 $f(-x) = f(x)$, even function

26. $f(x) = 2x^3 - 6x^5$
 $f(-x) = 2(-x)^3 - 6(-x)^5$
 $f(-x) = -2x^3 + 6x^5$
 $f(-x) = -(2x^3 - 6x^5)$
 $f(-x) = -f(x)$, odd function

27. $f(x) = x\sqrt{1-x^2}$
 $f(-x) = -x\sqrt{1-(-x)^2}$
 $f(-x) = -x\sqrt{1-x^2}$
 $= -\left(x\sqrt{1-x^2}\right)$
 $f(-x) = -f(x)$, odd function

28. $f(x) = x^2\sqrt{1-x^2}$
 $f(-x) = (-x)^2\sqrt{1-(-x)^2}$
 $f(-x) = x^2\sqrt{1-x^2}$
 $f(-x) = f(x)$, even function

29. The graph is symmetric with respect to the y-axis.
The function is even.

30. The graph is symmetric with respect to the origin.
The function is odd.

31. The graph is symmetric with respect to the origin.
The function is odd.

32. The graph is not symmetric with respect to the y-axis
or the origin. The function is neither even nor odd.

33. a. domain: $(-\infty, \infty)$

b. range: $[-4, \infty)$

c. x-intercepts: 1, 7

d. y-intercept: 4

e. $(4, \infty)$

f. $(0, 4)$

g. $(-\infty, 0)$

h. $x = 4$

i. $y = -4$

j. $f(-3) = 4$

k. $f(2) = -2$ and $f(6) = -2$

l. neither ; $f(-x) \neq x$, $f(-x) \neq -x$

34. a. domain: $(-\infty, \infty)$


b. range: $(-\infty, 4]$

c. x-intercepts: -4, 4

d. y-intercept: 1


e. $(-\infty, -2)$ or $(0, 3)$

- f. $(-2, 0)$ or $(3, \infty)$
- g. $(-\infty, -4]$ or $[4, \infty)$
- h. $x = -2$ and $x = 3$
- i. $f(-2) = 4$ and $f(3) = 2$
- j. $f(-2) = 4$
- k. $x = -4$ and $x = 4$
- l. neither; $f(-x) \neq x$, $f(-x) \neq -x$
35. a. domain: $(-\infty, 3]$
- b. range: $(-\infty, 4]$
- c. x-intercepts: $-3, 3$
- d. $f(0) = 3$
- e. $(-\infty, 1)$
- f. $(1, 3)$
- g. $(-\infty, -3]$
- h. $f(1) = 4$
- i. $x = 1$
- j. positive; $f(-1) = +2$
36. a. domain: $(-\infty, 6]$
- b. range: $(-\infty, 1]$
- c. zeros of f : $-3, 3$
- d. $f(0) = 1$
- e. $(-\infty, -2)$
- f. $(2, 6)$
- g. $(-2, 2)$
- h. $(-3, 3)$
- i. $x = -5$ and $x = 5$
- j. negative; $f(4) = -1$
- k. neither
- l. no; $f(2)$ is not greater than the function values to the immediate left.
37. a. $f(-2) = 3(-2) + 5 = -1$
- b. $f(0) = 4(0) + 7 = 7$
- c. $f(3) = 4(3) + 7 = 19$
38. a. $f(-3) = 6(-3) - 1 = -19$
- b. $f(0) = 7(0) + 3 = 3$
- c. $f(4) = 7(4) + 3 = 31$
39. a. $g(0) = 0 + 3 = 3$
- b. $g(-6) = -(-6 + 3) = -(-3) = 3$
- c. $g(-3) = -3 + 3 = 0$
40. a. $g(0) = 0 + 5 = 5$
- b. $g(-6) = -(-6 + 5) = -(-1) = 1$
- c. $g(-5) = -5 + 5 = 0$
41. a. $h(5) = \frac{5^2 - 9}{5 - 3} = \frac{25 - 9}{2} = \frac{16}{2} = 8$
- b. $h(0) = \frac{0^2 - 9}{0 - 3} = \frac{-9}{-3} = 3$
- c. $h(3) = 6$
42. a. $h(7) = \frac{7^2 - 25}{7 - 5} = \frac{49 - 25}{2} = \frac{24}{2} = 12$
- b. $h(0) = \frac{0^2 - 25}{0 - 5} = \frac{-25}{-5} = 5$
- c. $h(5) = 10$
43. a.



- b. range: $[0, \infty)$


44. a.



$$f(x) = \begin{cases} x & \text{if } x < 0 \\ -x & \text{if } x \geq 0 \end{cases}$$

b. range: $(-\infty, 0]$


45. a.



$$f(x) = \begin{cases} 2x & \text{if } x \leq 0 \\ 2 & \text{if } x > 0 \end{cases}$$

b. range: $(-\infty, 0] \cup \{2\}$


46. a.



$$f(x) = \begin{cases} \frac{1}{2}x & \text{if } x \leq 0 \\ 3 & \text{if } x > 0 \end{cases}$$

b. range: $(-\infty, 0] \cup \{3\}$


47. a.



$$f(x) = \begin{cases} x + 3 & \text{if } x < -2 \\ x - 3 & \text{if } x \geq -2 \end{cases}$$

b. range: $(-\infty, \infty)$


48. a.



$$f(x) = \begin{cases} x + 2 & \text{if } x < -3 \\ x - 2 & \text{if } x \geq -3 \end{cases}$$

b. range: $(-\infty, \infty)$


49. a.



$$f(x) = \begin{cases} 3 & \text{if } x \leq -1 \\ -3 & \text{if } x > -1 \end{cases}$$

b. range: $\{-3, 3\}$


50. a.



$$f(x) = \begin{cases} 4 & \text{if } x \leq -1 \\ -4 & \text{if } x > -1 \end{cases}$$

b. range: $\{-4, 4\}$


51. a.



$$f(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } x < 1 \\ 2x - 1 & \text{if } x \geq 1 \end{cases}$$

b. range: $[0, \infty)$


52. a.



$$f(x) = \begin{cases} -\frac{1}{2}x^2 & \text{if } x < 1 \\ 2x + 1 & \text{if } x \geq 1 \end{cases}$$

b. range: $(-\infty, 0] \cup [3, \infty)$


53. a.



$$f(x) = \begin{cases} 0 & \text{if } x < -4 \\ -x & \text{if } -4 \leq x < 0 \\ x^2 & \text{if } x \geq 0 \end{cases}$$

 b. range: $[0, \infty)$

54. a.



$$f(x) = \begin{cases} 0 & \text{if } x < -3 \\ -x & \text{if } -3 \leq x < 0 \\ x^2 - 1 & \text{if } x \geq 0 \end{cases}$$

 b. range: $[-1, \infty)$

$$\begin{aligned} 55. \quad & \frac{f(x+h) - f(x)}{h} \\ &= \frac{4(x+h) - 4x}{h} \\ &= \frac{4x + 4h - 4x}{h} \\ &= \frac{4h}{h} \\ &= 4 \end{aligned}$$

$$\begin{aligned} 56. \quad & \frac{f(x+h) - f(x)}{h} \\ &= \frac{7(x+h) - 7x}{h} \\ &= \frac{7x + 7h - 7x}{h} \\ &= \frac{7h}{h} \\ &= 7 \end{aligned}$$

$$\begin{aligned} 57. \quad & \frac{f(x+h) - f(x)}{h} \\ &= \frac{3(x+h) + 7 - (3x+7)}{h} \\ &= \frac{3x + 3h + 7 - 3x - 7}{h} \\ &= \frac{3h}{h} \\ &= 3 \end{aligned}$$

$$\begin{aligned} 58. \quad & \frac{f(x+h) - f(x)}{h} \\ &= \frac{6(x+h) + 1 - (6x+1)}{h} \\ &= \frac{6x + 6h + 1 - 6x - 1}{h} \\ &= \frac{6h}{h} \\ &= 6 \end{aligned}$$

$$\begin{aligned} 59. \quad & \frac{f(x+h) - f(x)}{h} \\ &= \frac{(x+h)^2 - x^2}{h} \\ &= \frac{x^2 + 2xh + h^2 - x^2}{h} \\ &= \frac{2xh + h^2}{h} \\ &= \frac{h(2x+h)}{h} \\ &= 2x+h \end{aligned}$$

$$\begin{aligned} 60. \quad & \frac{f(x+h) - f(x)}{h} \\ &= \frac{2(x+h)^2 - 2x^2}{h} \\ &= \frac{2(x^2 + 2xh + h^2) - 2x^2}{h} \\ &= \frac{2x^2 + 4xh + 2h^2 - 2x^2}{h} \\ &= \frac{4xh + 2h^2}{h} \\ &= \frac{h(4x+2h)}{h} \\ &= 4x+2h \end{aligned}$$

$$\begin{aligned}
 61. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{(x+h)^2 - 4(x+h) + 3 - (x^2 - 4x + 3)}{h} \\
 &= \frac{x^2 + 2xh + h^2 - 4x - 4h + 3 - x^2 + 4x - 3}{h} \\
 &= \frac{2xh + h^2 - 4h}{h} \\
 &= \frac{h(2x + h - 4)}{h} \\
 &= 2x + h - 4
 \end{aligned}$$

$$\begin{aligned}
 62. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{(x+h)^2 - 5(x+h) + 8 - (x^2 - 5x + 8)}{h} \\
 &= \frac{x^2 + 2xh + h^2 - 5x - 5h + 8 - x^2 + 5x - 8}{h} \\
 &= \frac{2xh + h^2 - 5h}{h} \\
 &= \frac{h(2x + h - 5)}{h} \\
 &= 2x + h - 5
 \end{aligned}$$

$$\begin{aligned}
 63. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{2(x+h)^2 + (x+h) - 1 - (2x^2 + x - 1)}{h} \\
 &= \frac{2x^2 + 4xh + 2h^2 + x + h - 1 - 2x^2 - x + 1}{h} \\
 &= \frac{4xh + 2h^2 + h}{h} \\
 &= \frac{h(4x + 2h + 1)}{h} \\
 &= 4x + 2h + 1
 \end{aligned}$$

$$\begin{aligned}
 64. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{3(x+h)^2 + (x+h) + 5 - (3x^2 + x + 5)}{h} \\
 &= \frac{3x^2 + 6xh + 3h^2 + x + h + 5 - 3x^2 - x - 5}{h} \\
 &= \frac{6xh + 3h^2 + h}{h} \\
 &= \frac{h(6x + 3h + 1)}{h} \\
 &= 6x + 3h + 1
 \end{aligned}$$

$$\begin{aligned}
 65. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{-(x+h)^2 + 2(x+h) + 4 - (-x^2 + 2x + 4)}{h} \\
 &= \frac{-x^2 - 2xh - h^2 + 2x + 2h + 4 + x^2 - 2x - 4}{h} \\
 &= \frac{-2xh - h^2 + 2h}{h} \\
 &= \frac{h(-2x - h + 2)}{h} \\
 &= -2x - h + 2
 \end{aligned}$$

$$\begin{aligned}
 66. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{-(x+h)^2 - 3(x+h) + 1 - (-x^2 - 3x + 1)}{h} \\
 &= \frac{-x^2 - 2xh - h^2 - 3x - 3h + 1 + x^2 + 3x - 1}{h} \\
 &= \frac{-2xh - h^2 - 3h}{h} \\
 &= \frac{h(-2x - h - 3)}{h} \\
 &= -2x - h - 3
 \end{aligned}$$

$$\begin{aligned}
 67. \quad & \frac{f(x+h)-f(x)}{h} \\
 &= \frac{-2(x+h)^2 + 5(x+h) + 7 - (-2x^2 + 5x + 7)}{h} \\
 &= \frac{-2x^2 - 4xh - 2h^2 + 5x + 5h + 7 + 2x^2 - 5x - 7}{h} \\
 &= \frac{-4xh - 2h^2 + 5h}{h} \\
 &= \frac{h(-4x - 2h + 5)}{h} \\
 &= -4x - 2h + 5
 \end{aligned}$$

$$\begin{aligned}
 68. \quad & \frac{f(x+h)-f(x)}{h} \\
 &= \frac{-3(x+h)^2 + 2(x+h) - 1 - (-3x^2 + 2x - 1)}{h} \\
 &= \frac{-3x^2 - 6xh - 3h^2 + 2x + 2h - 1 + 3x^2 - 2x + 1}{h} \\
 &= \frac{-6xh - 3h^2 + 2h}{h} \\
 &= \frac{h(-6x - 3h + 2)}{h} \\
 &= -6x - 3h + 2
 \end{aligned}$$

$$\begin{aligned}
 69. \quad & \frac{f(x+h)-f(x)}{h} \\
 &= \frac{-2(x+h)^2 - (x+h) + 3 - (-2x^2 - x + 3)}{h} \\
 &= \frac{-2x^2 - 4xh - 2h^2 - x - h + 3 + 2x^2 + x - 3}{h} \\
 &= \frac{-4xh - 2h^2 - h}{h} \\
 &= \frac{h(-4x - 2h - 1)}{h} \\
 &= -4x - 2h - 1
 \end{aligned}$$

$$\begin{aligned}
 70. \quad & \frac{f(x+h)-f(x)}{h} \\
 &= \frac{-3(x+h)^2 + (x+h) - 1 - (-3x^2 + x - 1)}{h} \\
 &= \frac{-3x^2 - 6xh - 3h^2 + x + h - 1 + 3x^2 - x + 1}{h} \\
 &= \frac{-6xh - 3h^2 + h}{h} \\
 &= \frac{h(-6x - 3h + 1)}{h} \\
 &= -6x - 3h + 1
 \end{aligned}$$

$$71. \quad \frac{f(x+h)-f(x)}{h} = \frac{6-6}{h} = \frac{0}{h} = 0$$

$$72. \quad \frac{f(x+h)-f(x)}{h} = \frac{7-7}{h} = \frac{0}{h} = 0$$

$$\begin{aligned}
 73. \quad & \frac{f(x+h)-f(x)}{h} \\
 &= \frac{\frac{1}{x+h} - \frac{1}{x}}{h} \\
 &= \frac{\frac{x}{x(x+h)} + \frac{-(x+h)}{x(x+h)}}{h} \\
 &= \frac{\frac{x-x-h}{x(x+h)}}{h} \\
 &= \frac{\frac{-h}{x(x+h)}}{h} \\
 &= \frac{-h}{x(x+h)} \cdot \frac{1}{h} \\
 &= \frac{-1}{x(x+h)}
 \end{aligned}$$

$$\begin{aligned}
 74. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{\frac{1}{2(x+h)} - \frac{1}{2x}}{h} \\
 &= \frac{\frac{x}{2x(x+h)} - \frac{x+h}{2x(x+h)}}{h} \\
 &= \frac{-h}{2x(x+h)} \\
 &= \frac{-h}{2x(x+h)} \cdot \frac{1}{h} \\
 &= \frac{-1}{2x(x+h)}
 \end{aligned}$$


$$\begin{aligned}
 75. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{\sqrt{x+h} - \sqrt{x}}{h} \\
 &= \frac{\sqrt{x+h} - \sqrt{x}}{h} \cdot \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\
 &= \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\
 &= \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\
 &= \frac{1}{\sqrt{x+h} + \sqrt{x}}
 \end{aligned}$$

$$\begin{aligned}
 76. \quad & \frac{f(x+h) - f(x)}{h} \\
 &= \frac{\sqrt{x+h-1} - \sqrt{x-1}}{h} \\
 &= \frac{\sqrt{x+h-1} - \sqrt{x-1}}{h} \cdot \frac{\sqrt{x+h-1} + \sqrt{x-1}}{\sqrt{x+h-1} + \sqrt{x-1}} \\
 &= \frac{x+h-1 - (x-1)}{h(\sqrt{x+h-1} + \sqrt{x-1})} \\
 &= \frac{x+h-1-x+1}{h(\sqrt{x+h-1} + \sqrt{x-1})} \\
 &= \frac{h}{h(\sqrt{x+h-1} + \sqrt{x-1})} \\
 &= \frac{1}{\sqrt{x+h-1} + \sqrt{x-1}}
 \end{aligned}$$


$$\begin{aligned}
 77. \quad & \sqrt{f(-1.5) + f(-0.9)} - [f(\pi)]^2 + f(-3) \div f(1) \cdot f(-\pi) \\
 &= \sqrt{1+0} - [-4]^2 + 2 \div (-2) \cdot 3 \\
 &= \sqrt{1} - 16 + (-1) \cdot 3 \\
 &= 1 - 16 - 3 \\
 &= -18
 \end{aligned}$$

$$\begin{aligned}
 78. \quad & \sqrt{f(-2.5) - f(1.9)} - [f(-\pi)]^2 + f(-3) \div f(1) \cdot f(\pi) \\
 &= \sqrt{2 - (-2)} - [3]^2 + 2 \div (-2) \cdot (-4) \\
 &= \sqrt{4} - 9 + (-1)(-4) \\
 &= 2 - 9 + 4 \\
 &= -3
 \end{aligned}$$


$$79. \quad 30 + 0.30(t - 120) = 30 + 0.3t - 36 = 0.3t - 6$$




$$80. \quad 40 + 0.30(t - 200) = 40 + 0.3t - 60 = 0.3t - 20$$



$$81. \quad C(t) = \begin{cases} 50 & \text{if } 0 \leq t \leq 400 \\ 50 + 0.30(t - 400) & \text{if } t > 400 \end{cases}$$



82.
$$C(t) = \begin{cases} 60 & \text{if } 0 \leq t \leq 450 \\ 60 + 0.35(t - 450) & \text{if } t > 450 \end{cases}$$



83. increasing: (25, 55); decreasing: (55, 75)
84. increasing: (25, 65); decreasing: (65, 75)
85. The percent body fat in women reaches a maximum at age 55. This maximum is 38%.
86. The percent body fat in men reaches a maximum at age 65. This maximum is 26%.
87. domain: [25, 75]; range: [34, 38]
88. domain: [25, 75]; range: [23, 26]
89. This model describes percent body fat in men.
90. This model describes percent body fat in women.
91.
$$T(20,000) = 850 + 0.15(20,000 - 8500)$$

$$= 2575$$


 A single taxpayer with taxable income of \$20,000 owes \$2575.
92.
$$T(50,000) = 4750 + 0.25(50,000 - 34,500)$$

$$= 8625$$


 A single taxpayer with taxable income of \$50,000 owes \$8625.
93. $42,449 + 0.33(x - 174,400)$
94. $110,016.50 + 0.35(x - (x - 379,150))$
95. $f(3) = 0.85$
 The cost of mailing a first-class letter weighing 3 ounces is \$0.85.
96. $f(3.5) = 1.05$
 The cost of mailing a first-class letter weighing 3.5 ounces is \$1.05.
97. The cost to mail a letter weighing 1.5 ounces is \$0.65.

98. The cost to mail a letter weighing 1.8 ounces is \$0.65.


99. Pet Insurance




100. – 105. Answers will vary.




The number of doctor visits decreases during childhood and then increases as you get older. The minimum is (20.29, 3.99), which means that the minimum number of doctor visits, about 4, occurs at around age 20.




Increasing: $(-\infty, 1)$ or $(3, \infty)$
 Decreasing: $(1, 3)$




Increasing: $(-2, 0)$ or $(2, \infty)$
 Decreasing: $(-\infty, -2)$ or $(0, 2)$




Increasing: $(2, \infty)$
 Decreasing: $(-\infty, -2)$
 Constant: $(-2, 2)$





Increasing: $(1, \infty)$
 Decreasing: $(-\infty, 1)$




Increasing: $(0, \infty)$
 Decreasing: $(-\infty, 0)$



Increasing: $(-\infty, 0)$
 Decreasing: $(0, \infty)$



- c. Increasing: $(0, \infty)$
 Decreasing: $(-\infty, 0)$
- d. $f(x) = x^n$ is increasing from $(-\infty, \infty)$ when n is odd.



114. does not make sense; Explanations will vary.
 Sample explanation: It's possible the graph is not defined at a .

115. makes sense

116. makes sense

117. makes sense

118. answers will vary

119. answers will vary

120. a. h is even if both f and g are even or if both f and g are odd.

f and g are both even:

$$h(-x) = \frac{f(-x)}{g(-x)} = \frac{f(x)}{g(x)} = h(x)$$

f and g are both odd:

$$h(-x) = \frac{f(-x)}{g(-x)} = \frac{-f(x)}{-g(x)} = \frac{f(x)}{g(x)} = h(x)$$

- b. h is odd if f is odd and g is even or if f is even and g is odd.

f is odd and g is even:

$$h(-x) = \frac{f(-x)}{g(-x)} = \frac{-f(x)}{g(x)} = -\frac{f(x)}{g(x)} = -h(x)$$

f is even and g is odd:

$$h(-x) = \frac{f(-x)}{g(-x)} = \frac{f(x)}{-g(x)} = -\frac{f(x)}{g(x)} = -h(x)$$

121. answers will vary

$$122. \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 1}{-2 - (-3)} = \frac{3}{1} = 3$$

123. When $y = 0$:

$$4x - 3y - 6 = 0$$

$$4x - 3(0) - 6 = 0$$

$$4x - 6 = 0$$

$$4x = 6$$

$$x = \frac{3}{2}$$

The point is $\left(\frac{3}{2}, 0\right)$.

When $x = 0$:

$$4x - 3y - 6 = 0$$

$$4(0) - 3y - 6 = 0$$

$$-3y - 6 = 0$$

$$-3y = 6$$

$$y = -2$$

The point is $(0, -2)$.

$$124. 3x + 2y - 4 = 0$$

$$2y = -3x + 4$$

$$y = \frac{-3x + 4}{2}$$

or

$$y = -\frac{3}{2}x + 2$$

Section 1.4

Check Point Exercises

$$1. \quad a. \quad m = \frac{-2 - 4}{-4 - (-3)} = \frac{-6}{-1} = 6$$

$$b. \quad m = \frac{5 - (-2)}{-1 - 4} = \frac{7}{-5} = -\frac{7}{5}$$

2. Point-slope form:

$$y - y_1 = m(x - x_1)$$

$$y - (-5) = 6(x - 2)$$

$$y + 5 = 6(x - 2)$$

Slope-intercept form:

$$y + 5 = 6(x - 2)$$

$$y + 5 = 6x - 12$$

$$y = 6x - 17$$

$$3. \quad m = \frac{-6 - (-1)}{-1 - (-2)} = \frac{-5}{1} = -5,$$

so the slope is -5 .

Using the point $(-2, -1)$, we get the following point-slope equation:

$$y - y_1 = m(x - x_1)$$

$$y - (-1) = -5[x - (-2)]$$

$$y + 1 = -5(x + 2)$$

Using the point $(-1, -6)$, we get the following point-slope equation:

$$y - y_1 = m(x - x_1)$$

$$y - (-6) = -5[x - (-1)]$$

$$y + 6 = -5(x + 1)$$


Solve the equation for y :

$$y + 1 = -5(x + 2)$$


$$y + 1 = -5x - 10$$

$$y = -5x - 11.$$


4. The slope m is $\frac{3}{5}$ and the y -intercept is 1, so one point on the line is $(1, 0)$. We can find a second point on the line by using the slope $m = \frac{3}{5} = \frac{\text{Rise}}{\text{Run}}$: starting at the point $(0, 1)$, move 3 units up and 5 units to the right, to obtain the point $(5, 4)$.



5. $y = 3$ is a horizontal line.



6. All ordered pairs that are solutions of $x = -3$ have a value of x that is always -3 . Any value can be used for y .




7. $3x + 6y - 12 = 0$

$$6y = -3x + 12$$

$$y = \frac{-3}{6}x + \frac{12}{6}$$

$$y = -\frac{1}{2}x + 2$$



$$3x + 6y - 12 = 0$$

The slope is $-\frac{1}{2}$ and the y-intercept is 2.

8. Find the x-intercept:

$$3x - 2y - 6 = 0$$

$$3x - 2(0) - 6 = 0$$

$$3x - 6 = 0$$

$$3x = 6$$

$$x = 2$$

Find the y-intercept:


$$3x - 2y - 6 = 0$$

$$3(0) - 2y - 6 = 0$$

$$-2y - 6 = 0$$

$$-2y = 6$$

$$y = -3$$



$$3x - 2y = 6$$

9. First find the slope.

$$m = \frac{\text{Change in } y}{\text{Change in } x} = \frac{57.64 - 57.04}{354 - 317} = \frac{0.6}{37} \approx 0.016$$

Use the point-slope form and then find slope-intercept form.

$$y - y_1 = m(x - x_1)$$

$$y - 57.04 = 0.016(x - 317)$$

$$y - 57.04 = 0.016x - 5.072$$

$$y = 0.016x + 51.968$$

$$f(x) = 0.016x + 52.0$$

Find the temperature at a concentration of 600 parts per million.

$$f(x) = 0.016x + 52.0$$

$$f(600) = 0.016(600) + 52.0$$

$$= 61.6$$

The temperature at a concentration of 600 parts per million would be 61.6°F.

Concept and Vocabulary Check 1.4

1. scatter plot; regression

2. $\frac{y_2 - y_1}{x_2 - x_1}$

3. positive

4. negative

5. zero

6. undefined

7. $y - y_1 = m(x - x_1)$

8. $y = mx + b$; slope; y-intercept

9. (0, 3); 2; 5

10. horizontal

11. vertical

12. general

Exercise Set 1.4

1. $m = \frac{10 - 7}{8 - 4} = \frac{3}{4}$; rises

2. $m = \frac{4 - 1}{3 - 2} = \frac{3}{1} = 3$; rises

3. $m = \frac{2 - 1}{2 - (-2)} = \frac{1}{4}$; rises

4. $m = \frac{4 - 3}{2 - (-1)} = \frac{1}{3}$; rises

5. $m = \frac{2 - (-2)}{3 - 4} = \frac{0}{-1} = 0$; horizontal

6. $m = \frac{-1 - (-1)}{3 - 4} = \frac{0}{-1} = 0$; horizontal
7. $m = \frac{-1 - 4}{-1 - (-2)} = \frac{-5}{1} = -5$; falls
8. $m = \frac{-2 - (-4)}{4 - 6} = \frac{2}{-2} = -1$; falls
9. $m = \frac{-2 - 3}{5 - 5} = \frac{-5}{0}$ undefined; vertical
10. $m = \frac{5 - (-4)}{3 - 3} = \frac{9}{0}$ undefined; vertical
11. $m = 2, x_1 = 3, y_1 = 5$;
point-slope form: $y - 5 = 2(x - 3)$;
slope-intercept form: $y - 5 = 2x - 6$
 $y = 2x - 1$
12. point-slope form: $y - 3 = 4(x - 1)$;
 $m = 4, x_1 = 1, y_1 = 3$;
slope-intercept form: $y = 4x - 1$
13. $m = 6, x_1 = -2, y_1 = 5$;
point-slope form: $y - 5 = 6(x + 2)$;
slope-intercept form: $y - 5 = 6x + 12$
 $y = 6x + 17$
14. point-slope form: $y + 1 = 8(x - 4)$;
 $m = 8, x_1 = 4, y_1 = -1$;
slope-intercept form: $y = 8x - 33$
15. $m = -3, x_1 = -2, y_1 = -3$;
point-slope form: $y + 3 = -3(x + 2)$;
slope-intercept form: $y + 3 = -3x - 6$
 $y = -3x - 9$
16. point-slope form: $y + 2 = -5(x + 4)$;
 $m = -5, x_1 = -4, y_1 = -2$;
slope-intercept form: $y = -5x - 22$
17. $m = -4, x_1 = -4, y_1 = 0$;
point-slope form: $y - 0 = -4(x + 4)$;
slope-intercept form: $y = -4(x + 4)$
 $y = -4x - 16$
18. point-slope form: $y + 3 = -2(x - 0)$
 $m = -2, x_1 = 0, y_1 = -3$;
slope-intercept form: $y = -2x - 3$
19. $m = -1, x_1 = \frac{-1}{2}, y_1 = -2$;
point-slope form: $y + 2 = -1\left(x + \frac{1}{2}\right)$;
slope-intercept form: $y + 2 = -x - \frac{1}{2}$
 $y = -x - \frac{5}{2}$
20. point-slope form: $y + \frac{1}{4} = -1(x + 4)$;
 $m = -1, x_1 = -4, y_1 = -\frac{1}{4}$;
slope-intercept form: $y = -x - \frac{17}{4}$
21. $m = \frac{1}{2}, x_1 = 0, y_1 = 0$;
point-slope form: $y - 0 = \frac{1}{2}(x - 0)$;
slope-intercept form: $y = \frac{1}{2}x$
22. point-slope form: $y - 0 = \frac{1}{3}(x - 0)$;
 $m = \frac{1}{3}, x_1 = 0, y_1 = 0$;
slope-intercept form: $y = \frac{1}{3}x$
23. $m = -\frac{2}{3}, x_1 = 6, y_1 = -2$;
point-slope form: $y + 2 = -\frac{2}{3}(x - 6)$;
slope-intercept form: $y + 2 = -\frac{2}{3}x + 4$
 $y = -\frac{2}{3}x + 2$
24. point-slope form: $y + 4 = -\frac{3}{5}(x - 10)$;
 $m = -\frac{3}{5}, x_1 = 10, y_1 = -4$;
slope-intercept form: $y = -\frac{3}{5}x + 2$

$$25. \quad m = \frac{10-2}{5-1} = \frac{8}{4} = 2;$$

point-slope form: $y - 2 = 2(x - 1)$ using
 $(x_1, y_1) = (1, 2)$, or $y - 10 = 2(x - 5)$ using

$(x_1, y_1) = (5, 10)$;

slope-intercept form: $y - 2 = 2x - 2$ or

$$y - 10 = 2x - 10,$$

$$y = 2x$$

$$26. \quad m = \frac{15-5}{8-3} = \frac{10}{5} = 2;$$

point-slope form: $y - 5 = 2(x - 3)$ using

$(x_1, y_1) = (3, 5)$, or $y - 15 = 2(x - 8)$ using

$(x_1, y_1) = (8, 15)$;

slope-intercept form: $y = 2x - 1$

$$27. \quad m = \frac{3-0}{0-(-3)} = \frac{3}{3} = 1;$$

point-slope form: $y - 0 = 1(x + 3)$ using

$(x_1, y_1) = (-3, 0)$, or $y - 3 = 1(x - 0)$ using

$(x_1, y_1) = (0, 3)$; slope-intercept form: $y = x + 3$

$$28. \quad m = \frac{2-0}{0-(-2)} = \frac{2}{2} = 1;$$

point-slope form: $y - 0 = 1(x + 2)$ using

$(x_1, y_1) = (-2, 0)$, or $y - 2 = 1(x - 0)$ using

$(x_1, y_1) = (0, 2)$;

slope-intercept form: $y = x + 2$

$$29. \quad m = \frac{4-(-1)}{2-(-3)} = \frac{5}{5} = 1;$$

point-slope form: $y + 1 = 1(x + 3)$ using

$(x_1, y_1) = (-3, -1)$, or $y - 4 = 1(x - 2)$ using

$(x_1, y_1) = (2, 4)$; slope-intercept form:

$$y + 1 = x + 3 \text{ or}$$

$$y - 4 = x - 2$$

$$y = x + 2$$

$$30. \quad m = \frac{-1-(-4)}{1-(-2)} = \frac{3}{3} = 1;$$

point-slope form: $y + 4 = 1(x + 2)$ using

$(x_1, y_1) = (-2, -4)$, or $y + 1 = 1(x - 1)$ using

$(x_1, y_1) = (1, -1)$

slope-intercept form: $y = x - 2$

$$31. \quad m = \frac{6-(-2)}{3-(-3)} = \frac{8}{6} = \frac{4}{3};$$

point-slope form: $y + 2 = \frac{4}{3}(x + 3)$ using

$(x_1, y_1) = (-3, -2)$, or $y - 6 = \frac{4}{3}(x - 3)$ using

$(x_1, y_1) = (3, 6)$;

slope-intercept form: $y + 2 = \frac{4}{3}x + 4$ or

$$y - 6 = \frac{4}{3}x - 4,$$

$$y = \frac{4}{3}x + 2$$

$$32. \quad m = \frac{-2-6}{3-(-3)} = \frac{-8}{6} = -\frac{4}{3};$$

point-slope form: $y - 6 = -\frac{4}{3}(x + 3)$ using

$(x_1, y_1) = (-3, 6)$, or $y + 2 = -\frac{4}{3}(x - 3)$ using

$(x_1, y_1) = (3, -2)$;

slope-intercept form: $y = -\frac{4}{3}x + 2$

$$33. \quad m = \frac{-1-(-1)}{4-(-3)} = \frac{0}{7} = 0;$$

point-slope form: $y + 1 = 0(x + 3)$ using

$(x_1, y_1) = (-3, -1)$, or $y + 1 = 0(x - 4)$ using

$(x_1, y_1) = (4, -1)$;

slope-intercept form: $y + 1 = 0$, so

$$y = -1$$

$$34. \quad m = \frac{-5-(-5)}{6-(-2)} = \frac{0}{8} = 0;$$

point-slope form: $y + 5 = 0(x + 2)$ using

$(x_1, y_1) = (-2, -5)$, or $y + 5 = 0(x - 6)$ using

$(x_1, y_1) = (6, -5)$;

slope-intercept form: $y + 5 = 0$, so

$$y = -5$$

$$35. \quad m = \frac{0-4}{-2-2} = \frac{-4}{-4} = 1;$$

point-slope form: $y - 4 = 1(x - 2)$ using

$(x_1, y_1) = (2, 4)$, or $y - 0 = 1(x + 2)$ using

$(x_1, y_1) = (-2, 0)$;

slope-intercept form: $y - 9 = x - 2$, or

$$y = x + 2$$

$$36. \quad m = \frac{0 - (-3)}{-1 - 1} = \frac{3}{-2} = -\frac{3}{2}$$

point-slope form: $y + 3 = -\frac{3}{2}(x - 1)$ using

$$(x_1, y_1) = (1, -3), \text{ or } y - 0 = -\frac{3}{2}(x + 1) \text{ using}$$

$$(x_1, y_1) = (-1, 0);$$

slope-intercept form: $y + 3 = -\frac{3}{2}x + \frac{3}{2}$, or

$$y = -\frac{3}{2}x - \frac{3}{2}$$

$$37. \quad m = \frac{4 - 0}{0 - (-\frac{1}{2})} = \frac{4}{\frac{1}{2}} = 8;$$

point-slope form: $y - 4 = 8(x - 0)$ using

$$(x_1, y_1) = (0, 4), \text{ or } y - 0 = 8(x + \frac{1}{2}) \text{ using}$$

$$(x_1, y_1) = (-\frac{1}{2}, 0); \text{ or } y - 0 = 8(x + \frac{1}{2})$$

slope-intercept form: $y = 8x + 4$

$$38. \quad m = \frac{-2 - 0}{0 - 4} = \frac{-2}{-4} = \frac{1}{2};$$


point-slope form: $y - 0 = \frac{1}{2}(x - 4)$ using

$$(x_1, y_1) = (4, 0),$$


or $y + 2 = \frac{1}{2}(x - 0)$ using $(x_1, y_1) = (0, -2);$

slope-intercept form: $y = \frac{1}{2}x - 2$


$$39. \quad m = 2; b = 1$$




$$40. \quad m = 3; b = 2$$




$$41. \quad m = -2; b = 1$$




$$42. \quad m = -3; b = 2$$




$$43. \quad m = \frac{3}{4}; b = -2$$




$$44. \quad m = \frac{3}{4}; b = -3$$




$$45. \quad m = -\frac{3}{5}; b = 7$$












46. $m = -\frac{2}{5}; b = 6$



47. $m = -\frac{1}{2}; b = 0$




48. $m = -\frac{1}{3}; b = 0$



57. $3x - 18 = 0$

$3x = 18$


$x = 6$



58. $3x + 12 = 0$

$3x = -12$

$x = -4$




59. a. $3x + y - 5 = 0$

$y - 5 = -3x$

$y = -3x + 5$

b. $m = -3; b = 5$

c.




60. a. $4x + y - 6 = 0$

$y - 6 = -4x$

$y = -4x + 6$

b. $m = -4; b = 6$

c.



61. a. $2x + 3y - 18 = 0$

$2x - 18 = -3y$


$-3y = 2x - 18$

$y = \frac{2}{-3}x - \frac{18}{-3}$

$y = -\frac{2}{3}x + 6$

b. $m = -\frac{2}{3}; b = 6$

c.



62. a. $4x + 6y + 12 = 0$

$4x + 12 = -6y$


$-6y = 4x + 12$

$y = \frac{4}{-6}x + \frac{12}{-6}$

$y = -\frac{2}{3}x - 2$

b. $m = -\frac{2}{3}; b = -2$

c.



63. a. $8x - 4y - 12 = 0$

$8x - 12 = 4y$


$4y = 8x - 12$

$y = \frac{8}{4}x - \frac{12}{4}$

$y = 2x - 3$

b. $m = 2; b = -3$

c.



64. a. $6x - 5y - 20 = 0$

$$6x - 20 = 5y$$


$$5y = 6x - 20$$

$$y = \frac{6}{5}x - \frac{20}{5}$$

$$y = \frac{6}{5}x - 4$$

b. $m = \frac{6}{5}; b = -4$

c.




65. a. $3y - 9 = 0$

$$3y = 9$$

$$y = 3$$

b. $m = 0; b = 3$

c.




66. a. $4y + 28 = 0$

$$4y = -28$$

$$y = -7$$

b. $m = 0; b = -7$

c.



67. Find the x -intercept:

$$6x - 2y - 12 = 0$$

$$6x - 2(0) - 12 = 0$$

$$6x - 12 = 0$$

$$6x = 12$$

$$x = 2$$

Find the y -intercept:


$$6x - 2y - 12 = 0$$

$$6(0) - 2y - 12 = 0$$

$$-2y - 12 = 0$$

$$-2y = 12$$

$$y = -6$$



$6x - 2y - 12 = 0$

68. Find the x -intercept:

$$6x - 9y - 18 = 0$$

$$6x - 9(0) - 18 = 0$$

$$6x - 18 = 0$$

$$6x = 18$$

$$x = 3$$

Find the y -intercept:


$$6x - 9y - 18 = 0$$

$$6(0) - 9y - 18 = 0$$

$$-9y - 18 = 0$$

$$-9y = 18$$

$$y = -2$$



69. Find the
- x
- intercept:

$$2x + 3y + 6 = 0$$

$$2x + 3(0) + 6 = 0$$

$$2x + 6 = 0$$

$$2x = -6$$

$$x = -3$$

- Find the
- y
- intercept:


$$2x + 3y + 6 = 0$$

$$2(0) + 3y + 6 = 0$$

$$3y + 6 = 0$$

$$3y = -6$$

$$y = -2$$



70. Find the
- x
- intercept:

$$3x + 5y + 15 = 0$$

$$3x + 5(0) + 15 = 0$$

$$3x + 15 = 0$$

$$3x = -15$$

$$x = -5$$

- Find the
- y
- intercept:


$$3x + 5y + 15 = 0$$

$$3(0) + 5y + 15 = 0$$

$$5y + 15 = 0$$

$$5y = -15$$

$$y = -3$$



71. Find the
- x
- intercept:

$$8x - 2y + 12 = 0$$

$$8x - 2(0) + 12 = 0$$

$$8x + 12 = 0$$

$$8x = -12$$

$$\frac{8x}{8} = \frac{-12}{8}$$

$$x = \frac{-3}{2}$$

- Find the
- y
- intercept:


$$8x - 2y + 12 = 0$$

$$8(0) - 2y + 12 = 0$$

$$-2y + 12 = 0$$

$$-2y = -12$$

$$y = -6$$



72. Find the
- x
- intercept:

$$6x - 3y + 15 = 0$$

$$6x - 3(0) + 15 = 0$$

$$6x + 15 = 0$$

$$6x = -15$$

$$\frac{6x}{6} = \frac{-15}{6}$$

$$x = -\frac{5}{2}$$

- Find the
- y
- intercept:


$$6x - 3y + 15 = 0$$

$$6(0) - 3y + 15 = 0$$

$$-3y + 15 = 0$$

$$-3y = -15$$

$$y = 5$$



$$73. \quad m = \frac{0-a}{b-0} = \frac{-a}{b} = -\frac{a}{b}$$

Since a and b are both positive, $-\frac{a}{b}$ is negative. Therefore, the line falls.

$$74. \quad m = \frac{-b-0}{0-(-a)} = \frac{-b}{a} = -\frac{b}{a}$$

Since a and b are both positive, $-\frac{b}{a}$ is negative. Therefore, the line falls.

$$75. \quad m = \frac{(b+c)-b}{a-a} = \frac{c}{0}$$

The slope is undefined.
The line is vertical.

$$76. \quad m = \frac{(a+c)-c}{a-(a-b)} = \frac{a}{b}$$

Since a and b are both positive, $\frac{a}{b}$ is positive.
Therefore, the line rises.

$$77. \quad Ax + By = C$$

$$By = -Ax + C$$

$$y = -\frac{A}{B}x + \frac{C}{B}$$

The slope is $-\frac{A}{B}$ and the y -intercept is $\frac{C}{B}$.

$$78. \quad Ax = By - C$$

$$Ax + C = By$$

$$\frac{A}{B}x + \frac{C}{B} = y$$

The slope is $\frac{A}{B}$ and the y -intercept is $\frac{C}{B}$.

$$79. \quad -3 = \frac{4-y}{1-3}$$

$$-3 = \frac{4-y}{-2}$$

$$6 = 4 - y$$

$$2 = -y$$

$$-2 = y$$

$$80. \quad \frac{1}{3} = \frac{-4-y}{4-(-2)}$$

$$\frac{1}{3} = \frac{-4-y}{4+2}$$

$$\frac{1}{3} = \frac{-4-y}{6}$$

$$6 = 3(-4-y)$$

$$6 = -12 - 3y$$


$$18 = -3y$$

$$-6 = y$$

$$81. \quad 3x - 4f(x) = 6$$

$$-4f(x) = -3x + 6$$


$$f(x) = \frac{3}{4}x - \frac{3}{2}$$



$$82. \quad 6x - 5f(x) = 20$$

$$-5f(x) = -6x + 20$$

$$f(x) = \frac{6}{5}x - 4$$



83. Using the slope-intercept form for the equation of a line:

$$-1 = -2(3) + b$$

$$-1 = -6 + b$$

$$5 = b$$

$$84. \quad -6 = -\frac{3}{2}(2) + b$$

$$-6 = -3 + b$$

$$-3 = b$$

$$85. \quad m_1, m_3, m_2, m_4$$

86. b_2, b_1, b_4, b_3

87. a. First, find the slope using
- $(20, 38.9)$
- and
- $(10, 31.1)$
- .

$$m = \frac{38.9 - 31.1}{20 - 10} = \frac{7.8}{10} = 0.78$$

Then use the slope and one of the points to write the equation in point-slope form.

$$y - y_1 = m(x - x_1)$$

$$y - 31.1 = 0.78(x - 10)$$

or

$$y - 38.9 = 0.78(x - 20)$$

b. $y - 31.1 = 0.78(x - 10)$

$$y - 31.1 = 0.78x - 7.8$$

$$y = 0.78x + 23.3$$

$$f(x) = 0.78x + 23.3$$

c. $f(40) = 0.78(40) + 23.3 = 54.5$

The linear function predicts the percentage of never married American females, ages 25 – 29, to be 54.5% in 2020.

88. a. First, find the slope using
- $(20, 51.7)$
- and
- $(10, 45.2)$
- .

$$m = \frac{51.7 - 45.2}{20 - 10} = \frac{6.5}{10} = 0.65$$

Then use the slope and one of the points to write the equation in point-slope form.

$$y - y_1 = m(x - x_1)$$

$$y - 45.2 = 0.65(x - 10)$$

or

$$y - 51.7 = 0.65(x - 20)$$

b. $y - 45.2 = 0.65(x - 10)$

$$y - 45.2 = 0.65x - 6.5$$


$$y = 0.65x + 38.7$$

$$f(x) = 0.65x + 38.7$$

c. $f(35) = 0.65(35) + 38.7 = 61.45$

The linear function predicts the percentage of never married American males, ages 25 – 29, to be 61.45% in 2015.

89. a. Life Expectancy for United States Males, by Year of Birth



b. $m = \frac{\text{Change in } y}{\text{Change in } x} = \frac{74.3 - 70.0}{40 - 20} = 0.215$

$$y - y_1 = m(x - x_1)$$

$$y - 70.0 = 0.215(x - 20)$$

$$y - 70.0 = 0.215x - 4.3$$

$$y = 0.215x + 65.7$$

$$E(x) = 0.215x + 65.7$$


c. $E(x) = 0.215x + 65.7$

$$E(60) = 0.215(60) + 65.7$$

$$= 78.6$$

The life expectancy of American men born in 2020 is expected to be 78.6.

90. a. Life Expectancy for United States Females, by Year of Birth



b. $m = \frac{\text{Change in } y}{\text{Change in } x} = \frac{79.7 - 74.7}{40 - 10} \approx 0.17$

$$y - y_1 = m(x - x_1)$$

$$y - 74.7 = 0.17(x - 10)$$

$$y - 74.7 = 0.17x - 1.7$$

$$y = 0.17x + 73$$

$$E(x) = 0.17x + 73$$

c. $E(x) = 0.17x + 73$

$$E(60) = 0.17(60) + 73$$

$$= 83.2$$

The life expectancy of American women born in 2020 is expected to be 83.2.

91. (10, 230) (60, 110) Points may vary.

$$m = \frac{110 - 230}{60 - 10} = -\frac{120}{50} = -2.4$$

$$y - 230 = -2.4(x - 10)$$

$$y - 230 = -2.4x + 24$$


$$y = -2.4x + 254$$

Answers will vary for predictions.

92. – 99. Answers will vary.

100. Two points are (0, 4) and (10, 24).


$$m = \frac{24 - 4}{10 - 0} = \frac{20}{10} = 2.$$



101. Two points are (0, 6) and (10, -24).


$$m = \frac{-24 - 6}{10 - 0} = \frac{-30}{10} = -3.$$

Check: $y = mx + b$: $y = -3x + 6$.



102. Two points are (0, -5) and (10, -10).


$$m = \frac{-10 - (-5)}{10 - 0} = \frac{-5}{10} = -\frac{1}{2}.$$



103. Two points are (0, -2) and (10, 5.5).


$$m = \frac{5.5 - (-2)}{10 - 0} = \frac{7.5}{10} = 0.75 \text{ or } \frac{3}{4}.$$

Check: $y = mx + b$: $y = \frac{3}{4}x - 2$.



104. a. Enter data from table.

b.




- c. $a = -22.96876741$

$$b = 260.5633751$$

$$r = -0.8428126855$$

d.



105. does not make sense; Explanations will vary.
Sample explanation: Linear functions never change from increasing to decreasing.

106. does not make sense; Explanations will vary.
Sample explanation: Since college cost are going up, this function has a positive slope.

107. does not make sense; Explanations will vary.
Sample explanation: The slope of line's whose equations are in this form can be determined in several ways. One such way is to rewrite the equation in slope-intercept form.

108. makes sense

109. false; Changes to make the statement true will vary.
A sample change is: It is possible for m to equal b .

110. false; Changes to make the statement true will vary.
A sample change is: Slope-intercept form is $y = mx + b$. Vertical lines have equations of the form $x = a$. Equations of this form have undefined slope and cannot be written in slope-intercept form.

111. true

112. false; Changes to make the statement true will vary.
A sample change is: The graph of $x = 7$ is a vertical line through the point (7, 0).

- 113.** We are given that the x -intercept is -2 and the y -intercept is 4 . We can use the points $(-2, 0)$ and $(0, 4)$ to find the slope.

$$m = \frac{4 - 0}{0 - (-2)} = \frac{4}{0 + 2} = \frac{4}{2} = 2$$

Using the slope and one of the intercepts, we can write the line in point-slope form.

$$y - y_1 = m(x - x_1)$$

$$y - 0 = 2(x - (-2))$$

$$y = 2(x + 2)$$

$$y = 2x + 4$$

$$-2x + y = 4$$

Find the x - and y -coefficients for the equation of the line with right-hand-side equal to 12. Multiply both sides of $-2x + y = 4$ by 3 to obtain 12 on the right-hand-side.

$$-2x + y = 4$$

$$3(-2x + y) = 3(4)$$

$$-6x + 3y = 12$$

Therefore, the coefficient of x is -6 and the coefficient of y is 3 .

- 114.** We are given that the y -intercept is -6 and the slope is $\frac{1}{2}$.

So the equation of the line is $y = \frac{1}{2}x - 6$.

We can put this equation in the form $ax + by = c$ to find the missing coefficients.

$$y = \frac{1}{2}x - 6$$

$$y - \frac{1}{2}x = -6$$

$$2\left(y - \frac{1}{2}x\right) = 2(-6)$$

$$2y - x = -12$$

$$x - 2y = 12$$

Therefore, the coefficient of x is 1 and the coefficient of y is -2 .

- 115.** Answers will vary.

- 116.** Let $(25, 40)$ and $(125, 280)$ be ordered pairs (M, E) where M is degrees Madonna and E is degrees Elvis. Then

$$m = \frac{280 - 40}{125 - 25} = \frac{240}{100} = 2.4. \text{ Using } (x_1, y_1) = (25, 40),$$

point-slope form tells us that

$$E - 40 = 2.4(M - 25) \text{ or}$$

$$E = 2.4M - 20.$$

- 117.** Answers will vary.

- 118.** Since the slope is the same as the slope of $y = 2x + 1$, then $m = 2$.

$$y - y_1 = m(x - x_1)$$

$$y - 1 = 2(x - (-3))$$

$$y - 1 = 2(x + 3)$$

$$y - 1 = 2x + 6$$

$$y = 2x + 7$$

- 119.** Since the slope is the negative reciprocal of $-\frac{1}{4}$,

then $m = 4$.

$$y - y_1 = m(x - x_1)$$

$$y - (-5) = 4(x - 3)$$

$$y + 5 = 4x - 12$$

$$-4x + y + 17 = 0$$

$$4x - y - 17 = 0$$

$$\begin{aligned} \mathbf{120.} \quad \frac{f(x_2) - f(x_1)}{x_2 - x_1} &= \frac{f(4) - f(1)}{4 - 1} \\ &= \frac{4^2 - 1^2}{4 - 1} \\ &= \frac{15}{3} \\ &= 5 \end{aligned}$$