

# INSTRUCTOR'S SOLUTIONS MANUAL

## PROBABILITY AND STATISTICAL INFERENCE

EIGHTH EDITION

Robert V. Hogg

*University of Iowa*

Elliot A. Tanis

*Hope College*

Prentice Hall  
is an imprint of

PEARSON



**This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.**

The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expresses or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson Prentice Hall from electronic files supplied by the author.

Copyright ©2010 Pearson Education, Inc.

Publishing as Pearson Prentice Hall, Upper Saddle River, NJ 07458.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.

ISBN-13: 978-0-321-58476-2

ISBN-10: 0-321-58476-7

**Prentice Hall**  
is an imprint of



[www.pearsonhighered.com](http://www.pearsonhighered.com)

# Contents

|                                                                     |           |
|---------------------------------------------------------------------|-----------|
| <b>Preface</b>                                                      | <b>v</b>  |
| <b>1 Probability</b>                                                | <b>1</b>  |
| 1.1 Basic Concepts . . . . .                                        | 1         |
| 1.2 Properties of Probability . . . . .                             | 2         |
| 1.3 Methods of Enumeration . . . . .                                | 3         |
| 1.4 Conditional Probability . . . . .                               | 4         |
| 1.5 Independent Events . . . . .                                    | 6         |
| 1.6 Bayes's Theorem . . . . .                                       | 7         |
| <b>2 Discrete Distributions</b>                                     | <b>11</b> |
| 2.1 Random Variables of the Discrete Type . . . . .                 | 11        |
| 2.2 Mathematical Expectation . . . . .                              | 15        |
| 2.3 The Mean, Variance, and Standard Deviation . . . . .            | 16        |
| 2.4 Bernoulli Trials and the Binomial Distribution . . . . .        | 19        |
| 2.5 The Moment-Generating Function . . . . .                        | 22        |
| 2.6 The Poisson Distribution . . . . .                              | 24        |
| <b>3 Continuous Distributions</b>                                   | <b>27</b> |
| 3.1 Continuous-Type Data . . . . .                                  | 27        |
| 3.2 Exploratory Data Analysis . . . . .                             | 30        |
| 3.3 Random Variables of the Continuous Type . . . . .               | 37        |
| 3.4 The Uniform and Exponential Distributions . . . . .             | 45        |
| 3.5 The Gamma and Chi-Square Distributions . . . . .                | 48        |
| 3.6 The Normal Distribution . . . . .                               | 50        |
| 3.7 Additional Models . . . . .                                     | 54        |
| <b>4 Bivariate Distributions</b>                                    | <b>57</b> |
| 4.1 Distributions of Two Random Variables . . . . .                 | 57        |
| 4.2 The Correlation Coefficient . . . . .                           | 59        |
| 4.3 Conditional Distributions . . . . .                             | 61        |
| 4.4 The Bivariate Normal Distribution . . . . .                     | 66        |
| <b>5 Distributions of Functions of Random Variables</b>             | <b>69</b> |
| 5.1 Functions of One Random Variable . . . . .                      | 69        |
| 5.2 Transformations of Two Random Variables . . . . .               | 73        |
| 5.3 Several Independent Random Variables . . . . .                  | 76        |
| 5.4 The Moment-Generating Function Technique . . . . .              | 79        |
| 5.5 Random Functions Associated with Normal Distributions . . . . . | 81        |
| 5.6 The Central Limit Theorem . . . . .                             | 84        |
| 5.7 Approximations for Discrete Distributions . . . . .             | 86        |

|           |                                                                        |            |
|-----------|------------------------------------------------------------------------|------------|
| <b>6</b>  | <b>Estimation</b>                                                      | <b>91</b>  |
| 6.1       | Point Estimation . . . . .                                             | 91         |
| 6.2       | Confidence Intervals for Means . . . . .                               | 94         |
| 6.3       | Confidence Intervals for the Difference of Two Means . . . . .         | 95         |
| 6.4       | Confidence Intervals for Variances . . . . .                           | 97         |
| 6.5       | Confidence Intervals for Proportions . . . . .                         | 99         |
| 6.6       | Sample Size . . . . .                                                  | 100        |
| 6.7       | A Simple Regression Problem . . . . .                                  | 101        |
| 6.8       | More Regression . . . . .                                              | 107        |
| <b>7</b>  | <b>Tests of Statistical Hypotheses</b>                                 | <b>115</b> |
| 7.1       | Tests about Proportions . . . . .                                      | 115        |
| 7.2       | Tests about One Mean . . . . .                                         | 117        |
| 7.3       | Tests of the Equality of Two Means . . . . .                           | 120        |
| 7.4       | Tests for Variances . . . . .                                          | 123        |
| 7.5       | One-Factor Analysis of Variance . . . . .                              | 124        |
| 7.6       | Two-Factor Analysis of Variance . . . . .                              | 127        |
| 7.7       | Tests Concerning Regression and Correlation . . . . .                  | 128        |
| <b>8</b>  | <b>Nonparametric Methods</b>                                           | <b>131</b> |
| 8.1       | Chi-Square Goodness-of-Fit Tests . . . . .                             | 131        |
| 8.2       | Contingency Tables . . . . .                                           | 135        |
| 8.3       | Order Statistics . . . . .                                             | 136        |
| 8.4       | Distribution-Free Confidence Intervals for Percentiles . . . . .       | 138        |
| 8.5       | The Wilcoxon Tests . . . . .                                           | 140        |
| 8.6       | Run Test and Test for Randomness . . . . .                             | 144        |
| 8.7       | Kolmogorov-Smirnov Goodness of Fit Test . . . . .                      | 147        |
| 8.8       | Resampling Methods . . . . .                                           | 149        |
| <b>9</b>  | <b>Bayesian Methods</b>                                                | <b>157</b> |
| 9.1       | Subjective Probability . . . . .                                       | 157        |
| 9.2       | Bayesian Estimation . . . . .                                          | 158        |
| 9.3       | More Bayesian Concepts . . . . .                                       | 159        |
| <b>10</b> | <b>Some Theory</b>                                                     | <b>161</b> |
| 10.1      | Sufficient Statistics . . . . .                                        | 161        |
| 10.2      | Power of a Statistical Test . . . . .                                  | 162        |
| 10.3      | Best Critical Regions . . . . .                                        | 166        |
| 10.4      | Likelihood Ratio Tests . . . . .                                       | 168        |
| 10.5      | Chebyshev's Inequality and Convergence in Probability . . . . .        | 169        |
| 10.6      | Limiting Moment-Generating Functions . . . . .                         | 170        |
| 10.7      | Asymptotic Distributions of Maximum<br>Likelihood Estimators . . . . . | 171        |
| <b>11</b> | <b>Quality Improvement Through Statistical Methods</b>                 | <b>173</b> |
| 11.1      | Time Sequences . . . . .                                               | 173        |
| 11.2      | Statistical Quality Control . . . . .                                  | 176        |
| 11.3      | General Factorial and $2^k$ Factorial Designs . . . . .                | 179        |

# Preface

This solutions manual provides answers for the even-numbered exercises in *Probability and Statistical Inference*, 8th edition, by Robert V. Hogg and Elliot A. Tanis. Complete solutions are given for most of these exercises. You, the instructor, may decide how many of these answers you want to make available to your students. Note that the answers for the odd-numbered exercises are given in the textbook.

All of the figures in this manual were generated using *Maple*, a computer algebra system. Most of the figures were generated and many of the solutions, especially those involving data, were solved using procedures that were written by Zaven Karian from Denison University. We thank him for providing these. These procedures are available free of charge for your use. They are available on the CD-ROM in the textbook. Short descriptions of these procedures are provided in the “Maple Card” that is on the CD-ROM. Complete descriptions of these procedures are given in *Probability and Statistics: Explorations with MAPLE*, second edition, 1999, written by Zaven Karian and Elliot Tanis, published by Prentice Hall (ISBN 0-13-021536-8).

**REMARK** Note that *Probability and Statistics: Explorations with MAPLE*, second edition, written by Zaven Karian and Elliot Tanis, is available for download from Pearson Education’s online catalog. It has been slightly revised and now contains references to several of the exercises in the 8th edition of *Probability and Statistical Inference*. ♦

Our hope is that this solutions manual will be helpful to each of you in your teaching. If you find an error or wish to make a suggestion, send these to Elliot Tanis at [tanis@hope.edu](mailto:tanis@hope.edu) and he will post corrections on his web page, <http://www.math.hope.edu/tanis/>.

R.V.H.  
E.A.T.



# Chapter 1

## Probability

### 1.1 Basic Concepts

1.1-2 (a)  $S = \{\text{bbb, gbb, bgb, bbg, bgg, gbg, ggb, ggg}\};$

(b)  $S = \{\text{female, male}\};$

(c)  $S = \{000, 001, 002, 003, \dots, 999\}.$

1.1-4 (a) 

|              |   |   |   |    |    |    |    |    |    |    |    |
|--------------|---|---|---|----|----|----|----|----|----|----|----|
| Clutch size: | 4 | 5 | 6 | 7  | 8  | 9  | 10 | 11 | 12 | 13 | 14 |
| Frequency:   | 3 | 5 | 7 | 27 | 26 | 37 | 8  | 2  | 0  | 1  | 1  |




Figure 1.1-4: Clutch sizes for the common gallinule

(c) 9.

**1.1-6 (a)**

|            |    |    |    |   |    |   |    |    |    |    |    |    |    |    |    |
|------------|----|----|----|---|----|---|----|----|----|----|----|----|----|----|----|
| No. Boxes: | 4  | 5  | 6  | 7 | 8  | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 19 | 24 |
| Frequency: | 10 | 19 | 13 | 8 | 13 | 7 | 9  | 5  | 2  | 4  | 4  | 2  | 2  | 1  | 1  |




Figure 1.1-6: Number of boxes of cereal

**1.1-8 (a)**  $f(1) = \frac{2}{10}, f(2) = \frac{3}{10}, f(3) = \frac{3}{10}, f(4) = \frac{2}{10}.$

**1.1-10** This is an experiment.

**1.1-12 (a)**  $50/204 = 0.245; 93/329 = 0.283;$

**(b)**  $124/355 = 0.349; 21/58 = 0.362;$

**(c)**  $174/559 = 0.311; 114/387 = 0.295;$

**(d)** Although James' batting average is higher than Hrbek's on both grass and artificial turf, Hrbek's is higher over all. Note the different numbers of at bats on grass and artificial turf and how this affects the batting averages.

## 1.2 Properties of Probability

**1.2-2** Sketch a figure and fill in the probabilities of each of the disjoint sets.

Let  $A = \{\text{insure more than one car}\}$ ,  $P(A) = 0.85$ .

Let  $B = \{\text{insure a sports car}\}$ ,  $P(B) = 0.23$ .

Let  $C = \{\text{insure exactly one car}\}$ ,  $P(C) = 0.15$ .

It is also given that  $P(A \cap B) = 0.17$ . Since  $P(A \cap C) = 0$ , it follows that

$P(A \cap B \cap C') = 0.17$ . Thus  $P(A' \cap B \cap C') = 0.06$  and  $P(A' \cap B' \cap C) = 0.09$ .

**1.2-4 (a)**  $S = \{\text{HHHH, HHHT, HHTH, HTHH, THHH, HHTT, HTTH, TTHH, HTHT, THTH, THHT, HTTT, THTT, TTHT, TTTH, TTTT}\};$

**(b)** (i)  $5/16$ , (ii)  $0$ , (iii)  $11/16$ , (iv)  $4/16$ , (v)  $4/16$ , (vi)  $9/16$ , (vii)  $4/16$ .

**1.2-6 (a)**  $1/6;$

**(b)**  $P(B) = 1 - P(B') = 1 - P(A) = 5/6;$

**(c)**  $P(A \cup B) = P(S) = 1.$

**1.2-8 (a)**  $P(A \cup B) = 0.4 + 0.5 - 0.3 = 0.6;$

**(b)**

$$\begin{aligned} A &= (A \cap B') \cup (A \cap B) \\ P(A) &= P(A \cap B') + P(A \cap B) \\ 0.4 &= P(A \cap B') + 0.3 \\ P(A \cap B) &= 0.1; \end{aligned}$$

**(c)**  $P(A' \cup B') = P[(A \cap B)'] = 1 - P(A \cap B) = 1 - 0.3 = 0.7.$

**1.2-10** Let  $A = \{\text{lab work done}\}$ ,  $B = \{\text{referral to a specialist}\}$ ,

$$P(A) = 0.41, P(B) = 0.53, P([A \cup B]') = 0.21.$$

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ 0.79 &= 0.41 + 0.53 - P(A \cap B) \\ P(A \cap B) &= 0.41 + 0.53 - 0.79 = 0.15. \end{aligned}$$

**1.2-12**

$$\begin{aligned} A \cup B \cup C &= A \cup (B \cup C) \\ P(A \cup B \cup C) &= P(A) + P(B \cup C) - P[A \cap (B \cup C)] \\ &= P(A) + P(B) + P(C) - P(B \cap C) - P[(A \cap B) \cup (A \cap C)] \\ &= P(A) + P(B) + P(C) - P(B \cap C) - P(A \cap B) - P(A \cap C) \\ &\quad + P(A \cap B \cap C). \end{aligned}$$

**1.2-14 (a)**  $1/3$ ; **(b)**  $2/3$ ; **(c)**  $0$ ; **(d)**  $1/2$ .

**1.2-16 (a)**  $S = \{(1, 2), (1, 3), (1, 4), (1, 5), (2, 3), (2, 4), (2, 5), (3, 4), (3, 5), (4, 5)\};$

**(b) (i)**  $1/10$ ; **(ii)**  $5/10$ .

**1.2-18**  $P(A) = \frac{2[r - r(\sqrt{3}/2)]}{2r} = 1 - \frac{\sqrt{3}}{2}.$

**1.2-20** Note that the respective probabilities are  $p_0$ ,  $p_1 = p_0/4$ ,  $p_2 = p_0/4^2, \dots$

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{p_0}{4^k} &= 1 \\ \frac{p_0}{1 - 1/4} &= 1 \\ p_0 &= \frac{3}{4} \\ 1 - p_0 - p_1 &= 1 - \frac{15}{16} = \frac{1}{16}. \end{aligned}$$

## 1.3 Methods of Enumeration

**1.3-2**  $(4)(3)(2) = 24.$

**1.3-4 (a)**  $(4)(5)(2) = 40$ ; **(b)**  $(2)(2)(2) = 8.$

**1.3-6 (a)**  $4 \binom{6}{3} = 80;$

**(b)**  $4(2^6) = 256;$

**(c)**  $\frac{(4-1+3)!}{(4-1)!3!} = 20.$

**1.3-8**  ${}_9P_4 = \frac{9!}{5!} = 3024.$

**1.3-10**  $S = \{ \text{HHH, HHCH, HCHH, CHHH, HHCCH, HCHCH, CHHCH, HCCHH, CHCHH, CCHHH, CCC, CCHC, CHCC, HCCC, CCHHC, CHCHC, HCCHC, CHHCC, HCHCC, HHCCC} \}$  so there are 20 possibilities.

**1.3-12**  $3 \cdot 3 \cdot 2^{12} = 36,864.$

**1.3-14** 
$$\binom{n-1}{r} + \binom{n-1}{r-1} = \frac{(n-1)!}{r!(n-1-r)!} + \frac{(n-1)!}{(r-1)!(n-r)!}$$

$$= \frac{(n-r)(n-1)! + r(n-1)!}{r!(n-r)!} = \frac{n!}{r!(n-r)!} = \binom{n}{r}.$$

**1.3-16** 
$$0 = (1-1)^n = \sum_{r=0}^n \binom{n}{r} (-1)^r (1)^{n-r} = \sum_{r=0}^n (-1)^r \binom{n}{r}.$$

$$2^n = (1+1)^n = \sum_{r=0}^n \binom{n}{r} (1)^r (1)^{n-r} = \sum_{r=0}^n \binom{n}{r}.$$

**1.3-18** 
$$\binom{n}{n_1, n_2, \dots, n_s} = \binom{n}{n_1} \binom{n-n_1}{n_2} \binom{n-n_1-n_2}{n_3} \dots \binom{n-n_1-\dots-n_{s-1}}{n_s}$$

$$= \frac{n!}{n_1!(n-n_1)!} \cdot \frac{(n-n_1)!}{n_2!(n-n_1-n_2)!}$$

$$\cdot \frac{(n-n_1-n_2)!}{n_3!(n-n_1-n_2-n_3)!} \dots \frac{(n-n_1-n_2-\dots-n_{s-1})!}{n_s!0!}$$

$$= \frac{n!}{n_1!n_2! \dots n_s!}.$$

**1.3-20** (a) 
$$\frac{\binom{19}{3} \binom{52-19}{6}}{\binom{52}{9}} = \frac{102,486}{351,325} = 0.2917;$$

(b) 
$$\frac{\binom{19}{3} \binom{10}{2} \binom{7}{1} \binom{3}{0} \binom{5}{1} \binom{2}{0} \binom{6}{2}}{\binom{52}{9}} = \frac{7,695}{1,236,664} = 0.00622.$$

**1.3-22** 
$$\binom{45}{36} = 886,163,135.$$

## 1.4 Conditional Probability

**1.4-2** (a)  $\frac{1041}{1456};$

(b)  $\frac{392}{633};$

(c)  $\frac{649}{823}.$

(d) The proportion of women who favor a gun law is greater than the proportion of men who favor a gun law.

$$1.4-4 \quad (\text{a}) \quad P(\text{HH}) = \frac{13}{52} \cdot \frac{12}{51} = \frac{1}{17};$$

$$(\text{b}) \quad P(\text{HC}) = \frac{13}{52} \cdot \frac{13}{51} = \frac{13}{204};$$

$$\begin{aligned} (\text{c}) \quad & P(\text{Non-Ace Heart, Ace}) + P(\text{Ace of Hearts, Non-Heart Ace}) \\ &= \frac{12}{52} \cdot \frac{4}{51} + \frac{1}{52} \cdot \frac{3}{51} = \frac{51}{52 \cdot 51} = \frac{1}{52}. \end{aligned}$$

1.4-6 Let  $A = \{3 \text{ or } 4 \text{ kings}\}$ ,  $B = \{2, 3, \text{ or } 4 \text{ kings}\}$ .

$$\begin{aligned} P(A|B) &= \frac{P(A \cap B)}{P(B)} = \frac{N(A)}{N(B)} \\ &= \frac{\binom{4}{3}\binom{48}{10} + \binom{4}{4}\binom{48}{9}}{\binom{4}{2}\binom{48}{11} + \binom{4}{3}\binom{48}{10} + \binom{4}{4}\binom{48}{9}} = 0.170. \end{aligned}$$

1.4-8 Let  $H = \{\text{died from heart disease}\}$ ;  $P = \{\text{at least one parent had heart disease}\}$ .

$$P(H|P) = \frac{N(H \cap P)}{N(P)} = \frac{110}{648}.$$

$$1.4-10 \quad (\text{a}) \quad \frac{3}{20} \cdot \frac{2}{19} \cdot \frac{1}{18} = \frac{1}{1140};$$

$$(\text{b}) \quad \frac{\binom{3}{2}\binom{17}{1}}{\binom{20}{3}} \cdot \frac{1}{17} = \frac{1}{760};$$

$$(\text{c}) \quad \sum_{k=1}^9 \frac{\binom{3}{2}\binom{17}{2k-2}}{\binom{20}{2k}} \cdot \frac{1}{20-2k} = \frac{35}{76} = 0.4605.$$

(d) Draw second. The probability of winning in  $1 - 0.4605 = 0.5395$ .

$$1.4-12 \quad \frac{\binom{2}{0}\binom{8}{5}}{\binom{10}{5}} \cdot \frac{2}{5} + \frac{\binom{2}{1}\binom{8}{4}}{\binom{10}{5}} \cdot \frac{1}{5} = \frac{1}{5}.$$

$$1.4-14 \quad (\text{a}) \quad P(A) = \frac{52}{52} \cdot \frac{51}{52} \cdot \frac{50}{52} \cdot \frac{49}{52} \cdot \frac{48}{52} \cdot \frac{47}{52} = \frac{8,808,975}{11,881,376} = 0.74141;$$

$$(\text{b}) \quad P(A') = 1 - P(A) = 0.25859.$$

$$1.4-16 \quad (\text{a}) \quad \text{It doesn't matter because } P(B_1) = \frac{1}{18}, \quad P(B_5) = \frac{1}{18}, \quad P(B_{18}) = \frac{1}{18};$$

$$(\text{b}) \quad P(B) = \frac{2}{18} = \frac{1}{9} \quad \text{on each draw.}$$

$$1.4-18 \quad \frac{3}{5} \cdot \frac{5}{8} + \frac{2}{5} \cdot \frac{4}{8} = \frac{23}{40}.$$

- 1.4-20** (a)  $P(A_1) = 30/100$ ;  
 (b)  $P(A_3 \cap B_2) = 9/100$ ;  
 (c)  $P(A_2 \cup B_3) = 41/100 + 28/100 - 9/100 = 60/100$ ;  
 (d)  $P(A_1 | B_2) = 11/41$ ;  
 (e)  $P(B_1 | A_3) = 13/29$ .

## 1.5 Independent Events

**1.5-2** (a)  $P(A \cap B) = P(A)P(B) = (0.3)(0.6) = 0.18$ ;  
 $P(A \cup B) = P(A) + P(B) - P(A \cap B)$   
 $= 0.3 + 0.6 - 0.18$   
 $= 0.72$ .

(b)  $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0}{0.6} = 0$ .

**1.5-4** Proof of (b):  $P(A' \cap B) = P(B)P(A'|B)$   
 $= P(B)[1 - P(A|B)]$   
 $= P(B)[1 - P(A)]$   
 $= P(B)P(A')$ .

Proof of (c):  $P(A' \cap B') = P[(A \cup B)']$   
 $= 1 - P(A \cup B)$   
 $= 1 - P(A) - P(B) + P(A \cap B)$   
 $= 1 - P(A) - P(B) + P(A)P(B)$   
 $= [1 - P(A)][1 - P(B)]$   
 $= P(A')P(B')$ .

**1.5-6**  $P[A \cap (B \cap C)] = P[A \cap B \cap C]$   
 $= P(A)P(B)P(C)$   
 $= P(A)P(B \cap C)$ .

$$\begin{aligned} P[A \cap (B \cup C)] &= P[(A \cap B) \cup (A \cap C)] \\ &= P(A \cap B) + P(A \cap C) - P(A \cap B \cap C) \\ &= P(A)P(B) + P(A)P(C) - P(A)P(B)P(C) \\ &= P(A)[P(B) + P(C) - P(B \cap C)] \\ &= P(A)P(B \cup C). \end{aligned}$$

$$\begin{aligned} P[A' \cap (B \cap C')] &= P(A' \cap C' \cap B) \\ &= P(B)[P(A' \cap C') | B] \\ &= P(B)[1 - P(A \cup C | B)] \\ &= P(B)[1 - P(A \cup C)] \\ &= P(B)P[(A \cup C)'] \\ &= P(B)P(A' \cap C') \\ &= P(B)P(A')P(C') \\ &= P(A')P(B)P(C') \\ &= P(A')P(B \cap C'). \end{aligned}$$

$$\begin{aligned} P[A' \cap B' \cap C'] &= P[(A \cup B \cup C)'] \\ &= 1 - P(A \cup B \cup C) \\ &= 1 - P(A) - P(B) - P(C) + P(A)P(B) + P(A)P(C) + \\ &\quad P(B)P(C) - P(A)P(B)P(C) \\ &= [1 - P(A)][1 - P(B)][1 - P(C)] \\ &= P(A')P(B')P(C'). \end{aligned}$$

$$1.5-8 \quad \frac{1}{6} \cdot \frac{2}{6} \cdot \frac{3}{6} + \frac{1}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} + \frac{5}{6} \cdot \frac{2}{6} \cdot \frac{3}{6} = \frac{2}{9}.$$

$$1.5-10 \quad (\text{a}) \quad \frac{3}{4} \cdot \frac{3}{4} = \frac{9}{16};$$

$$(\text{b}) \quad \frac{1}{4} \cdot \frac{3}{4} + \frac{3}{4} \cdot \frac{2}{4} = \frac{9}{16};$$

$$(\text{c}) \quad \frac{2}{4} \cdot \frac{1}{4} + \frac{2}{4} \cdot \frac{4}{4} = \frac{10}{16}.$$

$$1.5-12 \quad (\text{a}) \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2;$$

$$(\text{b}) \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2;$$

$$(\text{c}) \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2;$$

$$(\text{d}) \quad \frac{5!}{3!2!} \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2.$$

$$1.5-14 \quad (\text{a}) \quad 1 - (0.4)^3 = 1 - 0.064 = 0.936;$$

$$(\text{b}) \quad 1 - (0.4)^8 = 1 - 0.00065536 = 0.99934464.$$

$$1.5-16 \quad (\text{a}) \quad \sum_{k=0}^{\infty} \frac{1}{5} \left(\frac{4}{5}\right)^{2k} = \frac{5}{9};$$

$$(\text{b}) \quad \frac{1}{5} + \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{1}{3} + \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{2}{3} \cdot \frac{1}{2} \cdot \frac{1}{1} = \frac{3}{5}.$$

$$1.5-18 \quad (\text{a}) \quad 7; (\text{b}) \quad (1/2)^7; (\text{c}) \quad 63; (\text{d}) \quad \text{No! } (1/2)^{63} = 1/9,223,372,036,854,775,808.$$

|        |     |        |        |        |        |        |
|--------|-----|--------|--------|--------|--------|--------|
| 1.5-20 | n   | 3      | 6      | 9      | 12     | 15     |
|        | (a) | 0.7037 | 0.6651 | 0.6536 | 0.6480 | 0.6447 |
|        | (b) | 0.6667 | 0.6319 | 0.6321 | 0.6321 | 0.6321 |

(c) Very little when  $n > 15$ , sampling with replacement

Very little when  $n > 10$ , sampling without replacement.

(d) Convergence is faster when sampling with replacement.

## 1.6 Bayes's Theorem

$$1.6-2 \quad (\text{a}) \quad \begin{aligned} P(G) &= P(A \cap G) + P(B \cap G) \\ &= P(A)P(G|A) + P(B)P(G|B) \\ &= (0.40)(0.85) + (0.60)(0.75) = 0.79; \end{aligned}$$

$$\begin{aligned} (\text{b}) \quad P(A|G) &= \frac{P(A \cap G)}{P(G)} \\ &= \frac{(0.40)(0.85)}{0.79} = 0.43. \end{aligned}$$

**1.6-4** Let event  $B$  denote an accident and let  $A_1$  be the event that age of the driver is 16–25. Then

$$\begin{aligned} P(A_1 | B) &= \frac{(0.1)(0.05)}{(0.1)(0.05) + (0.55)(0.02) + (0.20)(0.03) + (0.15)(0.04)} \\ &= \frac{50}{50 + 110 + 60 + 60} = \frac{50}{280} = 0.179. \end{aligned}$$

**1.6-6** Let  $B$  be the event that the policyholder dies. Let  $A_1, A_2, A_3$  be the events that the deceased is standard, preferred and ultra-preferred, respectively. Then

$$\begin{aligned} P(A_1 | B) &= \frac{(0.60)(0.01)}{(0.60)(0.01) + (0.30)(0.008) + (0.10)(0.007)} \\ &= \frac{60}{60 + 24 + 7} = \frac{60}{91} = 0.659; \\ P(A_2 | B) &= \frac{24}{91} = 0.264; \\ P(A_3 | B) &= \frac{7}{91} = 0.077. \end{aligned}$$

**1.6-8** Let  $A$  be the event that the DVD player is under warranty.

$$\begin{aligned} P(B_1 | A) &= \frac{(0.40)(0.10)}{(0.40)(0.10) + (0.30)(0.05) + (0.20)(0.03) + (0.10)(0.02)} \\ &= \frac{40}{40 + 15 + 6 + 2} = \frac{40}{63} = 0.635; \\ P(B_2 | A) &= \frac{15}{63} = 0.238; \\ P(B_3 | A) &= \frac{6}{63} = 0.095; \\ P(B_4 | A) &= \frac{2}{63} = 0.032. \end{aligned}$$

**1.6-10 (a)**  $P(AD) = (0.02)(0.92) + (0.98)(0.05) = 0.0184 + 0.0490 = 0.0674$ ;

**(b)**  $P(N | AD) = \frac{0.0490}{0.0674} = 0.727$ ;  $P(A | AD) = \frac{0.0184}{0.0674} = 0.273$ ;

**(c)**  $P(N | ND) = \frac{(0.98)(0.95)}{(0.02)(0.08) + (0.98)(0.95)} = \frac{9310}{16 + 9310} = 0.998$ ;  $P(A | ND) = 0.002$ .

**(d)** Yes, particularly those in part (b).

**1.6-12** Let  $D = \{\text{has the disease}\}$ ,  $DP = \{\text{detects presence of disease}\}$ . Then

$$\begin{aligned} P(D | DP) &= \frac{P(D \cap DP)}{P(DP)} \\ &= \frac{P(D) \cdot P(DP | D)}{P(D) \cdot P(DP | D) + P(D') \cdot P(DP | D')} \\ &= \frac{(0.005)(0.90)}{(0.005)(0.90) + (0.995)(0.02)} \\ &= \frac{0.0045}{0.0045 + 0.199} = \frac{0.0045}{0.2035} = 0.0221. \end{aligned}$$

**1.6-14** Let  $D = \{\text{defective roll}\}$  Then

$$\begin{aligned} P(I|D) &= \frac{P(I \cap D)}{P(D)} \\ &= \frac{P(I) \cdot P(D|I)}{P(I) \cdot P(D|I) + P(II) \cdot P(D|II)} \\ &= \frac{(0.60)(0.03)}{(0.60)(0.03) + (0.40)(0.01)} \\ &= \frac{0.018}{0.018 + 0.004} = \frac{0.018}{0.022} = 0.818. \end{aligned}$$



## Chapter 2

# Discrete Distributions

### 2.1 Random Variables of the Discrete Type

2.1-2 (a)

$$f(x) = \begin{cases} 0.6, & x = 1, \\ 0.3, & x = 5, \\ 0.1, & x = 10, \end{cases}$$

(b)




Figure 2.1-2: A probability histogram

2.1-4 (a)  $f(x) = \frac{1}{10}$ ,  $x = 0, 1, 2, \dots, 10$ ;

(b)  $\mathcal{N}(\{0\})/150 = 11/150 = 0.073$ ;  $\mathcal{N}(\{5\})/150 = 13/150 = 0.087$ ;  
 $\mathcal{N}(\{1\})/150 = 14/150 = 0.093$ ;  $\mathcal{N}(\{6\})/150 = 22/150 = 0.147$ ;  
 $\mathcal{N}(\{2\})/150 = 13/150 = 0.087$ ;  $\mathcal{N}(\{7\})/150 = 16/150 = 0.107$ ;  
 $\mathcal{N}(\{3\})/150 = 12/150 = 0.080$ ;  $\mathcal{N}(\{8\})/150 = 18/150 = 0.120$ ;  
 $\mathcal{N}(\{4\})/150 = 16/150 = 0.107$ ;  $\mathcal{N}(\{9\})/150 = 15/150 = 0.100$ .

(c)




Figure 2.1-4: Michigan daily lottery digits

**2.1-6 (a)**  $f(x) = \frac{6 - |7 - x|}{36}$ ,  $x = 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12$ .

(b)




Figure 2.1-6: Probability histogram for the sum of a pair of dice

**2.1-8 (a)** The space of  $W$  is  $S = \{0, 1, 2, 3, 4, 5, 6, 7\}$ .

$$P(W = 0) = P(X = 0, Y = 0) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8}, \text{ assuming independence.}$$

$$P(W = 1) = P(X = 0, Y = 1) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 2) = P(X = 2, Y = 0) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 3) = P(X = 2, Y = 1) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 4) = P(X = 0, Y = 4) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 5) = P(X = 0, Y = 5) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 6) = P(X = 2, Y = 4) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 7) = P(X = 2, Y = 5) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8}.$$

$$\text{That is, } f(w) = P(W = w) = \frac{1}{8}, \quad w \in S.$$

(b)




Figure 2.1-8: Probability histogram of sum of two special dice

**2.1-10 (a)** 
$$\frac{\binom{3}{1}\binom{47}{9}}{\binom{50}{10}} = \frac{39}{98};$$

**(b)** 
$$\sum_{x=0}^1 \frac{\binom{3}{x}\binom{47}{10-x}}{\binom{50}{10}} = \frac{221}{245}.$$

$$\mathbf{2.1-12} \quad OC(0.04) = \frac{\binom{1}{0}\binom{24}{5}}{\binom{25}{5}} + \frac{\binom{1}{1}\binom{24}{4}}{\binom{25}{5}} = 1.000;$$

$$OC(0.08) = \frac{\binom{2}{0}\binom{23}{5}}{\binom{25}{5}} + \frac{\binom{2}{1}\binom{23}{4}}{\binom{25}{5}} = 0.967;$$

$$OC(0.12) = \frac{\binom{3}{0}\binom{22}{5}}{\binom{25}{5}} + \frac{\binom{3}{1}\binom{22}{4}}{\binom{25}{5}} = 0.909;$$

$$OC(0.16) = \frac{\binom{4}{0}\binom{21}{5}}{\binom{25}{5}} + \frac{\binom{4}{1}\binom{21}{4}}{\binom{25}{5}} = 0.834.$$

$$\mathbf{2.1-14} \quad P(X \geq 1) = 1 - P(X = 0) = 1 - \frac{\binom{3}{0}\binom{17}{5}}{\binom{20}{5}} = 1 - \frac{91}{228} = \frac{137}{228} = 0.60.$$

**2.1-16 (a)** Let  $Y$  equal the number of  $H$  chips that are selected. Then

$X = |Y - (10 - Y)| = |2Y - 10|$  and the p.m.f. of  $Y$  is

$$g(y) = \frac{\binom{10}{y}\binom{10}{10-y}}{\binom{20}{10}}, \quad y = 0, 1, \dots, 10.$$

The p.m.f. of  $X$  is as follows:

|                     |                       |                         |                         |                       |                    |
|---------------------|-----------------------|-------------------------|-------------------------|-----------------------|--------------------|
| $f(0) = g(5)$       | $f(2) = 2g(6)$        | $f(4) = 2g(7)$          | $f(6) = 2g(8)$          | $f(8) = 2g(9)$        | $f(10) = 2g(10)$   |
| $\frac{1}{184,756}$ | $\frac{2025}{92,378}$ | $\frac{22,050}{46,189}$ | $\frac{22,050}{46,189}$ | $\frac{2025}{92,378}$ | $\frac{1}{92,378}$ |

**(b)** The mode is equal to 2.

**2.1-18 (a)**  $P(2, 1, 6, 10)$  means that 2 is in position 1 so 1 cannot be selected. Thus

$$P(2, 1, 6, 10) = \frac{\binom{1}{0}\binom{1}{1}\binom{8}{5}}{\binom{10}{6}} = \frac{56}{210} = \frac{4}{15};$$

$$\mathbf{(b)} \quad P(i, r, k, n) = \frac{\binom{i-1}{r-1}\binom{1}{1}\binom{n-i}{k-r}}{\binom{n}{k}}.$$