Epidemiology of Women's Health Senie TestBank – Chapter 1

Multiple Choice

1.	What is the primary goal of epidemiological research?
	 A. to eliminate all disease B. to identify new diseases C. to identify risk factors associated with diseases D. to gauge effectiveness of treatment options
	Ans. C See page 8.
2.	Women have the greatest risk of developing cancer in which part of the body?
	A. pancreasB. lungC. colonD. breast
	Ans. D See page 12.
3.	Treatment during pregnancy with has been associated with birth defects.
	A. thalidomideB. theophyllineC. acetaminophenD. diphenhydramine
	Ans. A See page 9.
4.	Chronic conditions are often characterized by requiring medical treatment.
	A. musculoskeletal painB. persistent nauseaC. hypotension

	D. tinnitus	
	Ans. A See page 12.	
5.	Which of the following conditions has had the highest mortality rate for the last 50 years?	
	A. cancerB. heart diseaseC. strokeD. diabetes	
	Ans. B See page 11.	
6.	What is considered essential for maintenance of an alert mind in older women?	
	A. sufficient hydrationB. moderate exerciseC. social interactionD. supplemental nutrition	
	Ans. C See page 13.	
True or False		
7.	The obesity epidemic is due in part to the availability of an overabundance of fresh and processed foods.	
	A. True B. False	
	Ans. True See page 7.	

8.	With more women surviving to older ages, lower rates of chronic conditions are anticipated.
	A. True B. False
	Ans. False See page 8.
9.	Despite 30 years of research and new treatments, human immunodeficiency virus (HIV) infection is still inevitably fatal.
	A. True B. False
	Ans. False See page 9.
10.	Suicide is more frequent among women with bipolar disease than among women with other psychologic conditions.
	A. True B. False
	Ans. True See page 10.
11.	Autoimmune diseases are a major cause of infertility and pregnancy loss.
	A. True B. False
	Ans. True See page 10.
12.	Environmental exposures can influence an individual's risk of disease and age at onset.

A. True

B. False

Ans. True See page 13.

Short Answer

13. What public health measures have successfully reduced the sale of cigarettes and diminished interest in smoking?

Ans. These public health measures include controls on cigarette advertising, increased taxes making cigarettes less affordable, and laws prohibiting smoking in most public indoor and outdoor settings, including parks and beaches.

See page 6.

14. List three of the goals defined in the office of research women's health strategic plan for 2020.

Ans. Responses should include any three of the following:

- Increase the study of sex and gender differences in basic biomedical and behavioral research.
- Incorporate findings of sex and gender differences into the design and application of new technologies, medical devices, and therapeutic drugs.
- Actualize personalized prevention, diagnostics, and therapeutics for women and girls.
- Create strategic alliances and partnerships to maximize the domestic and global impact of women's health research.
- Develop and implement new communication and social networking technologies to increase understanding and appreciation of women's health and wellness research.
- Employ innovative strategies to build a well-trained, diverse, and vigorous women's health research workforce.

See page 4.

15. Explain why life expectancy has significantly increased since World War II.

Epidemiology of Women's Health Senie TestBank – Chapter 1

Ans. Life expectancy has significantly increased since World War II in large part because of medical innovations and drug discoveries such as antibiotics, vaccines, and enhanced clinical devices and procedures.

See page 5.