5) During interphase of the cell cycle, \_\_\_\_\_.

C) sister chromatids move to opposite poles

A) the nuclear membrane disappears

D) DNA content essentially doubles

B) RNA replicates

E) DNA recombines


Answer: D

6) If a typical somatic cell has 64 chromosomes, how many chromosomes are expected in each					6)
gamete of that A) 128 Answer: D	organism? B) 16	C) 8	D) 32	E) 64	
	7) In an organism with 52 chromosomes, how many bivalents would be expected to form during				
meiosis? A) 26 Answer: A	B) 13	C) 52	D) 208	E) 104	
8) The ant <i>Myrmecia pilosula</i> is found in Australia and is named <i>bulldog</i> because of its aggressive behavior. It is particularly interesting because it carries all its genetic information in a single pair of chromosomes. In other words, $2n = 2$ . (Males are haploid and have just one chromosome.) Which of the following figures would most likely represent a correct configuration of chromosomes in a metaphase I cell of a female?					8)
A.	B.	C.	D.	E.	
A) A Answer: A	B) B	C) C	D) D	E) E	
pilosula contain cell of a female A) 32 picogra B) 16 picogra C) 8 picogram D) 4 picogram	is 2 picograms of DN? ? ms ms	JA. How much D	omatic cell nucleus in NA would be expected he question.	•	9)
10) An interesting group of ants consists of several virtually identical, closely related species, with females having chromosome numbers of 18, 20, 32, 48, 60, 62, and 64. If one crossed a female of species (A) with 32 chromosomes and a male species (B) with 9 chromosomes (males are haploid, and each gamete contains the <i>n</i> complement), how many chromosomes would one expect in the body (somatic) cells of the female offspring?				10)	
A) 25 Answer: A	B) 4.5	C) 41	D) 9	E) 32	

		11)
·····		
Answer: E		
2) In a healthy male, how many sperm cells would be expected to form from (a) 400 prima	ary	12)
spermatocytes? (b) 400 secondary spermatocytes?		
A) (a) 1600; (b) 1600		
B) (a) 1600; (b) 800		
C) (a) 400; (b) 400		
D) (a) 800; (b) 800		
E) (a) 400; (b) 800		
Answer: B		
3) In a healthy female, how many secondary occytes would be expected to form from 100		13)
	<b>J</b>	
	,	
ΓANSWER. Write the word or phrase that best completes each statement or answers th	ne question	
4) Which type of cell structure is the nucleolus organizer (NOR) responsible for	14)	
7 (113 VV CI : 11 DO SOTTIC		
5) Name two cellular organelles, each having genetic material, that are involved in either	15)	
photosynthesis or respiration.		
Answer: chloroplasts and mitochondria		
// List form towns and to decomine the marked promphalogical with many set to amb matic of	1/\	
·	16)	
Answer: metacentric, submetacentric, acrocentric, telocentric		
7) Homologous chromosomes can be matched by their similar structure and function	17)	
within a nucleus. Which chromosomes making up a genome do not follow the same	·	
characteristics of homology?		
Answer: sex-determining chromosomes		
	18)	
Answer: 2; 2		
9) How many haploid sets of chromosomes are present in an individual cell that is	19)	
tetraploid (4 <i>n</i> )?	,	
Answer: 4		
	spermatocytes? (b) 400 secondary spermatocytes? A) (a) 1600: (b) 1600 B) (a) 1600: (b) 800 C) (a) 400: (b) 800 E) (a) 400: (b) 800 E) (a) 400: (b) 800 Answer: B  3) In a healthy female, how many secondary oocytes would be expected to form from 100 primary oocytes? How many first polar bodies would be expected from 100 primary ocytes? How many first polar bodies would be expected from 100 primary ocytes? How many first polar bodies would be expected from 100 primary ocytes? How many first polar bodies would be expected from 100 primary ocytes? How many first polar bodies would be expected from 100 primary ocytes? Ocytes and the expected from 100 primary ocytes? Ocytes and the expected from 100 primary ocytes? Ocytes and the expected from 100 primary ocytes. Ocytes and the expected from 100 primary ocytes and the expected from 100 primary ocytes. Ocytes and 100 primary ocytes.	A) dyad formation B) monad movement to opposite poles C) chiasma segregation D) side-by-side alignment of nonhomologous chromosomes E) side-by-side alignment of homologous chromosomes Answer: E  2) In a healthy male, how many sperm cells would be expected to form from (a) 400 primary spermatocytes? A) (a) 1600: (b) 1600 B) (a) 1600: (b) 1600 C) (a) 400: (b) 800 C) (a) 400: (b) 800 E) (a) 400: (b) 800 Answer: B  3) In a healthy female, how many secondary oocytes would be expected to form from 100 primary oocytes? A) 200: 200 B) 50: 50 C) 100: 100 D) 100: 50 E) 200: 50 Answer: C  FANSWER. Write the word or phrase that best completes each statement or answers the question 4) Which type of cell structure is the nucleolus organizer (NOR) responsible for producing? Answer: ribosome  5) Name two cellular organelles, each having genetic material, that are involved in either photosynthesis or respiration. Answer: chloroplasts and mitochondria 6) List four terms used to describe the normal morphologies, with respect to arm ratio, of eukaryotic chromosomes. Answer: metacentric, submetacentric, acrocentric, telocentric 7) Homologous chromosomes can be matched by their similar structure and function within a nucleus. Which chromosomes making up a genome do not follow the same characteristics of homology? Answer: ex-determining chromosomes 8) How many haploid sets of chromosomes are present in a diploid individual cell with a chromosome number of 467 32? Answer: 2; 2  9) How many haploid sets of chromosomes are present in an individual cell that is 19) tetraploid (4n)?

20)	In which stage of the cell cycle is G0 located?  Answer: G1	20)
21)	When cells withdraw from the continuous cell cycle and enter a "quiescent" phase, which stage are they said to be in? Answer: G0	21)
22)	The house fly, <i>Musca domestica</i> , has a haploid chromosome number of 6. How many chromatids should be present in a diploid, somatic, metaphase cell?  Answer: 24	22)
23)	Regarding the mitotic cell cycle, what is meant by a checkpoint?  Answer: A checkpoint is the portion of a cell cycle that is sensitive to a variety of conditions that impact the eventual health of the cell or individual. It serves as a potential halting point to facilitate DNA repair.	23)
24)	After which meiotic stage (meiosis I or II) would one expect monads to be formed?  Answer: meiosis II	24)
25)	Assume that the somatic cells of a male contain one pair of homologous chromosomes (e.g., AaAb) and an additional chromosome without a homolog (e.g., W). Which chromosomal combinations would be expected in the meiotic products (spermatids) of a single primary spermatocyte? (There may be more than one answer.)  Answer: AaW, AaW, Ab, Ab or Aa, Aa, AbW, AbW	25)
26)	The accompanying sketch depicts a cell from an organism in which $2n = 2$ and each chromosome is metacentric.	26)

(a) Circle the correct stage for the cell in this sketch:



anaphase of mitosis anaphase of meiosis I anaphase of meiosis II telophase of mitosis

(b) Given that each G1 nucleus from this organism contains 16 picograms of DNA, how many picograms of chromosomal DNA would you expect in the cell shown here?

Answer: (a) anaphase of meiosis II

(b) 16


27)	You may have heard through various media of an animal alleged to be the hybrid of a	27)
	rabbit and a cat. Given that the cat ( <i>Felis domesticus</i> ) has a diploid chromosome number of 38 and a rabbit ( <i>Oryctolagus cuniculus</i> ) has a diploid chromosome number of 44, what would be the expected chromosome number in the somatic tissues of this alleged hybrid?	
	Answer: 41	
28)	The horse ( <i>Equus caballus</i> ) has 32 pairs of chromosomes, whereas the donkey ( <i>Equus asinus</i> ) has 31 pairs of chromosomes. How many chromosomes would be expected in	28)
	the somatic tissue of a mule? Answer: 63	
29)	Name two evolutionarily significant benefits of meiosis that are not present in mitosis.  Answer: reshuffling of homologous chromosomes and crossing over	29)
30)	What is meant by the term <i>chiasma</i> ?	30)
ŕ	Answer: Chiasma are areas where chromatids intertwine during meiosis and where nonsister chromatids have undergone genetic exchange.	
31)	List in order of occurrence the phases of mitosis.	31)
	Answer: prophase, prometaphase, metaphase, anaphase, telophase	
32)	The two terms <i>reductional</i> and <i>equational</i> generally refer to which stages of meiosis (I or II)?	32)
	Answer: meiosis I and meiosis II, respectively	
33)	Trisomy 21 or Down syndrome occurs when there is a normal diploid chromosomal complement of 46 chromosomes plus one (extra) chromosome 21. Such individuals therefore have 47 chromosomes. Assume that a mating occurs between a female with Down syndrome and a normal 46-chromosome male. What proportion of the offspring would be expected to have Down syndrome? Justify your answer.	33)
	Answer: One-half of the offspring would be expected to have Down syndrome because of 2 $\times$ 1 segregation of chromosome 21 at anaphase I.	
34)	Normal diploid somatic (body) cells of the mosquito Culex pipiens contain six	34)
	chromosomes. Assign the symbols AmAp, BmBp, and CmCp to the three homologous chromosomal pairs. The "m" superscript indicates that the homolog is maternally derived; the "p" indicates a paternally derived homolog. Assume that in the genus	
	Culex, the sex chromosomes are morphologically identical.	
	(a) For each of the cell types given below, draw and label (with reference to the symbols defined above) an expected chromosomal configuration.	
	mitotic metaphase metaphase of meiosis I metaphase of meiosis II	
	(b) The stage at which "sister chromatids go to opposite poles" immediately follows which of the stages listed in (a)?	

(c) Assuming that all nuclear DNA is restricted to chromosomes and that the amount of nuclear DNA essentially doubles during the S phase of interphase, how much nuclear DNA would be present in each cell listed in part (a)? *Note*: Assume that the G1 nucleus of a mosquito cell contains  $3.0 \times 10^{-12}$  grams of DNA.

34)


(d) Given that the sex of *Culex* is determined by alleles of one gene-males heterozygous, *Mm*, and females homozygous, *mm*-illustrate a labeled chromosomal configuration (involving the symbols AmAp, BmBp, and CmCp and the *M* locus) in a primary spermatocyte at metaphase. Assume that the *M* locus is on the AmAp chromosome and that crossing over has not occurred between the *M* locus and the centromere.

Answer:



(b) metaphase of meiosis II and mitotic metaphase


(c) 6, 6, 3



- 35) *Drosophila melanogaster*, the fruit fly, has a 2*n* chromosome number of 8. Assume that you are microscopically examining the mitotic and meiotic cells of this organism. You note that in the female, two chromosomal pairs are metacentric and two pairs are acrocentric.
  - (a) Draw the chromosomal configurations as you would expect to see them at the stages listed:

35)


- (b) Given that the previously mentioned cells are from individuals heterozygous for two independently segregating, autosomal loci, *plum eyes* and *curled wings*, place appropriate symbols (of your designation) on chromosomes in the drawings you made for part (a). Assume no crossing over, and there may be more than one correct answer in some cases.
- (c) Assuming that a somatic G2 nucleus from the individuals mentioned above contains about 8.0 picograms of DNA, how much nuclear DNA would you expect in each of the cells mentioned in part (a)?



36) Down syndrome, or trisomy 21, in humans is caused by an extra copy of the relatively small, acrocentric 21. Including only chromosome 21, the X chromosome (medium in size and somewhat metacentric) and the Y chromosome (small and acrocentric), draw one possible array of chromosomes in the four sperm cells produced by the complete meiosis of one primary spermatocyte. For the purposes of this question, assume that males with Down syndrome produce normal ratios of sperm cells. (More than one answer is possible.)


36)

Answer:



37) Assume that an organism has a diploid chromosome number of 6. Two chromosomal pairs are telocentric, and the other pair is metacentric. Assume that the sex chromosomes are morphologically identical. Draw chromosomes as you would expect them to appear at the following stages: primary oocyte (metaphase) secondary spermatocyte (metaphase) first polar body (metaphase)





38) There is about as much nuclear DNA in a primary spermatocyte as in \_\_\_\_\_\_ (how many) spermatids?
Answer: 4
39) List, in order of appearance, all the cell types expected to be formed during (a)
39)

spermatogenesis and (b) oogenesis.

Answer: (a) spermatogonia, primary spermatocyte, secondary spermatocyte, spermatozoa

39)

spermatozoa
(b) oogonium, primary oocyte, secondary oocyte and first polar body, ootid, ovum, and second polar body

TRUE/FALSE. Write 'T' if the statement is true and 'F' if the statement is false.

40) True or False: If a typical G1 nucleus is 2*n* and contains 2C (two complements) of DNA, a 40) \_\_\_\_\_ prophase I cell is 2*n* and contains 4C of DNA.

Answer: • True False

41) True or False: S phase is the part of interphase when DNA duplication takes place.

41) \_\_\_\_\_

Answer: • True False

42)	True or False: The cei	ntromere of a chromosome separates during anaphase.	42)
	Answer: • True	False	
43)	True or False: A chro mitotic or meiotic cel	mosome may contain one or two chromatids in different phases of the I cycle.	43)
	Answer: • True	False	
44)	True or False: If a typhaploid (n) contains	oical G1 nucleus contains 2C (two complements) of DNA, a gamete that is 1C of DNA.	44)
	Answer: • True	False	
45)		meiosis, chromosome number reduction takes place in anaphase II. False	45)
46)		lent at pachytene contains four chromatids. False	46)
47)	True or False: The me	eiotic cell cycle involves two cell divisions but only one DNA replication. False	47)
48)	chromosomes at the		48)
	Answer: • True	False	
49)	•	panism with a diploid chromosome number of 46 will produce 223 mosomes at the end of meiosis.	49)
	Answer: • True	False	