

/test-bank-physics-for-scientists-and-engineers-foundations-and-connections-1e-katz
Chapter 1 - Getting Started.

1. Which of the following products of ratios gives the conversion factor to convert miles per hour $\left(\frac{\text{mi}}{\text{h}}\right)$ to meters per second $\left(\frac{\text{m}}{\text{s}}\right)$?

- $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{1 \text{ in.}}{2.54 \text{ cm}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{1 \text{ h}}{3600 \text{ s}}$
- $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in.}} \cdot \frac{100 \text{ cm}}{1 \text{ m}} \cdot \frac{1 \text{ h}}{3600 \text{ s}}$
- $\frac{1 \text{ mi}}{5280 \text{ ft}} \cdot \frac{1 \text{ ft}}{12 \text{ in.}} \cdot \frac{1 \text{ in.}}{2.54 \text{ cm}} \cdot \frac{100 \text{ cm}}{1 \text{ m}} \cdot \frac{3600 \text{ s}}{1 \text{ h}}$
- $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in.}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{1 \text{ h}}{3600 \text{ s}}$
- $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in.}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{3600 \text{ s}}{1 \text{ h}}$

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

2. The density of an object is defined as

- the volume occupied by each unit of mass.
- the amount of mass for each unit of volume.
- the weight of each unit of volume.
- the amount of the substance that has unit volume and unit mass.
- the amount of the substance that contains as many particles as 12 grams of the carbon-12 isotope.

ANSWER:

b

POINTS:

1

DIFFICULTY:

Easy

3. If you drove day and night without stopping for one year without exceeding the legal highway speed limit in the United States, the maximum number of miles you could drive would be closest to

- 8700.
- 300,000.
- 500,000.
- 1,000,000.
- 32,000,000.

ANSWER:

c

POINTS:

2

DIFFICULTY:

Average

4. The term $\frac{1}{2} \rho v^2$ occurs in Bernoulli's equation in Chapter 15, with ρ being the density of a fluid and v its speed. The dimensions of this term are

Chapter 1 - Getting Started.

- a. $M^{-1}L^5T^2$.
- b. MLT^2 .
- c. $ML^{-1}T^{-2}$.
- d. $M^{-1}L^9T^{-2}$.
- e. $M^{-1}L^3T^{-2}$.

ANSWER:

c

POINTS:

2

DIFFICULTY:

Average

5. Which of the following quantities has the same dimensions as kinetic energy, $\frac{1}{2}mv^2$?
 Note: $[[a]] = [[g]] = LT^{-2}$; $[[h]] = L$ and $[[v]] = LT^{-1}$.

- a. ma
- b. mvx
- c. mvt
- d. mgh
- e. mgt

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

6. The quantity with the same units as force times time, Ft , with dimensions MLT^{-1} is

- a. mv
- b. mvr
- c. mv^2r
- d. ma
- e. $\frac{mv^2}{r}$

ANSWER:

a

POINTS:

2

DIFFICULTY:

Average

7. The equation for the change of position of a train starting at $x = 0$ m is given by $x = \frac{1}{2}at^2 + bt^3$. The dimensions of b are

- a. T^{-3}
- b. LT^{-3}
- c. LT^{-2}
- d. LT^{-1}
- e. $L^{-1}T^{-1}$

Chapter 1 - Getting Started.

ANSWER:

b

POINTS:

2

DIFFICULTY:

Average

8. One mole of the carbon-12 isotope contains 6.022×10^{23} atoms. What volume in m^3 would be needed to store one mole of cube-shaped children's blocks 2.00 cm long on each side?

- a. 4.8×10^{18}
- b. 1.2×10^{22}
- c. 6.0×10^{23}
- d. 1.2×10^{24}
- e. 4.8×10^{24}

ANSWER:

a

POINTS:

2

DIFFICULTY:

Average

9. Which of the following products of ratios gives the conversion factors to convert meters per second $\left(\frac{\text{m}}{\text{s}}\right)$ to miles per hour $\left(\frac{\text{mi}}{\text{h}}\right)$?

- a. $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in.}} \cdot \frac{100 \text{ cm}}{1 \text{ m}} \cdot \frac{3600 \text{ s}}{1 \text{ h}}$
- b. $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{1 \text{ in.}}{2.54 \text{ cm}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{1 \text{ h}}{3600 \text{ s}}$
- c. $\frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in.}}{1 \text{ ft}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in.}} \cdot \frac{100 \text{ cm}}{1 \text{ m}} \cdot \frac{1 \text{ h}}{3600 \text{ s}}$
- d. $\frac{1 \text{ mi}}{5280 \text{ ft}} \cdot \frac{1 \text{ ft}}{12 \text{ in.}} \cdot \frac{1 \text{ in.}}{2.54 \text{ cm}} \cdot \frac{100 \text{ cm}}{1 \text{ m}} \cdot \frac{3600 \text{ s}}{1 \text{ h}}$
- e. $\frac{1 \text{ mi}}{5280 \text{ ft}} \cdot \frac{1 \text{ ft}}{12 \text{ in.}} \cdot \frac{1 \text{ in.}}{2.54 \text{ cm}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{3600 \text{ s}}{1 \text{ h}}$

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

10. One U.S. fluid gallon contains a volume of 231 cubic inches. How many liters of gasoline would you have to buy in Canada to fill a 14-gallon tank? (Note: $1 \text{ L} = 10^3 \text{ cm}^3$.)

- a. 53
- b. 21
- c. 14
- d. 8.0
- e. 4.0

Chapter 1 - Getting Started.

ANSWER:

a

POINTS:

3

DIFFICULTY:

Challenging

11. John and Linda are arguing about the definition of density. John says the density of an object is proportional to its mass. Linda says the object's mass is proportional to its density and to its volume. Which one, if either, is correct?

- They are both wrong.
- John is correct, but Linda is wrong.
- John is wrong, but Linda is correct.
- They are both correct.
- They are free to redefine density as they wish.

ANSWER:

d

POINTS:

1

DIFFICULTY:

Easy

12. Spike claims that dimensional analysis shows that the correct expression for change in velocity, $\vec{v}_f - \vec{v}_i$, is

$\vec{v}_f - \vec{v}_i = \frac{mt}{F}$, where m is mass, t is time, and F is the magnitude of force. Carla says that can't be true because the

dimensions of force are $\left[\left[\frac{ML}{T^2} \right] \right]$. Which one, if either, is correct?

- $[[\vec{v}]] = \left[\left[\frac{ML}{T} \right] \right]$
Spike, because

- $[[\vec{v}]] = \left[\left[\frac{T^2}{L} \right] \right]$
Spike, because

- $[[\vec{v}]] = \left[\left[\frac{L}{T} \right] \right]$
Carla, because

- $[[\vec{v}]] = \left[\left[\frac{L}{MT} \right] \right]$
Carla, because

- $[[\vec{F}]] = \left[\left[\frac{T^2}{ML} \right] \right]$
Spike, because the dimensions of force are

ANSWER:

c

POINTS:

2

DIFFICULTY:

Average

13. Which one of the quantities below has dimensions equal to $\left[\left[\frac{ML}{T^2} \right] \right]$?

- mv
- mv^2

Chapter 1 - Getting Started.

- c. $\frac{mv^2}{r}$
- d. mrv
- e. $\frac{mv^2}{r^2}$

ANSWER:

c

POINTS:

2

DIFFICULTY:

Average

14. The standard exam page is 8.50 inches by 11.0 inches. Its area in cm^2 is

- a. 19.5.
- b. 36.8.
- c. 93.5.
- d. 237.
- e. 603.

ANSWER:

e

POINTS:

2

DIFFICULTY:

Average

15. A standard exam page is 8.5 inches by 11 inches. An exam that is 2.0 mm thick has a volume of

- a. $1.9 \times 10^4 \text{ mm}^3$.
- b. $4.7 \times 10^4 \text{ mm}^3$.
- c. $1.2 \times 10^5 \text{ mm}^3$.
- d. $3.1 \times 10^5 \text{ mm}^3$.
- e. $3.1 \times 10^3 \text{ mm}^3$.

ANSWER:

c

POINTS:

3

DIFFICULTY:

Challenging

16. Which quantity can be converted from the English system to the metric system by the conversion

$$\text{factor} = \frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{12 \text{ in}}{1 \text{ ft}} \cdot \frac{2.54 \text{ cm}}{1 \text{ in.}} \cdot \frac{1 \text{ m}}{100 \text{ cm}} \cdot \frac{1 \text{ h}}{3600 \text{ s}} ?$$

- a. feet per second
- b. feet per hour
- c. miles per second
- d. miles per hour
- e. miles per minute

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

Chapter 1 - Getting Started.

17. The answer to a question is $[[\text{MLT}^{-1}]]$. The question is "What are the dimensions of

- a. $mr?$ "
- b. $mvr?$ "
- c. $ma?$ "
- d. $mat?$ "
- e. $\frac{mv^2}{r}?$ "

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

18. If each frame of a motion picture film is 35 cm high, and 24 frames go by in a second, estimate how many frames are needed to show a two-hour-long movie.

- a. 1400
- b. 25,000
- c. 50,000
- d. 170,000
- e. This cannot be determined without knowing how many reels were used.

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

19. One number has three significant figures and another number has four significant figures. If these numbers are added, subtracted, multiplied, or divided, which operation can produce the greatest number of significant figures?

- a. the addition
- b. the subtraction
- c. the multiplication
- d. the division
- e. All the operations result in the same number of significant figures.

ANSWER:

a

POINTS:

2

DIFFICULTY:

Average

20. A rectangle has a length of 1.323 m and a width of 4.16 m. Using significant figure rules, what is the area of this rectangle?

- a. 5.50368 m^2
- b. 5.5037 m^2
- c. 5.504 m^2
- d. 5.50 m^2
- e. 5.5 m^2

ANSWER:

d

POINTS:

2

DIFFICULTY:

Average

Chapter 1 - Getting Started.

21. The standard kilogram is a platinum-iridium cylinder 39 mm in height and 39 mm in diameter. What is the density of the material?

ANSWER: 21475 kg/m³

POINTS: 2

DIFFICULTY: Average

22. A 2.00 m by 3.00 m plate of aluminum has a mass of 324 kg. What is the thickness of the plate? (The density of aluminum is 2.70×10^3 kg/m³.)

ANSWER: 2.00 cm

POINTS: 2

DIFFICULTY: Average

23. What is the mass of air in a room that measures 5.0 m $\times$ 8.0 m $\times$ 3.0 m? (The density of air is 1/800 that of water).

ANSWER: 150 kg

POINTS: 2

DIFFICULTY: Average

24. The basic function of a carburetor of an automobile is to atomize the gasoline and mix it with air to promote rapid combustion. As an example, assume that 30 cm³ of gasoline is atomized into N spherical droplets, each with a radius of 2.0×10^{-5} m. What is the total surface area of these N spherical droplets?

ANSWER: 4.5 m²

POINTS: 3

DIFFICULTY: Challenging